

NYSTATIN- nystatin cream
E. Fougera & Co. a division of Fougera Pharmaceuticals Inc.

NYSTATIN CREAM USP


FOR TOPICAL USE ONLY.

Rx only

NOT FOR OPHTHALMIC USE.

DESCRIPTION:

Nystatin is a polyene antifungal antibiotic obtained from *Streptomyces noursei*. Structural formula:



Molecular Weight 926.13

Molecular Formula C₄₇H₇₅NO₁₇

Nystatin cream is for dermatologic use.

Nystatin cream for topical use, contains 100,000 USP nystatin units per gram in an aqueous, vanishing cream base containing Polysorbate 60, Aluminum Hydroxide Compressed Gel, Titanium Dioxide, Glyceryl Monostearate, Polyethylene Glycol 400 Monostearate, Simethicone Emulsion, Sorbic Acid, Propylene Glycol, Ethylenediamine, Polyoxyethylene Fatty Alcohol Ether, Sorbitol Solution, Methylparaben, Propylparaben, Hydrochloric Acid, White Petrolatum, and Purified Water.

CLINICAL PHARMACOLOGY:

Pharmacokinetics: Nystatin is not absorbed from intact skin or mucous membrane.

Microbiology: Nystatin is an antifungal antibiotic which is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi, including *Candida albicans*, *C. parapsilosis*, *C. tropicalis*, *C. guilliermondi*, *C. pseudotropicalis*, *C. krusei*, *Torulopsis glabrata*, *Tricophyton rubrum*, *T. mentagrophytes*. Nystatin acts by binding to sterols in the cell membrane of susceptible species resulting in a change in membrane permeability and the subsequent leakage of intracellular components. On repeated subculturing with increasing levels of nystatin, *Candida albicans* does not develop resistance to nystatin. Generally, resistance to nystatin does not develop during therapy. However, other species of *Candida* (*C. tropicalis*, *C. guilliermondi*, *C. krusei*, and *C. stellatoidea*) become quite resistant on treatment with nystatin and simultaneously become cross resistant to amphotericin as well. This resistance is lost when the antibiotic is removed. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

INDICATIONS AND USAGE:

Nystatin cream is indicated in the treatment of cutaneous or mucocutaneous mycotic infections caused by *Candida albicans* and other susceptible *Candida* species.

This cream is not indicated for systemic, oral, intravaginal or ophthalmic use.

CONTRAINDICATIONS:

Nystatin cream is contraindicated in patients with a history of hypersensitivity to any of their components.

PRECAUTIONS:

General: Nystatin cream should not be used for the treatment of systemic, oral, intravaginal or ophthalmic infections.

If irritation or sensitization develops, treatment should be discontinued and appropriate measures taken as indicated. It is recommended that KOH smears, cultures, or other diagnostic methods be used to confirm the diagnosis of cutaneous or mucocutaneous candidiasis and to rule out infection caused by other pathogens.

INFORMATION FOR THE PATIENT: Patients using this medication should receive the following information and instructions:

1. The patient should be instructed to use this medication as directed (including the replacement of missed doses). This medication is not for any disorder other than that for which it is prescribed.
2. Even if symptomatic relief occurs within the first few days of treatment, the patient should be advised not to interrupt or discontinue therapy until the prescribed course of treatment is completed.
3. If symptoms of irritation develop, the patient should be advised to notify the physician promptly.

Laboratory Tests: If there is a lack of therapeutic response, KOH smears, cultures, or other diagnostic methods should be repeated.

Carcinogenesis, Mutagenesis, Impairment of Fertility: No long-term animal studies have been performed to evaluate the carcinogenic potential of nystatin. No studies have been performed to determine the mutagenicity of nystatin or the effects on male or female fertility.

Pregnancy: Teratogenic Effects: Category C: Animal reproduction studies have not been conducted with any nystatin cream. It also is not known whether this cream can cause fetal harm when used by a pregnant woman or can affect reproductive capacity. Nystatin cream should be prescribed for a pregnant woman only if the potential benefit to the mother outweighs the potential risk to the fetus.

Nursing Mothers: It is not known whether nystatin is excreted in human milk. Caution should be exercised when nystatin is prescribed for a nursing woman.

Pediatric Use: Safety and effectiveness have been established in the pediatric population from birth to 16 years. (See DOSAGE AND ADMINISTRATION.)

Geriatric Use: Clinical studies with nystatin cream did not include sufficient numbers of subjects aged 65 years and older to determine whether they respond differently than younger subjects. Other reported clinical experience has not identified differences in responses between elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.]

ADVERSE REACTIONS:

The frequency of adverse events reported in patients using nystatin cream is less than 0.1%. The more common events that were reported include allergic reactions, burning, itching, rash, eczema, and pain on application. (See PRECAUTIONS: General.)

DOSAGE AND ADMINISTRATION:

Adults and Pediatric Patients (Neonates and Older): Apply liberally to affected area twice daily or as indicated until healing is complete.

HOW SUPPLIED:

Nystatin Cream USP is available as follows:

NDC 0168-0054-15 15 gram tube

NDC 0168-0054-30 30 gram tube

Each tube provides 100,000 USP Nystatin Units per gram.

Store at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature].

Avoid freezing.

E. FOUGERA & CO.

A division of

Fougera

PHARMACEUTICALS INC.

Melville, New York 11747

I25415G

R07/13

#61

PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – 15G CONTAINER

NDC 0168-0054-15

FOUGERA®

NYSTATIN CREAM USP

100,000 USP Nystatin Units Per Gram

FOR EXTERNAL USE ONLY.

NOT FOR OPHTHALMIC USE.

Rx only

NET WT 15 grams

<p>NDC 0168-0054-15</p> <p>fougera®</p> <p>NYSTATIN CREAM USP 100,000 USP Nystatin Units Per Gram</p> <p>FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.</p> <p>USUAL DOSAGE: Apply liberally to affected area twice daily. See insert for full prescribing information. WARNING: Keep this and all drugs out of reach of children. TO OPEN: Use cap to puncture seal. IMPORTANT: Do not use if seal has been punctured or is not visible.</p> <p>E. FOUGERA & CO. A division of Fougera Pharmaceuticals Inc. Melville, New York 11747</p>	<p>Rx only</p> <p>Each gram contains: 100,000 USP Nystatin Units, Polysorbate 60, Aluminum Hydroxide Compressed Gel, Titanium Dioxide, Glyceryl Monostearate, Polyethylene Glycol 400 Monostearate, Simethicone Emulsion, Sorbic Acid, Propylene Glycol, Ethylenediamine, Polyoxyethylene Fatty Alcohol Ether, Sorbitol Solution, Methylparaben, Propylparaben, Hydrochloric Acid, White Petrolatum, and Purified Water.</p> <p>NET WT 15 grams</p> <p>Store at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature]. Avoid freezing.</p> <p>See crimp of tube for Control Number and Expiration Date. U4424C R07/13</p> 	<p>USE</p> 	
---	---	--	--

PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – 15 G CARTON

NDC 0168-0054-15

Rx only

FOUGERA®

NYSTATIN CREAM USP
100,000 USP Nystatin Units Per Gram

WARNING: Keep this and all drugs out of reach of children.

FOR EXTERNAL USE ONLY.
NOT FOR OPHTHALMIC USE.

NET WT 15 grams



NYSTATIN

nystatin cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:0168-0054
Route of Administration	TOPICAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
nystatin (nystatin)	nystatin	100000 [USP'U] in 1 g

Inactive Ingredients

Ingredient Name	Strength
Polysorbate 60	
Algeldrate	
Titanium Dioxide	
Glyceryl Monostearate	
PEG-8 Stearate	
Silicon Dioxide	
Sorbic Acid	
Propylene Glycol	
Ethylenediamine	
Sorbitol	
Methylparaben	

Propylparaben	
Hydrochloric Acid	
Petrolatum	
Water	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0168-0054-15	1 in 1 CARTON		
1		15 g in 1 TUBE		
2	NDC:0168-0054-30	1 in 1 CARTON		
2		30 g in 1 TUBE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA062129	02/01/1979	

Labeler - E. Fougera & Co. a division of Fougera Pharmaceuticals Inc. (043838424)

Revised: 5/2014

E. Fougera & Co. a division of Fougera Pharmaceuticals Inc.