

**LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride tablet**  
**Amerisource Bergen**

-----  
**Drug Facts**

**ACTIVE INGREDIENT (IN EACH CAPLET)**

Loperamide HCl USP, 2 mg

**PURPOSE**

Anti-diarrheal

**USE**

Controls symptoms of diarrhea, including Travelers' Diarrhea

**WARNINGS**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Do not use**

If you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease

**Ask a doctor or pharmacist before use if you are**

Taking antibiotics

**When using this product**

- tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- diarrhea lasts for more than 2 days
- symptoms get worse
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding**

Ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**DIRECTIONS**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

#### OTHER INFORMATION

- store between 20° – 25° C (68° – 77° F)
- **TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.**

#### INACTIVE INGREDIENTS

Anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no.10, FD&C blue no.1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

#### QUESTIONS?

Call **1800-406-7984**

#### PRINCIPAL DISPLAY PANEL

**GOOD NEIGHBOR PHARMACY®**

**NDC 46122-0169-08**

**\*Compare to Imodium® A-D active ingredient**

**Loperamide HCl Tablets USP, 2 mg**

**Anti-Diarrheal**

**Controls the Symptoms of Diarrhea**

**72 caplets ■**

**Each caplet (■ capsule-shaped tablet) contains Loperamide HCl USP, 2 mg**

**Distributed By: AmerisourceBergen**

**5098330/R0912**

TOP LABEL



INSIDE TOP LABEL

<p><b>Drug Facts (continued)</b></p> <p>Ask a doctor or pharmacist before use if you are taking antibiotics</p> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>■ tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ diarrhea lasts for more than 2 days</li> <li>■ symptoms get worse</li> <li>■ you get abdominal swelling or bulging. These may be signs of a serious condition.</li> </ul> <p>if pregnant or breast-feeding, ask a health professional before use.</p>	<p><b>Drug Facts (continued)</b></p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea</li> <li>■ find right dose on chart. If possible, use weight to dose; otherwise, use age.</li> </ul> <table border="1"> <tr> <td>adults and children 12 years and over</td> <td>2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours</td> </tr> </table>	adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours		

BASE LABEL

<p><b>Drug Facts (continued)</b></p> <table border="1"> <tr> <td>children 9-11 years (60-95 lbs)</td> <td>1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours</td> </tr> <tr> <td>children 6-8 years (49-59 lbs)</td> <td>1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours</td> </tr> <tr> <td>children under 6 years (up to 47 lbs)</td> <td>ask a doctor</td> </tr> </table>	children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours	children 6-8 years (49-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours	children under 6 years (up to 47 lbs)	ask a doctor	<p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 20° - 25° C (68° - 77° F)</li> <li>■ TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.</li> </ul> <p><b>Inactive ingredients:</b> anhydrous lactose, croscarmellose sodium, crospovidone, D&amp;C yellow no. 10, FD&amp;C blue no. 1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch</p> <p><b>Questions?</b> call 1-800-406-7984</p>
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours						
children 6-8 years (49-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours						
children under 6 years (up to 47 lbs)	ask a doctor						

\*This product is not manufactured or distributed by McNeil-PPC, Inc., distributor of Imodium® A-D. Imodium® is a registered trademark of Johnson & Johnson.

Distributed By: AmerisourceBergen  
1330 Morris Drive, Chesterbrook, PA 19387  
Visit us at [www.goodneighborpharmacy.com](http://www.goodneighborpharmacy.com)

# LOPERAMIDE HYDROCHLORIDE

loperamide hydrochloride tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:46 122-169
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77T135393C) (LOPERAMIDE - UNII:6X9OC3H4I)	LOPERAMIDE HYDROCHLORIDE	2 mg

## Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 68401960MK)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
TRISTEARIN (UNII: P6OCJ2551R)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	

## Product Characteristics

<b>Color</b>	green	<b>Score</b>	2 pieces
<b>Shape</b>	CAPSULE	<b>Size</b>	9mm

<b>Flavor</b>		<b>Imprint Code</b>	123
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46 122-169-08	72 in 1 BOTTLE		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074091	02/01/1993	

**Labeler** - Amerisource Bergen (007914906)

**Registrant** - Ranbaxy Pharmaceuticals Inc. (937890044)

**Establishment**

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		051565745	manufacture(46 122-169)