

# PENTAM 300- pentamidine isethionate injection, powder, lyophilized, for solution


Fresenius Kabi USA, LLC

## Pentam® 300 (pentamidine isethionate for injection)

### For Injection Only

### DESCRIPTION

Pentam 300 (pentamidine isethionate for injection), an anti-protozoal agent, is a sterile, nonpyrogenic, lyophilized product. After reconstitution, it should be administered by intramuscular (IM) or intravenous (IV) routes (see **DOSAGE AND ADMINISTRATION**). Pentamidine isethionate is a white crystalline powder soluble in water and glycerin, slightly soluble in alcohol and insoluble in ether, acetone, and chloroform. It is chemically designated as 4,4-[1,5-pentanediy]bis(oxy)]bis-benzenecarboximidamid with the following structural formula:



### Each vial contains:

Pentamidine isethionate . . . . . 300 mg

### CLINICAL PHARMACOLOGY

Pentamidine isethionate, an aromatic diamidine, is known to have activity against *Pneumocystis carinii*. The mode of action of pentamidine is not fully understood. *In vitro* studies indicate that the drug interferes with protozoal nuclear metabolism by inhibition of DNA, RNA, phospholipid and protein synthesis.

### Pharmacokinetics

Pharmacokinetic parameters following the administration of 4 mg/kg pentamidine isethionate as a single two-hour intravenous infusion or after a single intramuscular injection to 12 patients with AIDS are presented in the following table:

Mean ± SD	Cmax ng/mL	Clearance L/h	Half-life hours	Vdss L	Concentration ng/mL	
					8 hour	24 hour
2 hour I.V. infusion 4 mg/kg (N=6)	612 ± 371	248 ± 91	6.4 ± 1.3	821 ± 535	19.3 ± 16.9	2.9 ± 1.4
I.M. 4 mg/kg (N=6)	209 ± 48	305 ± 81	9.4 ± 2.0	2724 ± 1066	22.9 ± 8.0	6.6 ± 3.5

In seven patients treated with daily IM doses of pentamidine at 4 mg/kg for 10 to 12 days, plasma concentrations were between 300 to 500 ng/mL. The concentrations did not appreciably change with time after injection or from day to day. Higher plasma concentrations were encountered in patients with an elevated blood urea nitrogen. The patients continued to excrete decreasing amounts of pentamidine in urine up to 6 to 8 weeks after cessation of the treatment.

Following multiple intravenous administration of pentamidine isethionate (3.7 to 4 mg/kg/day infused over 4 hours) to 6 patients with AIDS being treated for PCP, the pharmacokinetic parameters obtained on Days 1,4 and 7 are summarized in the following table:

Mean ± SD	Cmax* ng/mL	Cmin* ng/mL	Clearance mL/min	Renal Clearance mL/min/1.73 m <sup>2</sup>	Creatinine Clearance mL/min/1.73 m <sup>2</sup>
Day 1	175.3 ± 54	---	5737 ± 1878	269 ± 149	97 ± 12
Day 4	210.9 ± 80	17.6 ± 9.5	3350 ± 1944	214 ± 145	93 ± 17
Day 7	256.7 ± 89	40.8 ± 16.1	1989 ± 566	134 ± 60	69 ± 17

\*derived from Lidman

Compared to the mean AUC on Day 1, AUC on Day 4 and Day 7 were about 2 and 3 fold higher, respectively, suggesting that steady state was not achieved by Day 7 of dosing.

In other published reports of pharmacokinetics of pentamidine following daily intravenous doses of 2 to 4 mg/kg/day, clearance ranged from 30 to 40 mL/min/kg and V<sub>dss</sub> ranged from 200 to 400 L/kg. Reported values for terminal half-lives of 2.8 to 12 days is suggestive of a deep peripheral compartment. In the urine, up to 12% of the administered dose has been recovered during a dosing interval as unchanged pentamidine.

Tissue distribution was studied in mice given a single intraperitoneal injection of pentamidine at 10 mg/kg. The concentration was highest in the kidneys followed by the liver. In mice, pentamidine was excreted unchanged, primarily via the kidneys with some elimination in the feces. The ratio of amounts excreted in the urine and feces (4:1) was constant over the period of study.

Tissue distribution has also been studied in normal and in renally impaired dogs (N = 3 each) given 13 mg/kg of pentamidine IV, in 2 doses separated by five weeks. The concentration of pentamidine was highest in the liver followed by kidneys and lungs. Pentamidine was concentrated in these organs approximately 70 to 1000 times that of the peak serum concentration. Similar findings were reported in normal and in renally impaired dogs (N = 2 each) given 97.5 mg/kg of pentamidine IV, in 15 daily doses. After repeated doses, the organs showed a further 3 to 7 fold accumulation while serum concentrations remained unchanged.

## INDICATIONS AND USAGE

Pentam 300 (pentamidine isethionate for injection) is indicated for the treatment of pneumonia due to *Pneumocystis carinii*.

## CONTRAINDICATIONS

Contraindicated in patients with a history of hypersensitivity to pentamidine isethionate.

## WARNINGS

Fatalities due to severe hypotension, hypoglycemia, acute pancreatitis and cardiac arrhythmias have been reported in patients treated with pentamidine isethionate, both by the IM and IV routes. Severe hypotension may result after a single IM or IV dose and is more likely with rapid IV administration (see **PRECAUTIONS**). The administration of the drug should, therefore, be limited to the patients in whom *Pneumocystis carinii* has been demonstrated. Patients should be closely monitored for the development of serious adverse reactions (see **PRECAUTIONS** and **ADVERSE REACTIONS**).

**Extravasations have been reported which, in some instances, proceeded to ulceration, tissue necrosis and/or sloughing at the injection site. While not common, surgical debridement and skin grafting has been necessary in some of these cases; long-term sequelae have been reported. Prevention is the most effective means of limiting the severity of extravasation. The intravenous needle or catheter must be properly positioned and closely observed throughout the period of pentamidine isethionate administration. If extravasation occurs, the injection should be discontinued immediately and restarted in another vein. Because there are no known local treatment measures which have proven to be useful, management of the extravasation should be symptomatic.**

## PRECAUTIONS

### General

Pentamidine isethionate should be used with caution in patients with hypertension, hypotension, ventricular tachycardia, hypoglycemia, hyperglycemia, hypocalcemia, pancreatitis, leukopenia, thrombocytopenia, anemia, hepatic or renal dysfunction and Stevens-Johnson syndrome.

**Patients may develop sudden, severe hypotension after a single dose of pentamidine isethionate, whether given IV or IM. Therefore, patients receiving the drug should be lying down and the blood pressure should be monitored closely during administration of the drug and several times thereafter until the blood pressure is stable. Equipment for emergency resuscitation should be readily available. If pentamidine isethionate is administered IV, it should be infused over a period of 60 to 120 minutes.**

Pentamidine isethionate-induced hypoglycemia has been associated with pancreatic islet cell necrosis and inappropriately high plasma insulin concentrations. Hyperglycemia and diabetes mellitus, with or without preceding hypoglycemia, have also occurred, sometimes several months after therapy with pentamidine isethionate. Therefore, blood glucose levels should be monitored daily during therapy with pentamidine isethionate, and several times thereafter.

### **Renal and Hepatic Impairment**

The efficacy or safety of alternative Pentam 300 dosing protocols have not been established for patients with impaired renal or hepatic function.

### **Laboratory Tests**

The following tests should be carried out before, during and after therapy:

- a) Daily blood urea nitrogen and serum creatinine determinations.
- b) Daily blood glucose determinations.
- c) Complete blood count and platelet count.
- d) Liver function test, including serum bilirubin, alkaline phosphatase, AST (SGOT), and ALT (SGPT).
- e) Serum calcium determinations.
- f) Electrocardiograms.

### **Drug Interactions**

No drug interaction studies with Pentam 300 have been conducted.

Because the nephrotoxic effects may be additive, the concomitant or sequential use of pentamidine isethionate and other nephrotoxic drugs such as aminoglycosides, amphotericin B, cisplatin, foscarnet, or vancomycin should be closely monitored and avoided, if possible.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

No studies have been conducted to evaluate the potential of pentamidine isethionate as a carcinogen, mutagen, or cause of impaired fertility.

### **Pregnancy-Pregnancy Category C**

Animal reproduction studies have not been conducted with pentamidine isethionate. It is also not known whether pentamidine isethionate can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Pentamidine isethionate should not be given to a pregnant woman unless the potential benefits are judged to outweigh the unknown risks.

### **Nursing Mothers**

It is not known whether pentamidine isethionate is excreted in human milk. Because of the potential for serious adverse reactions in nursing infants from pentamidine isethionate, a decision should be made whether to discontinue nursing or to discontinue

the drug, taking into account the importance of the drug to the mother. Because many drugs are excreted in human milk, pentamidine isethionate should not be given to a nursing mother unless the potential benefits are judged to outweigh the unknown risks.

## Pediatric Use

Intravenous and intramuscular pentamidine has been described as an effective treatment for *Pneumocystis carinii* pneumonia (PCP) in immunocompromised pediatric patients beyond 4 months of age. The efficacy and safety profiles in these pediatric patients were similar to those observed in adult patients (See **DOSAGE AND ADMINISTRATION** and **OVERDOSAGE**).

## ADVERSE REACTIONS

To report **SUSPECTED ADVERSE REACTIONS**, contact **Fresenius Kabi USA, LLC** at **1-800-551-7176** or **FDA** at **1-800-FDA-1088** or **www.fda.gov/medwatch**.

**CAUTION:** Fatalities due to severe hypotension, hypoglycemia, acute pancreatitis and cardiac arrhythmias have been reported in patients treated with pentamidine isethionate, both by the IM and IV routes. Nephrotoxic events (increased creatinine, impaired renal function, azotemia, and renal failure) are common with the parenteral administration of pentamidine isethionate. The administration of the drug should, therefore, be limited to the patients in whom *Pneumocystis carinii* has been demonstrated.

The most frequently reported spontaneous adverse events (1 to 30%) reported in clinical trials, regardless of their relation to pentamidine isethionate therapy were as follows (n=424):

Cardiovascular: Hypotension	5.0%
Gastrointestinal: Anorexia/Nausea	5.9%
Hematologic: Anemia Leukopenia Thrombocytopenia	1.2% 10.4% 2.6%
Hepatic: Elevated liver function tests	8.7%
Metabolic: Hypoglycemia	5.9%
Neurologic: Confusion/hallucinations	1.7%
Skin: Sterile abscess and/or necrosis, pain, or induration at the site of IM injection Rash	11.1% 3.3%
Special Senses: Bad taste	1.7%
Urogenital:	

Azotemia	8.5%
Elevated serum creatinine	23.6%
Elevated blood urea nitrogen	6.6%
Impaired renal function	28.8%

Adverse events with a frequency of less than 1% incidence were as follows (No causal relationship to treatment has been established for these adverse events):

Body as a whole:	Allergic reaction (i.e. urticaria, itching, rash), anaphylaxis, arthralgia, chills, extrapulmonary pneumocystosis, headache, night sweats, and Stevens-Johnson syndrome.
Cardiovascular:	Abnormal ST segment of electrocardiogram, cardiac arrhythmias, cerebrovascular accident, hypertension, palpitations, phlebitis, syncope, tachycardia, vasodilatation, vasculitis and ventricular tachycardia.
Gastrointestinal:	Abdominal pain, diarrhea, dry mouth, dyspepsia, hematochezia, hypersalivation, melena, pancreatitis, splenomegaly, and vomiting.
Hematological:	Defibrination, eosinophilia, neutropenia, pancytopenia, and prolonged clotting time.
Hepatic:	Hepatic dysfunction, hepatitis and hepatomegaly
Metabolic:	Hyperglycemia, hyperkalemia, hypocalcemia, and hypomagnesemia.
Neurological:	Anxiety, confusion, depression, dizziness, drowsiness, emotional lability, hypesthesia, insomnia, memory loss, neuropathy, nervousness, neuralgia, paranoia, paresthesia, peripheral neuropathy, seizure, tremors, unsteady gait, and vertigo.
Respiratory system:	Asthma, bronchitis, bronchospasm, chest congestion, chest tightness, coryza, cyanosis, eosinophilic or interstitial pneumonitis, gagging, hemoptysis, hyperventilation, laryngitis, laryngospasm, non-specific lung disorder, nasal congestion, pleuritis, pneumothorax, rales, rhinitis, shortness of breath, and tachypnea.
Skin:	Desquamation, dry and breaking hair, dry skin, erythema, dermatitis, pruritus, rash, and urticaria.
Special senses:	Blepharitis, blurred vision, conjunctivitis, contact lens discomfort, eye pain or discomfort, loss of hearing, loss of taste, and loss of smell.
Urogenital:	Flank pain, hematuria, incontinence, nephritis, renal dysfunction and renal failure.

From post-marketing clinical experience with pentamidine isethionate, the following adverse events have been reported: cough, diabetes mellitus/ketoacidosis, dyspnea, infiltration (extravasation—see **WARNINGS**), and torsades de pointes.

## OVERDOSAGE

A 17 month old infant inadvertently received 1600 mg of intravenous pentamidine isethionate which was followed by renal and hepatic function impairment, hypotension and cardiopulmonary arrest. Treatment included cardiopulmonary resuscitation, epinephrine, atropine and intubation. In addition, a four hour course of charcoal

hemoperfusion was accompanied by reduction of pentamidine serum concentration and stabilization of the patient's condition. The patient recovered from these adverse events, but later died due to an unknown cause. <sup>1</sup>

## **DOSAGE AND ADMINISTRATION**

**CAUTION: DO NOT USE SODIUM CHLORIDE INJECTION, USP FOR INITIAL RECONSTITUTION BECAUSE PRECIPITATION WILL OCCUR.** Pentamidine isethionate should be administered IM or IV only. The recommended regimen for adults and pediatric patients beyond 4 months of age is 4 mg/kg once a day for 14 to 21 days. Therapy for longer than 21 days with pentamidine isethionate has also been used but may be associated with increased toxicity.

### **Intramuscular Injection**

The contents of one vial (300 mg) should be dissolved in 3 mL of Sterile Water for Injection, USP at 22° - 30°C (72° - 86°F). The calculated daily dose should then be withdrawn and administered by deep IM injection.

### **Intravenous Injection**

The contents of one vial (300 mg) should first be dissolved in 3 to 5 mL of Sterile Water for Injection, USP, or 5% Dextrose Injection, USP at 22° - 30°C (72° - 86°F). The calculated dose of pentamidine isethionate should then be withdrawn and diluted further in 50 to 250 mL of 5% Dextrose Injection, USP.

**The diluted IV solutions containing pentamidine isethionate should be infused over a period of 60 to 120 minutes.**

Aseptic technique should be employed in preparation of all solutions. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

### **Stability**

After reconstitution with sterile water, the Pentam solution is stable for 48 hours in the original vial at room temperature if protected from light. To avoid crystallization, store at 22° - 30°C (72° - 86°F). Intravenous infusion solutions of pentamidine isethionate at 1 mg/mL and 2.5 mg/mL prepared in 5% Dextrose Injection, USP are stable at room temperature for up to 24 hours.

Intravenous (IV) solutions of pentamidine isethionate have been shown to be incompatible with fluconazole and foscarnet sodium. IV solutions of pentamidine isethionate have been shown to be compatible with IV solutions of zidovudine (AZT) and diltiazem hydrochloride.

## **HOW SUPPLIED**

PENTAM<sup>®</sup> 300 (pentamidine isethionate for injection)

<b>Product Code</b>	<b>Unit of Sale</b>	<b>Strength</b>	<b>Each</b>
---------------------	---------------------	-----------------	-------------

11310	NDC 63323-113-10 Unit of 10	300 mg	NDC 63323-113-01 Lyophilized Product in Single Dose Vial
-------	--------------------------------	--------	--

Store dry product at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from light.

Preservative Free. Discard unused portion.

## REFERENCE

1. Watts RG; Conte JE, Jr.; Zurlinden E; Waldo FB: Effect of charcoal hemoperfusion on clearance of pentamidine isethionate after accidental overdose. J Toxicol Clin Toxicol 1997;35:89-92.



[www.fresenius-kabi.com/us](http://www.fresenius-kabi.com/us)

45851D

Revised: May 2022

## **PACKAGE LABEL - PRINCIPAL DISPLAY - Pentam<sup>®</sup> 300 mg Single Dose Vial Label**

NDC 63323-113-01      11310

### **PENTAM<sup>®</sup> 300**

*(pentamidine isethionate for injection)*

300 mg

Lyophilized

### **For IM or IV Use**

Single Dose Vial

Rx only

**Preservative Free**

Sterile, Nonpyrogenic  
Each vial contains: Pentamidine Isethionate 300 mg.

**Usual Dosage:** See Package Insert.  
Reconstitute vial contents with Sterile Water for Injection or 5% Dextrose Injection only as directed in Package Insert.

Store dry product at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect dry product and reconstituted solution from light.

**Fresenius Kabi**

Lake Zurich, IL 60047  
www.fresenius-kabi.com/us

NDC 63323-113-01 11310

**PENTAM<sup>®</sup> 300**  
*(pentamidine isethionate for injection)*

**300 mg**  
Lyophilized

**For IM or IV Use**

Single Dose Vial

Rx only



3 63323-113-01 8

401894E

LOT/EXP

**PACKAGE LABEL - PRINCIPAL DISPLAY - Pentam<sup>®</sup> 300 mg Single Dose Tray Label**

NDC 63323-113-10 11310

**PENTAM<sup>®</sup> 300**

*(pentamidine isethionate for injection)*

**300 mg**

Lyophilized

**For Intramuscular or Intravenous Use**

Rx only

10 Single Dose Vials

NDC 63323-113-10 11310

**PENTAM<sup>®</sup> 300**  
*(pentamidine isethionate for injection)*

**300 mg**

Lyophilized

**For Intramuscular or Intravenous Use**

Rx only

10 Single Dose Vials

**Preservative Free**

Sterile, Nonpyrogenic

Each vial contains: Pentamidine Isethionate 300 mg.

**Dosage:** Recommended regimen for adults and pediatric patients is 4 mg/kg once a day for 14 days. The benefits and risks of therapy with pentamidine isethionate for more than 14 days are not well defined.

**ADMINISTRATION: Intramuscular Injection:** The contents of 1 vial (300 mg) should be dissolved in 3 mL of Sterile Water for Injection, USP. The calculated daily dose should then be withdrawn and administered by deep intramuscular injection.

**Intravenous Injection:** The contents of 1 vial should first be dissolved in 3-5 mL of Sterile Water for Injection, USP or 5% Dextrose Injection, USP only. The calculated dose of pentamidine isethionate should then be

withdrawn and diluted further in 50-250 mL of 5% dextrose solution. **The diluted intravenous solutions containing pentamidine isethionate should be infused over a period of 60 minutes.**

For additional information: See Package Insert.

Store dry product at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect dry product and reconstituted solution from light.

**FRESENIUS KABI**  
Lake Zurich, IL 60047  
www.fresenius-kabi.com/us



42956A

(01)20363323113104

**PENTAM 300**

pentamidine isethionate injection, powder, lyophilized, for solution

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:63323-113
<b>Route of Administration</b>	INTRAMUSCULAR, INTRAVENOUS		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>PENTAMIDINE ISETHIONATE</b> (UNII: V2P3K60DA2) (PENTAMIDINE - UNII:673LC5J4LQ)	PENTAMIDINE ISETHIONATE	300 mg in 3 mL

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:63323-113-10	10 in 1 TRAY	01/16/2001	
1	NDC:63323-113-01	3 mL in 1 VIAL; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA	NDA019264	01/16/2001	

**Labeler** - Fresenius Kabi USA, LLC (608775388)**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Fresenius Kabi USA, LLC		023648251	manufacture(63323-113)

Revised: 9/2022

Fresenius Kabi USA, LLC