

**GOOD SENSE ACID REDUCER- famotidine tablet, film coated**  
**L. Perrigo Company**

-----  
**Perrigo Acid Reducer Drug Facts**

**Active ingredient (in each tablet)**

Famotidine 10 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

**Inactive ingredients**

carnauba wax, hypromellose, iron oxide red, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, talc, titanium dioxide, triacetin

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

Original Strength

See New Warnings

Famotidine Tablets, 10mg

Acid Reducer

Actual Size

Just One Tablet!

-  
Prevents & Relieves Heartburn Due to Acid Indigestion

Compare to active ingredient of Pepcid® AC

100% SATISFACTION GUARANTEED

30 Tablets

GOODSENSE®

Famotidine Tablets, 10 mg

**Acid Reducer**

DO NOT USE IF BLISTER  
UNIT IS BROKEN OR TORN

JUST ONE TABLET prevents  
and relieves heartburn due to  
acid indigestion brought on by  
eating and drinking certain  
foods and beverages.

GOODSENSE®

NDC 0113-0141-65

**Original Strength**

**See New Warnings**

Famotidine Tablets, 10 mg

**Acid Reducer**

Actual  
Size

**Just One Tablet!**

**Prevents & Relieves Heartburn  
Due to Acid Indigestion**

Compare to active ingredient of  
Pepcid® AC

**100%**  
SATISFACTION  
GUARANTEED

30 Tablets

14165 C2 C2

OPEN OTHER END

### Drug Facts

**Active ingredient (in each tablet) Purpose**  
Famotidine 10 mg.....Acid reducer

### Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

### Warnings

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

### Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor
- with other acid reducers

### Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain ■ kidney disease

### Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

### Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

### Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F) ■ protect from moisture

**Inactive ingredients** camellia wax, hypromellose, iron oxide red, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, talc, titanium dioxide, triacetin

**Tips for Managing Heartburn** • Do not lie flat or bend over after eating • Do not wear tight-fitting clothing around the stomach • Do not eat before bedtime • Raise the head of your bed • Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables • Eat slowly and avoid big meals • If overweight, lose weight • Quit smoking

CONVENIENT RECLOSING TAB



## GOOD SENSE ACID REDUCER

famotidine tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0113-0141
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>FAMOTIDINE</b> (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	

### Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	L141
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0141-65	30 in 1 CARTON	10/23/2009	

<b>1</b>	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA075400	10/23/2009	

**Labeler -** L. Perrigo Company (006013346)

Revised: 7/2021

L. Perrigo Company