

OXYGEN - oxygen gas
Clinical 1 Home Medical, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Oxygen HP Label

OXYGEN COMPRESSED USP UN 1072 OXYGEN 2 C.A.S. NO. 7782-44-7

Clinical 1 Home Medical 65E Mathewson Drive Weymouth, MA 02189 (800) 261-5737

**WARNING: HIGH PRESSURE, OXIDIZING GAS, VIGOROUSLY ACCELERATES COMBUSTION
NO SMOKING IN THE PRESENCE OF OXYGEN OR A FIRE MAY RESULT.**

Keep oil and grease away. Open valve slowly. Store and use with adequate ventilation.

Use only with equipment cleaned for oxygen service and rated for cylinder pressure.

Use a back flow preventive device in the piping.

Close valve after each use and when empty. Cylinder temperature should not exceed 52 degrees C (125 degrees F).

Use in accordance with Material Safety Data Sheet (MSDS).

DO NOT REMOVE THIS PRODUCT LABEL.

WARNING: For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other

medical applications, Rx only. Uninterrupted use of high concentrations of oxygen over a long duration, without monitoring its effect on

oxygen content of arterial blood, may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not

attempt to use on patients who have stopped breathing unless used in conjunction with resuscitative equipment.

Produced by Air Liquefaction.

Size	Contents	Size	Contents
C	248 Liters	M	3455 Liters
D	415 Liters	H	6900 Liters
E	682 Liters	Other	____ Liters

OXYGEN, COMPRESSED USP

UN 1072

C.A.S. NO. 7782-44-7



Clinical 1 Home Medical

65E Mathewson Drive
Weymouth, MA 02189
(800) 261-5737

⚠ WARNING: HIGH PRESSURE, OXIDIZING GAS. VIGOROUSLY ACCELERATES COMBUSTION. NO SMOKING IN THE PRESENCE OF OXYGEN OR A FIRE MAY RESULT.

Keep oil and grease away. Open valve slowly. Store and use with adequate ventilation. Use only with equipment cleaned for oxygen service and rated for cylinder pressure.

Use a back flow preventative device in the piping.

Close valve after each use and when empty. Cylinder temperature should not exceed 52°C (125°F).

Use in accordance with Material Safety Data Sheet (MSDS).

DO NOT REMOVE THIS PRODUCT LABEL.

⚠ WARNING: For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications, **Rx only**. Uninterrupted use of high concentrations of oxygen over a long duration, without monitoring its effect on oxygen content of arterial blood, may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not attempt to use on patients who have stopped breathing unless used in conjunction with resuscitative equipment.

Produced by Air Liquefaction.

	Size	Contents		Size	Contents
<input type="checkbox"/>	C	248 Liters	<input type="checkbox"/>	M	3455 Liters
<input type="checkbox"/>	D	415 Liters	<input type="checkbox"/>	H	6900 Liters
<input type="checkbox"/>	E	682 Liters	<input type="checkbox"/>	Other	____ Liters

Oxygen Liq Label

OXYGEN REFRIGERATED LIQUID USP OXYGEN 2 UN 1073

CONTENTS: _____ LITERS

ALWAYS KEEP CONTAINER IN UPRIGHT POSITION

WARNING: DO NOT CHANGE OR FORCE FIT CONNECTIONS

EXTREMELY COLD OXIDIZING LIQUID AND GAS UNDER PRESSURE

VIGOROUSLY ACCELERATES COMBUSTION.

NO SMOKING IN THE PRESENCE OF OXYGEN OR A FIRE MAY RESULT

COMBUSTIBLES IN CONTACT WITH LIQUID OXYGEN MAY EXPLODE ON IGNITION OR IMPACT.

CAN CAUSE SEVERE FROSTBITE.

Keep oil, grease, and combustibles away.

Use only with equipment cleaned for oxygen service.

Store and use with adequate ventilation.

Do not get liquid in eyes, on skin or clothing.

For liquid withdraw, wear face shield and gloves.

Do not drop. Use suitable hand truck for container movement.

Avoid spills. Do not walk on or roll equipment over spills.

Container temperature should not exceed 52 degrees C (125 degrees F).

Close valve after each use and when empty.

Use a back flow preventive device in the piping.

Use in accordance with Material Safety Data Sheet (MSDS).

FIRST AID: IN CASE OF FROSTBITE, obtain medical treatment immediately.

WARNING:For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other

medical applications, Rx only.

Uninterrupted use of high concentrations of oxygen over a long duration, without monitoring its effect on oxygen content of arterial blood,

may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not attempt to use on patients who

have stopped breathing unless used in conjunction with resuscitative equipment.

Produced by Air Liquefaction

DO NOT REMOVE THIS LABEL.

Clinical 1 Home Medical

65 E Mathewson Drive

Weymouth, MA 02189

(800) 261-5737

CAS NO: 7782-44-7

**OXYGEN,
REFRIGERATED
LIQUID
USP**
UN 1073



CONTENTS: _____ LITERS

↑ ALWAYS KEEP CONTAINER IN UPRIGHT POSITION. ↑

⚠ WARNING: DO NOT CHANGE OR FORCE FIT CONNECTIONS
EXTREMELY COLD OXIDIZING LIQUID AND GAS UNDER
PRESSURE
VIGOROUSLY ACCELERATES COMBUSTION.
NO SMOKING IN THE PRESENCE OF OXYGEN OR A FIRE MAY RESULT
COMBUSTIBLES IN CONTACT WITH LIQUID OXYGEN MAY
EXPLODE ON IGNITION OR IMPACT.
CAN CAUSE SEVERE FROSTBITE.
Keep oil, grease, and combustibles away

Keep oil, grease, and combustibles away.

Use only with equipment cleaned for oxygen service.

Store and use with adequate ventilation.

Do not get liquid in eyes, on skin or clothing.

For liquid withdrawal, wear face shield and gloves.

Do not drop. Use suitable hand truck for container movement.

Avoid spills. Do not walk on or roll equipment over spills.

Container temperature should not exceed 52°C (125°F).

Close valve after each use and when empty.

Use a back flow preventive device in the piping.

Use in accordance with the Material Safety Data Sheet (MSDS).

△ **FIRST AID:** IN CASE OF FROSTBITE, obtain medical treatment immediately.

△ **WARNING:** For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications, **Rx Only.**

Uninterrupted use of high concentrations of oxygen over a long duration, without monitoring its effect on oxygen content of arterial blood, may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not attempt to use on patients who have stopped breathing, unless used in conjunction with resuscitative equipment.

Produced by Air Liquefaction

DO NOT REMOVE THIS PRODUCT LABEL.

Clinical 1 Home Medical

65E Mathewson Drive

Weymouth, MA 02189

(800) 261-5737

CAS NO: 7782-44-7

OXYGEN

oxygen gas

Product Information

Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:65919-001
Route of Administration	RESPIRATORY (INHALATION)	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYGEN (OXYGEN)	OXYGEN	99 L in 100 L

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65919-001-01	170 L in 1 CYLINDER		
2	NDC:65919-001-02	248 L in 1 CYLINDER		
3	NDC:65919-001-03	415 L in 1 CYLINDER		
4	NDC:65919-001-04	682 L in 1 CYLINDER		
5	NDC:65919-001-05	3455 L in 1 CYLINDER		
6	NDC:65919-001-06	6900 L in 1 CYLINDER		
7	NDC:65919-001-07	31 L in 1 DEWAR		
8	NDC:65919-001-08	41 L in 1 DEWAR		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved medical gas		06/01/2000	

Labeler - Clinical 1 Home Medical, Inc. (110832404)**Registrant** - Clinical 1 Home Medical, Inc. (110832404)**Establishment**

Name	Address	ID/FEI	Business Operations
Clinical 1 Home Medical, Inc.		110832404	repack(65919-001)

Revised: 12/2012

Clinical 1 Home Medical, Inc.