

**COOLING SEVERE DAYTIME COOLING SEVERE NIGHTTIME- acetaminophen,  
dextromethorphan hydrobromide, doxylamine succinate, phenylephrine  
hydrochloride, guaifenesin  
CVS Pharmacy**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**CVS Pharmacy, Inc. Cooling Severe Daytime Cooling Sever Nighttime Drug Facts**

**Active ingredients (in each 30 mL) – Night Time**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- sore throat
- runny nose and sneezing
- cough to help you sleep
- cough due to minor throat and bronchial irritation
- reduces swelling of nasal passages
- fever
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- be careful when driving a motor vehicle or operating machinery
- marked drowsiness may occur

- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks

**Stop use and ask a doctor if**

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- you get nervous, dizzy or sleepless
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 30 mL contains:** sodium 42 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients**

alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

**Questions or comments?**

**1-800-719-9260**

## **Active ingredients (in each 30 mL) - Day Time**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

## **Purpose**

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

## **Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- promotes nasal and/or sinus drainage

## **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- diabetes
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**When using this product**

**do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right

away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

### Other information

- **each 30 mL contains:** sodium 41 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

### Inactive ingredients

alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

### Questions or comments?

**1-800-719-9260**

### Package/Label Principal Display Panel

Compare to the active ingredients in Vicks® DayQuil® Severe+VapoCOOL™

Cooling

Severe Daytime

COLD & FLU RELIEF

ACETAMINOPHEN – Pain reliever/Fever reducer

PHENYLEPHRINE HCl – Nasal decongestant

DEXTROMETHORPHAN HBr – Cough suppressant

GUAIFENESIN – Expectorant

Relieves:

Minor aches, pains & fever

Nasal congestion & sinus pressure

Cough

Chest congestion

ALCOHOL 10%

12 FL OZ (355 mL)

Compare to the active ingredients in Vicks® NyQuil® Severe+VapoCOOL™

Cooling

Severe Nighttime

COLD & FLU RELIEF

ACETAMINOPHEN – Pain reliever/Fever reducer

PHENYLEPHRINE HCl – Nasal decongestant

DOXYLAMINE SUCCINATE - Antihistamine

DEXTROMETHORPHAN HBr – Cough suppressant

Relieves:

Minor aches, pains & fever

Nasal congestion & sinus pressure

Sneezing, runny nose

Cough

ALCOHOL 10%

12 FL OZ (355 mL)

<p><b>PARENTS:</b> Learn about teen medicine abuse <a href="http://www.StopMedicineAbuse.org">www.StopMedicineAbuse.org</a></p> <p><b>DO NOT USE IF PRINTED NECKBANDS BROKEN OR MISSING</b></p> <p>*These products are not manufactured or distributed by Packer &amp; Gamble, distributor of Vicks® DayQuil® Severe+VapoCOOL™ and Vicks® NyQuil® Severe+VapoCOOL™.</p> <p><b>Drug Facts (continued)</b> Inactive ingredients: alcohol, anhydrous citric acid, D&amp;C yellow #10, edetate disodium, FD&amp;C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose</p> <p>Questions or comments? 1-800-719-9260</p>	<p><b>CVS Health</b></p> <p>Compare to the active ingredients in Vicks® DayQuil® Severe+VapoCOOL™**</p>	<p><b>CVS Health</b></p> <p>Compare to the active ingredients in Vicks® NyQuil® Severe+VapoCOOL™**</p> <p>NDC 69842-737-02</p>	<p>Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 © 2019 CVS/pharmacy CVS.com® 1-800-SHOP CVS V-16430</p> <p><b>CVS Quality</b> Money Back Guarantee</p> <p><b>Drug Facts (continued)</b> Inactive ingredients: alcohol, anhydrous citric acid, D&amp;C yellow #10, edetate disodium, FD&amp;C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose</p> <p>Questions or comments? 1-800-719-9260</p>
	<p><b>Cooling Severe Daytime</b></p> <p><b>COLD &amp; FLU RELIEF</b></p> <p>ACETAMINOPHEN - Pain reliever/Fever reducer PHENYLEPHRINE HCl - Nasal decongestant DEXTROMETHORPHAN HBr - Cough suppressant GUAIFENESIN - Expectorant</p> <p><b>Relieves:</b></p> <ul style="list-style-type: none"> <li>• Minor aches, pains &amp; fever</li> <li>• Nasal congestion &amp; sinus pressure</li> <li>• Cough</li> <li>• Chest congestion</li> </ul> <p>ALCOHOL 10%</p> <p>12 FL OZ (355 mL)</p>	<p><b>Cooling Severe Nighttime</b></p> <p><b>COLD &amp; FLU RELIEF</b></p> <p>ACETAMINOPHEN - Pain reliever/Fever reducer PHENYLEPHRINE HCl - Nasal decongestant DOXYLAMINE SUCCINATE - Antihistamine DEXTROMETHORPHAN HBr - Cough suppressant</p> <p><b>Relieves:</b></p> <ul style="list-style-type: none"> <li>• Minor aches, pains &amp; fever</li> <li>• Nasal congestion &amp; sinus pressure</li> <li>• Sneezing, runny nose</li> <li>• Cough</li> </ul> <p>ALCOHOL 10%</p> <p>12 FL OZ (355 mL)</p>	

Drug Facts	Nighttime	Drug Facts	Daytime												
<b>Active ingredients (in each 30 mL)</b> Acetaminophen 650 mg Dextromethorphan HBr 20 mg Doxylamine succinate 12.5 mg Phenylephrine HCl 10 mg		<b>Active ingredients (in each 30 mL)</b> Acetaminophen 650 mg Dextromethorphan HBr 20 mg Guaifenesin 400 mg Phenylephrine HCl 10 mg													
<b>Purpose</b> Pain reliever/fever reducer Cough suppressant Antihistamine Nasal decongestant		<b>Purpose</b> Pain reliever/fever reducer Cough suppressant Expectorant Nasal decongestant													
<b>Uses</b> Temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> <li>nasal congestion</li> <li>sinus congestion and pressure</li> <li>minor aches and pains</li> <li>headache</li> <li>sore throat</li> <li>runny nose and sneezing</li> <li>cough to help you sleep</li> <li>promote minor throat and bronchial irritation</li> <li>reduce swelling of nasal passages</li> <li>fever</li> <li>promote nasal and/or sinus drainage</li> <li>temporarily restores free breathing through the nose</li> </ul>		<b>Uses</b> Temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> <li>nasal congestion</li> <li>sinus congestion and pressure</li> <li>cough due to minor throat and bronchial irritation</li> <li>minor aches and pains</li> <li>headache</li> <li>fever</li> <li>reduce swelling of nasal passages</li> <li>temporarily restores free breathing through the nose</li> <li>sore throat</li> <li>helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both so the mucus and malle coughs more productive</li> <li>promotes nasal and/or sinus drainage</li> </ul>													
<b>Warnings</b> Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none"> <li>more than 4,000 mg of acetaminophen in 24 hours</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> <li>skin redness</li> <li>hives</li> <li>itching</li> <li>swelling</li> <li>skin rash</li> </ul> Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		<b>Warnings</b> Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none"> <li>more than 4,000 mg of acetaminophen in 24 hours</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> <li>skin redness</li> <li>hives</li> <li>itching</li> <li>swelling</li> <li>skin rash</li> </ul> Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.													
<b>Do not use</b> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> <li>if you have ever had an allergic reaction to this product or any of its ingredients.</li> </ul>		<b>Do not use</b> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> <li>if you have ever had an allergic reaction to this product or any of its ingredients.</li> </ul>													
<b>Ask a doctor before use if you have</b> <ul style="list-style-type: none"> <li>liver disease</li> <li>heart disease</li> <li>high blood pressure</li> <li>breathing problem such as emphysema or chronic bronchitis</li> <li>thyroid disease</li> <li>diabetes</li> <li>glaucoma</li> <li>cough that occurs with too much phlegm (mucus)</li> <li>trouble urinating due to an enlarged prostate gland</li> <li>persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li> <li>a sodium-restricted diet</li> </ul>		<b>Ask a doctor before use if you have</b> <ul style="list-style-type: none"> <li>liver disease</li> <li>heart disease</li> <li>thyroid disease</li> <li>high blood pressure</li> <li>trouble urinating due to an enlarged prostate gland</li> <li>cough that occurs with too much phlegm (mucus)</li> <li>persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema</li> <li>diabetes</li> <li>a sodium-restricted diet</li> </ul>													
<b>Ask a doctor or pharmacist before use if you are</b> <ul style="list-style-type: none"> <li>taking sedatives or tranquilizers</li> <li>taking the blood thinning drug warfarin</li> </ul>		<b>Ask a doctor or pharmacist before use if you are</b> <ul style="list-style-type: none"> <li>taking the blood thinning drug warfarin</li> </ul>													
<b>When using this product</b> <ul style="list-style-type: none"> <li>do not use more than directed</li> <li>excitability may occur, especially in children</li> <li>be careful when driving a motor vehicle or operating machinery</li> <li>marked drowsiness may occur</li> <li>alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>avoid alcoholic drinks</li> </ul>		<b>When using this product</b> do not use more than directed													
<b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>pain, nasal congestion, or cough gets worse or lasts more than 7 days</li> <li>you get nervous, dizzy or sleepless</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</li> </ul>		<b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>you get nervous, dizzy or sleepless</li> <li>pain, nasal congestion, or cough gets worse or lasts more than 7 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</li> </ul>													
<b>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.		<b>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.													
<b>Directions</b> <ul style="list-style-type: none"> <li>take only as directed - see Overdose warning</li> <li>only use the dose supplied</li> <li>do not exceed 4 doses per 24 hrs</li> </ul> <table border="1"> <tr> <td>adults &amp; children 12 yrs &amp; over</td> <td>30 mL every 4 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>		adults & children 12 yrs & over	30 mL every 4 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use	<b>Directions</b> <ul style="list-style-type: none"> <li>take only as directed - see Overdose warning</li> <li>only use the dose provided</li> <li>do not exceed 4 doses per 24 hrs</li> </ul> <table border="1"> <tr> <td>adults &amp; children 12 yrs &amp; over</td> <td>30 mL every 4 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>		adults & children 12 yrs & over	30 mL every 4 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use
adults & children 12 yrs & over	30 mL every 4 hrs														
children 4 to under 12 yrs	ask a doctor														
children under 4 yrs	do not use														
adults & children 12 yrs & over	30 mL every 4 hrs														
children 4 to under 12 yrs	ask a doctor														
children under 4 yrs	do not use														
<b>Other information</b> <ul style="list-style-type: none"> <li>each 30 mL contains sodium 4 mg</li> <li>store at 20-25°C (68-77°F)</li> </ul>		<b>Other information</b> <ul style="list-style-type: none"> <li>each 30 mL contains sodium 41 mg</li> <li>store at 20-25°C (68-77°F). Do not refrigerate.</li> </ul>													

COOLING SEVERE DAYTIME COOLING SEVERE NIGHTTIME			
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride, guaifenesin kit			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69842-737
<b>Packaging</b>			
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>
1	NDC:69842-737-02	1 in 1 PACKAGE; Type 0: Not a Combination Product	07/23/2019
<b>Quantity of Parts</b>			
<b>Part #</b>	<b>Package Quantity</b>	<b>Total Product Quantity</b>	
Part 1	1 BOTTLE	355 mL	
Part 2	1 BOTTLE	355 mL	
<b>Part 1 of 2</b>			

# COOLING SEVERE NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride solution

## Product Information

Item Code (Source) NDC:69842-590

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-590-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

**Part 2 of 2****COOLING SEVERE DAYTIME**

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

**Product Information**

<b>Item Code (Source)</b>	NDC:69842-182
---------------------------	---------------

<b>Route of Administration</b>	ORAL
--------------------------------	------

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

**Inactive Ingredients**

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-182-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	06/20/2019	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/23/2019	

**Labeler** - CVS Pharmacy (062312574)

Revised: 1/2020

CVS Pharmacy