

**SOFT-SAN- lactic acid liquid  
IBA**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**SOFT-SAN**

**ACTIVE INGREDIENTS:**

Lactic Acid.....1.00%

**INERT INGREDIENTS**

Skin Conditioners.....35.00%

Other Inerts.....64.00%

(Includes Freeze Suppressants, Wintergreen,  
pH Buffers, Color Enhancers and Water)

Total.....100.00%

**GENERAL RECOMMENDATIONS**

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

**FIRST AID PROCEDURES**

**EYE CONTACT:** IMMEDIATELY FLUSH EYE LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

**SKIN CONTACT:** WASH ANY SENSITIVE AREAS.

**INGESTION:** DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION

INTO THE LUNGS.

## **RECOMMENDED PROCEDURES FOR TEAT SANITATION**

### **A. PRE-MILKING (PRE-DIPPING)**

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip or spray the cow's teats full length with Soft-San Teat Dip or your choice of another IBA pre-dip product.
4. Wait approximately 30 seconds.
5. Remove all teat dip by thoroughly drying the teats and teat orifices with a clean, single service paper towel(s).
6. Attach milking unit.

### **B. POST-MILKING (POST-DIPPING)**

Immediately after milking, dip or spray all teats full length with SOFT-SAN Teat Dip. Ensure good coverage of all sides of the teats and teat orifices.

## **WARNING**

**DO NOT DILUTE THIS PRODUCT.**

**DO NOT RETURN TEA DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.**

**ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.**

**PROTECT THIS PRODUCT FROM FREEZING.**

**Soft-San**

NDC 29983-1052-6



# SOFT-SAN

## ALL WEATHER TEAT DIP

A HIGHLY EFFECTIVE  
SANITIZING TEAT DIP

CONTAINS 35%  
SKIN CONDITIONERS

HELPS HEAL OR PREVENT DAMAGED TEATS DUE  
TO SUDDEN AND/OR SEVERE TEMPERATURE CHANGES

**ACTIVE INGREDIENT:**

Lactic Acid .....	1.00%
<b>INERT INGREDIENTS:</b>	
Skin Conditioners .....	35.00%
Other Inerts .....	64.00%
(Includes Freeze Suppressants, Wintergreen, pH Buffers, Color Enhancers and Water)	
<b>TOTAL:</b> .....	<b>100.00%</b>

**GENERAL RECOMMENDATIONS**

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.


**FIRST AID PROCEDURES**

**EYE CONTACT:** IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

**SKIN CONTACT:** WASH ANY SENSITIVE AREAS.

**INGESTION:** DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION  
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**



**RECOMMENDED PROCEDURES  
FOR TEAT SANITATION**

**A. PRE-MILKING (PRE-DIPPING)**

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip or spray the cow's teats full length with SOFT-SAN Teat Dip or your choice of another IBA pre-dip product.
4. Wait approximately 30 seconds.
5. Remove all teat dip by thoroughly drying the teats and teat orifices with a clean, single service paper towel(s).
6. Attach milking unit.

**B. POST-MILKING (POST-DIPPING)**

Immediately after milking, dip or spray all teats full length with SOFT-SAN Teat Dip. Ensure good coverage of all sides of the teats and teat orifices

**WARNING**

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

PROTECT THIS PRODUCT FROM FREEZING.

**A PART OF THE COMPLETE IBA SANITATION PROGRAM**

EXP. DATE:                      LOT NO.:                      NET CONTENTS: 15 GALLONS                      IBA STOCK # 260537                      1121W

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

## SOFT-SAN

lactic acid liquid

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:29983-1052
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	1 kg in 100 kg

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1052-8	215.04 kg in 1 DRUM		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/11/2014	

**Labeler** - IBA (019494160)

**Registrant** - IBA (019494160)

## Establishment

Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc. DBA Custom Chemical Formulators		028311595	manufacture, api manufacture, pack

Revised: 12/2023

IBA