

NIX LICE KILLING CREME RINSE- permethrin rinse
Geodis Logistics LLC

Nix Lice Killing Creme Rinse

Drug Facts

Active ingredient (in each fluid ounce)

Permethrin 280 mg (1%)

Purpose

Lice treatment

Uses

treats head lice

Warnings

For external use only

Do not use

- on children under 2 months of age
- near the eyes
- inside the nose, ear, mouth, or vagina
- on lice in eyebrows or eyelashes.
See your doctor.

Ask a doctor before use if you are

allergic to ragweed. May cause breathing difficulty or an asthmatic episode.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets into the eyes, immediately flush with large amounts of water
- scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

Important: Read warnings before use

- **Adults and children 2 months old and older**

Inspect

- all household members should be checked by another person for lice and/or nits (eggs)
- use a magnifying glass in bright light to help you see the lice and/or nits (eggs)
- use a tool, such as a comb or two unsharpened pencils to lift and part the hair
- look for tiny nits near scalp, beginning at the back of the neck and behind the ears
- small sections of hair (1-2 inches wide) should be examined at a time
- unlike dandruff, nits stick to the hair. Dandruff should move when lightly touched,
- if either lice or nits (or eggs) are found, treat with Nix[®] Creme Rinse

Treat

- wash hair with a shampoo without conditioner. Do not use a shampoo that contains a conditioner or a conditioner alone since this may decrease the activity of Nix[®]. Rinse with water.
- towel dry hair so it is damp but not wet
- shake the bottle of Nix[®] well
- completely saturate the hair and scalp with Nix[®]. Begin to apply Nix[®] behind the ears and at the back of the neck.
- keep Nix[®] out of the eyes. Protect the eyes with a washcloth or towel.
- leave Nix[®] on the hair for 10 minutes, but no longer
- rinse with warm water
- towel dry hair and comb out tangles
- if live lice are seen seven days or more after the first treatment, a second treatment should be given

Remove Lice/Nits

- remove nits by combing the hair with the special small tooth comb provided. Remaining nits may be removed by hand (using a throw-away glove), or cutting the nits out.
- use the nit comb provided and make sure the hair remains slightly damp while removing nits
- if the hair dries during combing, dampen it slightly with water
- part the hair into 4 sections. Work on one section at a time. Longer hair may take more time (1-2 hours).
- start at the top of the head on the section you have picked
- with one hand, lift a 1-2 inch wide strand of hair. Get the teeth of the comb as close to the scalp as possible and comb with a firm, even motion away from the scalp to the end of the hair.
- use clips to pin back each strand of hair after you have combed out the nits

- clean the comb completely as you go. Wipe the nits from the comb with a tissue and throw away the tissue in a sealed plastic bag to prevent the lice from coming back.
- after combing, recheck the entire head for nits and repeat combing if necessary
- check the affected head daily to remove any nits that you might have missed

Other information

- read all the directions in the Consumer Information Insert and warnings before use. Keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77°F)

Inactive ingredients

canada balsam, cetyl alcohol, citric acid, FD&C yellow no. 6, hydrolyzed animal protein, hydroxyethylcellulose, isopropyl alcohol, methylparaben, polyoxyethylene 10 cetyl ether, propylene glycol, propylparaben, stearylalkonium chloride, water

Questions or comments?

call **1-888- LICE (1-888-542-3546)**.

Monday to Friday, 9 AM - 5 PM EST

Package Labeling:

Nix

PERMETHRIN



Have questions?
Call our expert hotline
888.542.3546

Creme Rinse is Safe for use on children as young as 2 months old.

Lice Killing Creme Rinse

Drug Facts (continued)

Directions

Important: Read warnings before use

■ **Adults and children 2 months of age and over:**

Inspect

- all household members should be checked by another person for lice and/or nits (eggs)
- use a magnifying glass in bright light to help you see the lice and nits (eggs)
- use a tool, such as a comb or two unsharpened pencils to lift and part the hair
- look for tiny nits near the scalp, beginning at the back of the neck and behind the ears
- small sections of hair (1-2 inches wide) should be examined at a time
- unlike dandruff, nits stick to the hair. Dandruff should move when lightly touched.
- if either lice or nits (eggs) are found, treat with Nix® Creme Rinse

Treat

- Wash hair with a shampoo without conditioner. Do not use a shampoo that contains a conditioner or a conditioner alone since this may decrease the activity of Nix®, Rinse with water.
- towel dry hair so it is damp but not wet
- shake the bottle of Nix® well
- completely saturate the hair and scalp with Nix®. Begin to apply Nix® behind the ears and at the back of the neck.
- keep Nix® out of the eyes. Protect the eyes with a washcloth or towel.
- leave Nix® on the hair for 10 minutes, but no longer
- rinse with warm water
- towel dry hair and comb out tangles
- if live lice are seen seven days or more after the first treatment, a second treatment should be given

Remove Lice/Nits

- remove nits by combing the hair with the special small tooth comb provided. Remaining nits may be removed by hand (using a throw-away glove), or cutting the nits out.
- use the nit comb provided and make sure the hair remains slightly damp while removing nits
- if the hair dries during combing, dampen it slightly with water
- part the hair into 4 sections. Work on one section at a time. Longer hair may take more time (1-2 hours).
- start at the top of the head on the section you have picked
- with one hand, lift a 1-2 inch wide strand of hair. Get the teeth of the comb as close to the scalp as possible and comb with a firm, even motion away from the scalp to the end of the hair.
- use clips to pin back each strand of hair after you have combed out the nits
- clean the comb completely as you go. Wipe the nits from the comb with a tissue and throw away the tissue in a sealed plastic bag to prevent the lice from coming back.
- after combing, recheck the entire head for nits and repeat combing if necessary
- check the affected head daily to remove any nits that you might have missed

Drug Facts (continued)

Other information

■ Read all the directions in the Consumer Information Insert and warnings before use. Keep the carton. It contains important information.

- store at 20° to 25°C (68° to 77°F)

Inactive ingredients

canada balsam, cetyl alcohol, citric acid, FD&C yellow no. 6, hydrolyzed animal protein, hydroxyethylcellulose, isopropyl alcohol, methylparaben, polyoxyethylene 10 cetyl ether, propylene glycol, propylparaben, stearylaluminum chloride, water

Questions or comments?

call 1-888-LICE LINE (1-888-542-3546), Monday to Friday, 9 AM - 5 PM EST

LIMITED CONSUMER WARRANTY

If you are not satisfied with Nix® we will refund your purchase price. Send cash register receipt, original UPC code, and reason for dissatisfaction to: INSIGHT Pharmaceuticals LLC, 660 White Plains Rd., Tarrytown, NY 10591

nixlice.com

Dist. by: INSIGHT Pharmaceuticals LLC
660 White Plains Road
Tarrytown, NY 10591
A Prestige Consumer Healthcare company
© 2021 Trade dress is owned by Insight Pharmaceuticals LLC.
All rights reserved.
Creme Rinse made in India.
Comb made in China.

NXUS003102

Nix

PERMETHRIN

Have questions?
Call our expert hotline
888.542.3546



Creme Rinse is Safe for use on children as young as 2 months old.

Lice Killing Creme Rinse

- ✓ Kills lice and eggs in one application
- ✓ Prevents reinestation*
- ✓ Maximum Strength

Includes 1 bottle
Plus 1 nit comb
Spanish Instructions Included

*For up to 14 days



2 FL OZ (59 mL)

Drug Facts

Active ingredient (in each fluid ounce) Permethrin 280 mg (1%). **Purpose** Lice treatment

Use treats head lice

Warnings

For external use only

Do not use

- on children under 2 months of age
- near the eyes
- inside the nose, ear, mouth, or vagina
- on lice in eyebrows or eyelashes. *See your doctor.*

Ask a doctor before use if you are allergic to ragweed. May cause breathing difficulty or an asthmatic episode.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets into the eyes, immediately flush with large amounts of water
- scalp itching or redness may occur

Stop use and see a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.



3 63736 12002 3

Expiration Date

Lot #

Active ingredient (in each fluid ounce) Purpose
 Permethrin 280 mg (1%) Lice treatment

Use treats head lice

Warnings

For external use only

Do not use ■ on children under 2 months of age
 ■ near the eyes ■ inside the nose, ear, mouth, or vagina ■ on lice in eyebrows or eyelashes. See your doctor.

Ask a doctor before use if you are ■ allergic to ragweed. May cause breathing difficulty or an asthmatic episode.

When using this product ■ keep eyes tightly closed and protect eyes with a washcloth or towel
 ■ if product gets into the eyes, immediately flush with large amounts of water ■ scalp itching or redness may occur

Stop use and ask a doctor if ■ breathing difficulty occurs ■ eye irritation occurs ■ skin or scalp irritation continues or infection occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Nix
 Creme Rinse



✓ Kills lice and eggs in one application

✓ Prevents reinfestation*
 * For up to 14 days.

2 FL OZ (59 mL)

Directions See Consumer Information Insert for important directions on inspection for head lice, removing nits (eggs) and cleaning the home and personal items

Treat ■ wash hair with a shampoo without conditioner. Do not use a shampoo that contains a conditioner or a conditioner alone since this may decrease the activity of Nix®. Rinse with water. ■ towel dry hair so it is damp but not wet ■ shake the bottle of Nix® well ■ completely saturate the hair and scalp with Nix®. Begin to apply Nix® behind the ears and at the back of the neck. ■ keep Nix® out of the eyes. Protect the eyes with a washcloth or towel. ■ leave Nix® on the hair for 10 minutes, but no longer ■ rinse with warm water ■ towel dry hair and comb out tangles ■ if live lice are seen seven days or more after the first treatment, a second treatment should be given

Other information ■ Read all the directions on the Consumer Information Insert and warnings before use. Keep the carton. It contains important information ■ Store at 20° to 25° C (68° to 77° F)

Inactive ingredients canada balsam, cetyl alcohol, citric acid, FD&C yellow no. 6, hydrolyzed animal protein, hydroxyethylcellulose, isopropyl alcohol, methylparaben, polyoxyethylene 10 cetyl ether, propylene glycol, propylparaben, stearylkonium chloride, water

Questions? 1-888-LICE LINE (1-888-542-3546) nixlice.com

Distributed by: INSIGHT Pharmaceuticals LLC, Tarrytown, NY 10591, a Prestige Consumer Healthcare company
 ©2021 Trade dress is owned by Insight Pharmaceuticals LLC. All rights reserved. NXUS003201

LOT:

EXP:

NIX LICE KILLING CREME RINSE

permethrin rinse

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63471-3006(NDC:63736-030)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PERMETHRIN (UNII: 509F88P9SZ) (PERMETHRIN - UNII:509F88P9SZ)	PERMETHRIN	1 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
CANADA BALSAM (UNII: K304V198I3)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
CETETH-10 (UNII: LF9X1PN3XJ)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
STEARALKONIUM CHLORIDE (UNII: 00UO26BB88)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	orange (Light Orange)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63471-3006-2	1 in 1 CARTON	11/15/2013	
1		59 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019918	11/15/2013	

Labeler - Geodis Logistics LLC (877844097)

Revised: 2/2022

Geodis Logistics LLC