REZAMID- sulfur and resorcinol lotion Summers Laboratories Inc

SUMMERS LABS (as PLD) - REZAMID (11086-022)

ACTIVE INGREDIENTS

SULFUR 5%

RESORCINOL 2%

PURPOSE

ACNE TREATMENT LOTION

USE

DRIES UP ACNE PIMPLES, HELPS PREVENT NEW PIMPLES

WARNINGS

FOR EXTERNAL USE ONLY

DO NOT USE

- ON BROKEN SKIN
- ON LARGE AREAS OF THE BODY

WHEN USING THIS PRODUCT

- APPLY TO AFFECTED AREAS ONLY
- DO NOT GET INTO EYES
- skin irritation and dryness is more likely to occur if you use another topical acne medication at the same time. If irritation occurs, only use one topical acne medication at a time

STOP USE AND ASK A DOCTOR IF

IF SKIN IRRITATION OCCURS OR GETS WORSE

KEEP OUT OF REACH OF CHILDREN. IF SWALLOWED, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.

DIRECTIONS

- shake very well before using
- clean the skin thoroughly before applying this product
- cover the entire affected area with a thin layer one to three times daily
- because excessive drying of the skin may occur, start with one application daily, then gradually increase to two or three times daily if needed or as directed by a doctor
- if bothersome dryness or peeling occurs, reduce application to once a day or every

INACTIVE INGREDIENTS

water, SD-40 alcohol 28%, zinc oxide, talc, titanium dioxide, propylene glycol, attapulgite, lauramide DEA, iron oxides, sodium bisulfite, PEG-8 laurate, parachlorometaxylenol, hydroxyethylcellulose, sodium chloride, sodium polynapthalene sulfonate, EDTA, methyl paraben, xanthan gum, butylparaben, fragrance, simethicone.



REZAMID sulfur and resorcinol lotion Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:11086-022

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
SULFUR (UNII: 70FD1KFU70) (SULFUR - UNII:70FD1KFU70)	SULFUR	5 g in 100 mL		
RESORCINOL (UNII: YUL4L094HK) (RESORCINOL - UNII:YUL4L094HK)	RESORCINOL	2 g in 100 mL		

Inactive Ingredients				
Ingredient Name	Strength			
WATER (UNII: 059QF0KO0R)				
ALCOHOL (UNII: 3K9958V90M)				
ZINC OXIDE (UNII: SOI2LOH54Z)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
ATTAPULGITE (UNII: U6V729APAM)				
LAURIC DIETHANOLAMIDE (UNII: 12912VHG38)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
SODIUM BISULFITE (UNII: TZX5469Z6I)				
PEG-8 LAURATE (UNII: 76208IWA10)				
CHLOROXYLENOL (UNII: 0F32U78V2Q)				
HYDROXYETHYL CELLULOSE (140 MPA.S AT 5%) (UNII: 8136Y38GY5)				
SODIUM CHLORIDE (UNII: 451W47IQ8X)				
SODIUM NAPHTHALENESULFONATE (UNII: D3F8YRX7TP)				
EDETIC ACID (UNII: 9G34HU7RV0)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
XANTHAN GUM (UNII: TTV12P4NEE)				
BUTYLPARABEN (UNII: 3QPI1U3FV8)				
DIMETHICONE (UNII: 92RU3N3Y1O)				
SILICON (UNII: Z4152N8IUI)				

Pa	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
	NDC:11086- 022-01	56.7 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/30/2013		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M032	10/30/2013		

Revised: 10/2023 Summers Laboratories Inc