AUTO 157 - benzalkonium chloride, lidocaine, benzocaine, alcohol, ibuprofen, acetaminophen, aspirin
Genuine First Aid LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Auto 157

ACTIVE INGREDIENTS:

Benzalkonium Chloride 0.13%
Lidocaine HCL 0.5%

Purpose: First aid antiseptic, external analgesic

Uses: First aid to help prevent infection and for the temporary relief of pain and itching associated with:

Minor Cuts

Scrapes

Burns

Warnings:

For external use only

Do not use: In eyes, in large quantities, over raw blistered areas, or on deep puncture wounds, animal bites or serious burns, for more than one week

Keep out of reach of Children.

If ingested, contact a Poison Control Center right away.

Directions: Clean affected area, Apply small amount not more than 3 times daily.

May be covered with a sterile bandage.

Other Information:

Store at room temperature

LOT/EXP: Made in CHINA

20130301

Genuine First Aid Burn Cream

Antiseptic Pain Relief With Aloe

Net Wt 0.9g (1/32 oz)

Manufactured in CHINA for

Genuine First Aid.

Reorder BC-001

Genuine First Aid Burn Cream

Antiseptic Pain Relief With Aloe

ACTIVE INGREDIENTS:
Benzalkonium Chloride 0.13%
Lidocaine HCL 0.5%

Net Wt 0.9g - (1/32 oz)

Manufactured in CHINA for GENUINE FIRST AID®

Drug Facts

Purpose: First aid antiseptic, external analgesic

Uses: First aid to help prevent infection and for the temporary relief of pain and itching associated with: • minor cuts • scrapes • burns

Warnings:

For external use only

Do not use • in eyes • in large quantities • over raw blistered areas, or on deep puncture wounds, animal bites or serious burns • for more than one week

Keep out of reach of children. If ingested, contact a Poison Control Center right away.

Directions: • clean affected area • apply small amount not more than 3 times daily • may be covered with a sterile bandage

Other information: Store at room temperature

LOT 600609 2015/01

Active Ingredient: Purpose

Benzalkonium Chloride 0.40%..... First Aid Antiseptic

Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warning: For external use only.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.

Do not use in the eyes or over large areas of the body.

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive Ingredient: Purified water

LOT/EXP: Made in CHINA

20130301

Antiseptic Towelette

Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID

DRUG FACTS - Antiseptic Towelette

Active Ingredient: **Purpose:**
Benzalkonium Chloride 0.40%..First Aid Antiseptic
Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.
Warnings: For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
If unusual redness, swelling or other symptoms occur, consult a physician immediately.
Do not use: In the eyes, or over large areas of the body.
Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.
Inactive ingredient: Purified water.

Made in CHINA

- - TEAR HERE - - - -

Active Ingredient: Purpose:

Benzocaine, 6% w/v..... Topical Anesthetic

SD alcohol, 60% w/v..... Antiseptic

Use: For the temporary relief of pain and itching associated with minor burns, scrapes and insect bites.

Warnings: For external use only.

Avoid contact with eyes. If this happens, rinse thoroughly with water.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Flammable - keep away from fire or flame.

Do not use: In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.

Made in CHINA

LOT/EXP:

Insect Sting Relief Pad

Genuine First Aid LLC, Clearwater FL 33755
www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID

DRUG FACTS - Insect Sting Relief Pad

Active Ingredient:	**Purpose:**
Benzocaine, 6% w/v	Topical Anesthetic
SD alcohol, 60% w/v	Antiseptic

Use: For the temporary relief of pain and itching associated with minor burns, scrapes and insect bites.

Warnings: For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Flammable - keep away from fire or flame. Avoid contact with eyes. If this happens, rinse thoroughly with water.

Do not use: In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.

Made in CHINA

- - - - TEAR HERE - - - -

Active ingredient (in each tablet) Purpose

Ibuprofen USP (NSAID*) 200mg . . . . . .Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to:

the common cold

headache

toothache

muscular aches

backache

minor pain of arthritis

menstrual cramps temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: shock, facial swelling, asthma (wheezing) rash, skin reddening, blisters, hives If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach
bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinner (anticoagulant) or steroid drug, take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have a high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor before use if you are:
taking any other drug containing NSAID (prescription or nonprescription); taking aspirin for heart attack or stroke, because Ibuprofen may decrease this benefit of aspirin; taking any other drug

When using this product:
take with food or milk if stomach upset occurs

Stop use and ask a doctor If:
you experience any of the following signs of stomach bleeding; feel faint; vomit blood; have bloody or black stools; have stomach pain that does get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:
do not use more than directed; the smallest effective dose should be used; do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information: Store at controlled room temperature; avoid excessive heat 40 degree Celsius (104 degree Fahrenheit); tamper evident sealed packets; do not use any opened or torn packets

Inactive ingredients: cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

Distributed by GENUINE FIRST AID
600 Cleveland Str Suite 400, Clearwater, FL 33755

IBUPROFEN 2 Tablets

IBUPROFEN 2 Tablets

Active ingredient (in each tablet) ***Purpose***
Ibuprofen USP (NSAID*) 200mg . . . . . Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to: ■ the common cold ■ headache ■ toothache ■ muscular aches ■ backache ■ minor pain of arthritis ■ menstrual cramps temporarily reduces fever

Warnings **Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ shock ■ facial swelling ■ asthma (wheezing) ■ rash ■ skin reddening ■ blisters ■ hives If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking any other drug containing an NSAID (prescription or nonprescription) ▶

Warnings (continued)
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product
■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is presen in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ **do not use more than directed** ■ **the smallest effective dose should be used** ■ do not take longer than 10 days, unless directed by a doctor

Adults and Children (12 years and older): Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information ■ store at controlled room temperature ■ avoid excessive heat 40° C (104° F) ■ tamper evident sealed packets ■ do not use any opened or torn packets

Inactive ingredients cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin

Distributed by **GENUINE FIRST AID**
600 Cleveland Str Suite 400, Clearwater, FL 33755

Active Ingredient (in each tablet) Purpose

Acetaminophen 325 mg ..... Analgesic/antipyretic

Uses

temporary relief of minor aches and pains associated with:

common cold; headache; toothache; muscular aches; backache; arthritis; menstrual cramps; and reduction of fever

Warnings:

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if: adult takes more than 12 tablets in 24 hours, which is the maximum daily amount; child takes more than 5 doses in 24 hours; taken with other drugs containing acetaminophen; adult has 3 or more alcoholic drinks every day while using this product

Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; for more than 10 days for pain unless directed by a doctor; for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if: symptoms do not improve; pain gets worse or lasts for more than 10 days; fever gets worse or lasts for more than 3 days; new symptoms occur; redness or swelling is present; a rare sensitivity reaction occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Do not exceed recommended dosage

Directions

Adults and Children 12 years of age or older	Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.
Children 6-11 years of age	Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.
Children under 6 years of age	Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.

Store at 59-86 degree Fahrenheit (15-30 degree Celcius)

tamper evident sealed packets; do not use any open or torn packets

Distributed by GENUINE FIRST AID
600 Cleveland Str Suite 400, Clearwater, FL 33755

GENUINE FIRST AID 2 Tablets

NON-ASPIRIN

GENUINE FIRST AID. 2 Tablets

NON–ASPIRIN

Active ingredient (in each tablet) Purpose
Acetaminophen 325 mg . . . . . Analgesic/antipyretic

Uses
temporary relief of minor aches and pains associated with
■ common cold ■ headache ■ toothache
■ muscular aches ■ backache ■ arthritis
■ menstrual cramps ■ and reduction of fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 12 tablets in 24 hours, which is the maximum daily amount ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ for more than 10 days for pain unless directed by a doctor ■ for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin ▶

Warnings (continued)

Stop use and ask a doctor if ■ symptoms do not improve ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present ■ a rare sensitivity reaction occurs

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Do not exceed recommended dosage.**

Directions

Adults and children 12 years of age and older	Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.
Children 6-11 years of age	Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.
Children under 6 years of age	Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.

Other information ■ store at 59°-86°F (15°-30°C) ■ tamper evident sealed packets ■ do not use any open or torn packets

Inactive ingredients corn starch, hydroxypropyl methylcellulose, polyethylene glycol, pregelatinized starch, stearic acid. May contain povidone and sodium starch glycolates.

Distributed by **GENUINE FIRST AID**
600 Cleveland Str Suite 400, Clearwater, FL 33755

Active Ingredient (in each tablet) Purpose

Aspirin (NSAID*) 325 mg..... Pain Reliever / fever reducer

*nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and pains associated with:
headache ; muscular aches ; minor arthritis pain ; backache ; common cold ; toothache ; menstrual cramps ; Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox of flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, skin reddening, facial swelling, rash, asthma (wheezing), blisters, shock, If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinner (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed

Do not use: if you have ever had an allergic reaction to any other pain reliever/ fever reducer; right before or after heart surgery; if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if: stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

under a doctor's care for any serious condition; taking any other drug

When using this product: take with food or milk if stomach upset occurs

Stop use and ask a doctor if
you experience any of the following signs of stomach bleeding:
feel faint; vomit blood; have bloody or black stools; have stomach
pain that does not get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; you have difficulty swallowing; if ringing in the ears or loss of hearing occurs; redness or swelling is present in the painful areas; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or

contact a Poison Control Center right away.

Directions

do not use more than directed

the smallest effective dose should be used

drink a full glass of water with each dose

do not take longer than 10 days, unless directed by a doctor

Adults and children: (12 years and older) Take 1 or 2 tablets with

water every 4 hours as needed. Do not take more than 12 tablets in 24

hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Store at 59 - 86 degree Fahrenheit (15 - 30 degree Celsius); avoid

excessive heat and humidity; tamper evident sealed packets;

Do not use any opened or torn packets

Inactive Ingredients: hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by GENUINE FIRST AID
600 Clevelad Str Suite 400, Clearwater, FL 33755

ASPIRIN 2 Tablets

ASPIRIN

2 Tablets

Active ingredient (in each tablet) **Purpose**

Aspirin (NSAID*) 325 mg ..... Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses **Temporarily relieves minor aches and pains associated with**
■ headache ■ muscular aches ■ minor arthritis pain ■ backache
■ common cold ■ toothache ■ menstrual cramps

Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash ■ asthma (wheezing) ■ blisters ■ shock
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/ fever reducer ■ right before or after heart surgery ■ if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ▶

Warnings (continued)

Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing
■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ **do not use more than directed**
■ the smallest effective dose should be used
■ drink a full glass of water with each dose
■ do not take longer than 10 days, unless directed by a doctor

Adults and children: (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information ■ store at 59º-86ºF (15º-30ºC) ■ avoid excessive heat and humidity ■ tamper evident sealed packets ■ do not use any opened or torn packets

Inactive ingredients hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by **GENUINE FIRST AID**
600 Cleveland Str Suite 400, Clearwater, FL 33755

CARRYING CASE

1 Hard Case

1 Easy Access Pocket System

POCKET 1: SMALL CUTS and BURNS

50 Adhesive Plastic Bandages 3/4" X 3"

40 Junior Adhesive Plastic Bandages 3/8" X 1-1/2"

2 Knuckle Fabric Bandages

1 Adhesive Spot Bandages 7/8"X7/8"

1 Fingertip Fabric Bandage

3 Burn Cream

12 Antiseptic Towelettes

10 Cotton Tipped Applicators

POCKET 2: MEDIUM CUTS and SCRATCHES

6 Sterile Gauze Pads 2"X2"

2 Sterile Gauze Pads 3"X3"

1 Roller Gauze Bandage 2" X 4.1yds.

POCKET 3: SEVERE BLEEDING and BURNS

1 First Aid Tape Roll 1/2" X 5yds.

1 Combine Pad 5" X 9"

POCKET 4: CPR

1 CPR Breathing Barrier

POCKET 5: PROTECTION

2 Medical Grade Vinyl Gloves
1 Instant Cold Compress
1 Emergency Blanket 38" X 60"
2 Insect Sting Relief Pads
1 Triangular Bandage 42" X 42" X 59"
4 Ibuprofen 200mg
4 Non Aspirin 325mg
2 Aspirin 325mg

POCKET 6: INSTRUMENTS

1 Emergency First Aid Guide
2 Assorted Safety Pins

ESTUCHE PORTATIL

1 Estuche de Plastico
1 Sistema de fundas para rapido acceso

FUNDA 1: CORTADURAS Y QUEMADURAS MENORES

50 Vendas Adhesivas Plasticas 3/4" X 3"
40 Vendas Plasticas Tamano Junior 3/8" X 1-1/2"
2 Vendas de Tela para Nudillos
5 Vendas Adhesivas Redondas 7/8"X7/8"
1 Vendas de tela para yema del dedo
3 Cream para Quemaduras
12 Toallitas Antisepticas
10 Aplicadores con Puntas de Algodon

FUNDA 2: CORTADURAS Y RASGUNOS MEDIANOS

6 Almohadillas Esteril de Gasa 2"X2"
2 Almohadillas Esteril de Gasa 3"X3"
1 Vendas de Gasa en Rollo 2" X 4.1yds.

FUNDA 3: SANGRADO SEVERO Y QUEMADURAS

1 Cinta de papel para Primeros Auxilios 1/2" X 5yds.
1 Almohadilla de Uso Combinado 5" X 9"

FUNDA 4: CPR

1 Mascarilla CPR

FUNDA 5: PROTECCION

2 Guantes de Vinilo
1 Compresa de Frio Instantaneo
1 Manta Termica 38" X 60"

2 Toallitas para Picaduras de Insectos

1 Vendajes Triangulares 42" X 42" X 59"

4 Ibuprofen 200mg

4 Non Aspirin 325mg

2 Aspirin 325mg 325mg

FUNDA 6: INSTRUMENTOS

1 Manual de Primeros Auxilios
2 Imperdibles (alfileres de gancho)

Genuine First Aid LLC

814471010167

Copyright c 2010 Genuine First Aid LLC.

All rights reserved.

Manufactured in China for:

Genuine First Aid LLC.

600 Cleveland Street

Suite 400

Clearwater, FL 33755

www.GenuineFirstAid.com

GENUINE FIRST AID.



CARRYING CASE
1 Hard Case
1 Easy Access Pocket System

POCKET 1: SMALL CUTS & BURNS
50 Adhesive Plastic Bandages 3/4" x 3"
40 Junior Adhesive Plastic Bandages 3/8" x 1-1/2"
2 Knuckle Fabric Bandages
5 Adhesive Spot Bandages 7/8"x7/8"
1 Fingertip Fabric Bandage
3 Burn Cream
12 Antiseptic Towelettes
10 Cotton Tipped Applicators

POCKET 2: MEDIUM CUTS & SCRATCHES
6 Sterile Gauze Pads 2"x2"
2 Sterile Gauze Pads 3"x3"
1 Roller Gauze Bandage 2" x 4.1yds.

POCKET 3: SEVERE BLEEDING AND BURNS
1 First Aid Tape Roll 1/2" x 5yds.
1 Combine Pad 5" x 9"

POCKET 4: CPR
1 CPR Breathing Barrier

POCKET 5: PROTECTION
2 Medical Grade Vinyl Gloves
1 Instant Cold Compress
1 Emergency Blanket 38" x 60"
2 Insect Sting Relief Pads
1 Triangular Bandage 42" x 42" x 59"
4 Ibuprofen 200mg
4 Non Aspirin 325mg
2 Aspirin 325mg

POCKET 6: INSTRUMENTS
1 Emergency First Aid Guide
2 Assorted Safety Pins

ESTUCHE PORTATIL
1 Estuche de Plastico
1 Sistema de fundas para rapido acceso

FUNDA 1: CORTADURAS Y QUEMADURAS MENORES
50 Vendas Adhesivas Plasticas 3/4" x 3"
40 Vendas Plásticas Tamaño Júnior 3/8" x 1-1/2"
2 Vendas de Tela para Nudillos
5 Vendas Adhesivas Redondas 7/8"x7/8"
1 Vendas de tela para yema del dedo
3 Crema para Quemaduras
12 Toallitas Antisepticas
10 Aplicadores con Puntas de Algodon

FUNDA 2: CORTADURAS Y RASGUÑOS MEDIANOS
6 Almohadillas Esteril de Gasa 2"x2"
2 Almohadillas Esteril de Gasa 3"x3"
1 Vendas de Gasa en Rollo 2" x 4.1yds.

FUNDA 3: SANGRADO SEVERO Y QUEMADURAS
1 Cinta de Papel para Primeros Auxilios 1/2" x 5yds.
1 Almohadilla de Uso Combinado 5" x 9"

FUNDA 4: CPR
1 Mascarilla CPR

FUNDA 5: PROTECCIÓN
2 Guantes de Vinilo
1 Compresa de Frio Instantaneo
1 Manta Termica 38" x 60"
2 Toallitas para Picaduras de Insectos
1 Vendajes Triangulares 42" x 42" x 59"
4 Ibuprofen 200mg
4 Non Aspirin 325mg
2 Aspirin 325mg 325mg

FUNDA 6: INSTRUMENTOS
1 Manual de Primeros Auxilios
2 Imperdibles (alfileres de gancho)

Manufactured in China for:
Genuine First Aid LLC.
600 Cleveland Street
Suite 400
Clearwater, FL 33755
www.GenuineFirstAid.com

GENUINE FIRST AID.

Genuine First Aid LLC
814471010167

Copyright © 2010 Genuine First Aid LLC.
All rights reserved.

AUTO 157

benzalkonium chloride, lidocaine, benzocaine, alcohol, ibuprofen, acetaminophen, aspirin kit

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:52124-0118

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0118-1	1 in 1 KIT		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	3 PACKAGE	2.7 g
Part 2	12 PACKAGE	9.6 mL
Part 3	2 PACKAGE	1 mL
Part 4	2 PACKAGE	4
Part 5	2 PACKAGE	4
Part 6	1 PACKAGE	2

Part 1 of 6

GENUINE FIRST AID BURN ANTISEPTIC PAIN RELIEF WITH ALOE

benzalkonium chloride, lidocaine cream

Product Information			
Item Code (Source)	NDC:52124-0004		
Route of Administration	TOPICAL	**DEA Schedule**	

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.5 g in 100 g

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:52124-0004-1	0.9 g in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part345	05/07/2010	

Part 2 of 6

ANTISEPTIC TOWELETTE

benzalkonium chloride liquid

Product Information			
Item Code (Source)	NDC:52124-0001		
Route of Administration	TOPICAL	**DEA Schedule**	

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.40 mL in 100 mL

Inactive Ingredients	
Ingredient Name	**Strength**
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0001-1	0.8 mL in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333	05/08/2010	

Part 3 of 6

INSECT STING RELIEF PAD

benzocaine,alcohol liquid

Product Information

Item Code (Source)	NDC:52124-0008		
Route of Administration	TOPICAL	**DEA Schedule**	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	6 mL in 100 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	60 mL in 100 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0008-1	0.5 mL in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part348	05/08/2010	

Part 4 of 6

IBUPROFEN

ibuprofen tablet

Product Information

Item Code (Source)	NDC:52124-0009		
Route of Administration	ORAL	**DEA Schedule**	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE (UNII: 3NXW29V3WO)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white (White)	**Score**	no score
Shape	ROUND	**Size**	10mm
Flavor		**Imprint Code**	44;352
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0009-1	2 in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/08/2010	

Part 5 of 6

NON-ASPIRIN

acetaminophen tablet

Product Information			
Item Code (Source)	NDC:52124-0010		
Route of Administration	ORAL	**DEA Schedule**	

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients	
Ingredient Name	**Strength**
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics			
Color	white (WHITE)	**Score**	no score
Shape	ROUND	**Size**	11mm
Flavor		**Imprint Code**	AZ;234
Contains			

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:52124-0010-1	2 in 1 PACKAGE		

Marketing Information			
Marketing Category	**Application Number or Monograph Citation**	**Marketing Start Date**	**Marketing End Date**
OTC monograph final	part343	05/08/2010	

Part 6 of 6

ASPIRIN

aspirin tablet

Product Information			
Item Code (Source)	NDC:52124-0011		
Route of Administration	ORAL	**DEA Schedule**	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white (White)	**Score**	no score
Shape	ROUND	**Size**	11mm
Flavor		**Imprint Code**	44;157;ASPIRIN
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0011-1	2 in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	05/08/2010	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333	05/07/2010	

Labeler - Genuine First Aid LLC (619609857)

Establishment

Name	Address	ID/FEI	Business Operations
GFA Production (Xiamen) Co., Ltd		421256261	manufacture

Revised: 6/2010 Genuine First Aid LLC