

**IBUPROFEN- ibuprofen tablet, film coated**  
**Discount Drug Mart**

-----  
**Drug Mart 44-292**

***Active ingredient (in each brown caplet)***

Ibuprofen USP, 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - backache
  - muscular aches
  - toothache
  - menstrual cramps
  - headache
  - the common cold
  - minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- shock
- facial swelling
- blisters
- rash
- hives
- skin reddening
- asthma (wheezing)

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- are age 60 or older

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - leg swelling
  - slurred speech
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 caplet, 2 caplets may be used
  - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

***Other information***

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

***Inactive ingredients***

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

***Questions or comments?***

**Call 1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

**Principal Display Panel**

DISCOUNT

**drug  
mart**

FOOD MARKET

AN EMPLOYEE OWNERSHIP COMPANY

NDC 53943-292-15

MAY CONTAIN

**ANTI-THEFT  
DEVICE**

***IBUPROFEN***

TABLETS USP, 200 mg

*Pain Reliever / Fever Reducer (NSAID)*

Compare to the active ingredient in ADVIL® Caplets††

Actual

Size

**50 COATED CAPLETS†**

*†CAPSULE-SHAPED TABLETS*

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**SATISFACTION GUARANTEED**

IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE  
TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO  
STORE, SEND REASON FOR DISSATISFACTION, NAME,  
ADDRESS AND EMPTY PACKAGE TO:  
DISCOUNT DRUG MART  
211 COMMERCE DRIVE, MEDINA, OHIO 44256

††This product is not manufactured or distributed by  
PF Consumer Healthcare 1 LLC, owner of the registered  
trademark Advil® Caplets. 50844 REV1221B29215

**DISTRIBUTED BY: DRUG MART - FOOD FAIR  
211 COMMERCE DRIVE, MEDINA, OH 44256  
[www.discount-drugmart.com](http://www.discount-drugmart.com)**

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Drug Facts**  
**Active Ingredient (in each brown caplet)**  
ibuprofen USP, 200 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever/fever reducer

**Uses**  
• temporarily relieves minor aches and pains due to:  
• menstrual cramps  
• toothache  
• the common cold  
• headache  
• backache  
• minor pain of arthritis  
• temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
• rash  
• skin redness  
• asthma (wheezing)  
• facial swelling  
• shock  
• blisters  
• hives  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
• take more or for a longer time than directed  
• take a blood thinning (anticoagulant) or steroid drug  
• are age 60 or older  
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
• have had stomach ulcers or bleeding problems  
• have 3 or more alcoholic drinks every day while using this product

**Drug Facts (continued)**  
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Do not use**  
• if you have ever had an allergic reaction to any other pain reliever/fever reducer  
• right before or after heart surgery  
**Ask a doctor before use if**  
• you have a history of stomach problems, such as heartburn  
• you are taking a diuretic  
• you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
• stomach bleeding warning applies to you  
• you have problems or serious side effects from taking pain relievers or fever reducers  
**Ask a doctor or pharmacist before use if you are**  
• under a doctor's care for any serious condition  
• taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit or aspirin  
• taking any other drug  
**When using this product**  
• take with food or milk if stomach upset occurs  
**Stop use and ask a doctor if**  
• you experience any of the following signs of stomach bleeding:  
• feel faint  
• vomit blood  
• have bloody or black stools  
• have stomach pain that does not get better  
• you have symptoms of heart problems or stroke:  
• chest pain  
• leg swelling  
• trouble breathing  
• slurred speech  
• weakness in one part or side of body

**Drug Facts (continued)**  
glycolate, stearic acid, titanium dioxide  
polyethylene glycol, red iron oxide, sodium starch  
microcrystalline cellulose, polydextrose  
**Questions or comments?**  
Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday  
†This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil® Caplets. 50844 REV1221B29215  
DISTRIBUTED BY: DRUG MART - FOOD FAIR 211 COMMERCIAL DRIVE, MEDINA, OH 44256  
www.discount-drugmart.com

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**SATISFACTION GUARANTEED**  
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO:  
DISCOUNT DRUG MART  
211 COMMERCIAL DRIVE, MEDINA, OHIO 44256

**No print/No varnish Lot & Exp date**

0 93351 02767 3

**drug mart**  
FOOD MARKET  
ANTHI-THIEF  
MAY CONTAIN DEVICE

**IBUPROFEN**  
TABLETS USP, 200 mg  
Pain Reliever / Fever Reducer (NSAID)  
Compare to the active ingredient in ADVIL® Caplets††

**Actual Size**

**50 COATED CAPLETS\***  
\*CAPSULE-SHAPED TABLETS

NDC 53943-292-15

**Drug Facts (continued)**  
• pain gets worse or lasts more than 10 days  
• fever gets worse or lasts more than 3 days  
• redness or swelling is present in the painful area  
• any new symptoms appear  
**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.  
**Directions**  
• do not take more than directed  
• the smallest effective dose should be used  
• adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist  
• if pain or fever does not respond to 1 caplet, 2 caplets may be used  
• do not exceed 6 caplets in 24 hours, unless directed by a doctor  
• children under 12 years: ask a doctor  
**Other information**  
• store between 20°-25°C (68°-77°F)  
• avoid excessive heat 40°C (104°F)  
• see end flap for expiration date and lot number  
**Inactive ingredients** carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate,

## Drug Mart 44-292

### IBUPROFEN

ibuprofen tablet, film coated

#### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-292
Route of Administration	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDWL1A)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	44;292
<b>Contains</b>			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-292-15	1 in 1 CARTON	05/24/1988	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:53943-292-12	1 in 1 CARTON	05/24/1988	
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:53943-292-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(53943-292)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(53943-292)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(53943-292)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(53943-292)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(53943-292)

Revised: 3/2024

Discount Drug Mart