

GLYCERIN - glycerin suppository
Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DRUG FACTS

ACTIVE INGREDIENT

Glycerin, USP 80.7%

PURPOSE

Laxative

USE

For relief of occasional constipation

WARNINGS

For rectal use only. May cause rectal discomfort or a burning sensation.

Do not use

- more than one per day
- for a period of longer than one week unless directed by a doctor
- laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor
- if seal under product lid is damaged, missing or broken.

Ask a doctor

- before use if you have noticed a sudden change in bowel habits that persists over a period of two weeks
- if you are pregnant or nursing a baby

Stop use and ask a doctor

if you have rectal bleeding or fail to have a bowel movement after using a laxative. This may indicate a serious condition.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- Children 2 - 6 years of age: carefully insert one suppository well into the rectum
- Do not use more than once per day
- Generally produces a bowel movement in 15 minutes to 1 hour
- Children under 2 years of age: ask a doctor.

OTHER INFORMATION

- Store container tightly closed
- Store at room temperature: 15°- 30° C (59° - 86° F). Do not exceed 30° C (86° F).

INACTIVE INGREDIENTS

sodium stearate, purified water

PACKAGE LABEL

NDC 41250-404-12

meijer

**children's
Glycerin
Suppositories**

laxative

For fast, gentle relief of
occasional constipation

Compare to Fleet®
active ingredient*

Meijer Pharmacist Recommended

12 Suppositories

DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544

*This product is not manufactured or
distributed by C.B.Fleet co., owner of
the registered trademark Fleet®.

USE ONLY IF PRINTED
INNER SEAL IS NOT
BROKEN OR MISSING

NDC 41250-404-12



meijer
children's
Glycerin
Suppositories

laxative

FOR FAST, GENTLE RELIEF OF
OCCASIONAL CONSTIPATION

COMPARE TO FLEET®
active ingredient*

12 Suppositories



CODE AREA

Drug Facts	Rev. A	Purpose	Laxative
Active ingredient (per suppository)			Glycerin, USP 80.7%
Use			For relief of occasional constipation
Warnings			For rectal use only. May cause rectal discomfort or a burning sensation. Do not use more than one per day for a period longer than one week unless directed by a doctor. Laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor. If seal under product lid is damaged, missing, or broken. Ask a doctor before use if you have noticed a sudden change in bowel habits that persists over a period of two weeks. If you are pregnant or nursing a baby. Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after using a laxative. This may indicate a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions			Children 2 - 6 years of age: carefully insert one suppository well into the rectum. Do not use more than once per day. Generally produces a bowel movement in 15 minutes to 1 hour. Children under 2 years of age: ask a doctor
Other information			Store container tightly closed. Store at room temperature: 15° - 30° C (59° - 86° F). Do not exceed 30° C (86° F).
Inactive ingredients			sodium stearate, purified water


 DIST. BY MEIJER DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49544
*This product is not manufactured or distributed by C.B. Fleet Co., owner of the registered trademark Fleet®.


 7 19283 59237 8
 USE ONLY IF PRINTED
 INNER SEAL IS NOT
 BROKEN OR MISSING.

GLYCERIN

glycerin suppository

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-404
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
glycerin (UNII: PDC6A3C0OX) (glycerin - UNII:PDC6A3C0OX)	glycerin	1.3 g

Inactive Ingredients

Ingredient Name	Strength
sodium stearate (UNII: QU7E2XA9TG)	
water (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-404-12	12 in 1 JAR		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part334	02/01/2006	

Labeler - Meijer Distribution Inc (006959555)

