RANITIDINE- ranitidine hydrochloride tablet Shopko Stores Operating Co., LLC

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness**, **sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.
- store at 20° 25° C (68° 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

QUESTIONS?

Call 1-800-406-7984

PRINCIPAL DISPLAY PANEL

SHOPKO[®]

Compare to the Active Ingredient of Zantac 150®

Maximum Strength

Acid Reducer

Ranitidine Tablets, USP 150 mg

Acid Reducer

Prevents & Relieves: Heartburn Associated with Acid Indigestion & Sour Stomach

50 Tablets

Packaged by: Ohm Laboratories Inc.

5093015/0212



Maximum Strength

Acid Reducer

Ranitidine Tablets, USP 150 mg

SHOPKO

Ingredient of Zantac 150®

Maximum Strength

Acid Reducer

Ranitidine Tablets, USP 150 mg

Acid Reducer

Prevents & Relieves: Heartburn Associated with Acid Indigestion & Sour Stomach



50 Tablets

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

Batch No.

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

- This product is sodium and sugar free
 Toole at 20° 25° C (68° 77° F)
 Stool at 20° 25° C (68° 77° F)
 Rovid excessive heat or furnibity
 Rovid excessive heat or furnibity Other information

 - can be used up to twice daily (do not take more than 2 tablets in 24 hours) 100d of drinking beverages that cause heartburn
 - to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating

Drug Facts (continued)

adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water

DIRECTIONS

ceep out of reach of children. In case of overdose, get medical help or contact a Polson Control Center right away It pregnant or breast-feeding, ask a health professional before use.

 your heartburn continues or worsens
 you need to take this product for more than 14 days Stop use and ask a doctor if

- cheat pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightnesdedness
 - had heartburn over 3 months. This may be a sign of a more serious condition
 heartburn with lightheadedness, sweating or dizziness
 - stomach pain


 - In frequent cheet pain
 In frequent wheezing, particularly with heartburn
 In nexylained weight loss
 In nesusea or vomiting
 In the paint of th

Ask a doctor before use if you have

- If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor me with other exid reducers
 If you have kidney disease, except under the advice and supervision of a doctor. Do not use

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers Warnings

■ prevents hearthurn associated with acid indigestion and sour stomach brought on by certain foods and beverages



Packaged by: Ohm Laboratories Inc. 1385 Livingston Avenue North Brunswick, NJ 08902

This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals Inc., owner of the registered trademark Zantac 150®



311	Uses Institute associated with soid indinestion and sour stomach
Purpose Reducer	Active ingredient (in each tablet) Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)
	Drug Facts

RANITIDINE

ranitidine tablet

Product TypeHUMAN OTC DRUG LABELItem Code (Source)NDC:370 12-351

Route of Administration ORAL DEA Schedule

Active Ingredient/Active Moiety

Ingredient NameBasis of StrengthStrengthRANITIDINE HYDRO CHLO RIDE (RANITIDINE)RANITIDINE150 mg

Inactive Ingredients

Ingredient Name	Strength
COLLO IDAL SILICON DIO XIDE	
CROSCARMELLOSE SODIUM	
HYPROMELLOSES	
FERRIC O XIDE RED	
MAGNESIUM STEARATE	
CELLULO SE, MICRO CRYSTALLINE	
POLYETHYLENE GLYCOLS	

TITANIUM DIO XIDE

TALC

Product Characteristics

Color	white (Off-White to Pale Yellow)	Score	no score
Shape	OVAL	Size	7mm
Flavor		Imprint Code	9 R
Contains			

Packaging

ı	ue.mg.mg			
ı	# Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1 NDC:37012-351-50	50 in 1 BOTTLE		

Marketing Category Application Number or Monograph Citation Marketing Start Date Marketing End Date

ANDA	ANDA200536	03/30/2012

Labeler - Shopko Stores Operating Co., LLC (023252638)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment				
Name	Address	ID/FEI	Business Operations	
Shasun Pharmaceuticals Limited		915786829	manufacture(37012-351)	

Revised: 2/2013 Shopko Stores Operating Co., LLC