CAREONE MILK OF MAGNESIA- magnesium hydroxide suspension American Sales Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

American Sales Company Milk of Magnesia Drug Facts

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredient in Phillips'® Milk of Magnesia

MILK OF MAGNESIA

Magnesium Hydroxide 1200 mg per 15 mL

Saline Laxative

Wild Cherry Flavor

Comfortable, Cramp Free Relief

Stimulant Free

Low Sodium

Gluten Free

ALCOHOL 0.4%

OUR PHARMACISTS RECOMMEND

12 FL OZ (355mL)



Drug Facts DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Active ingredient (in each 15 mL)

Purpose

Magnesium hydroxide 1200 mg..

Saline laxative

Uses ■ relieves occasional constipation (irregularity)

quenerally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease a magnesium-restricted diet stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

■ you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided ■ mL = milliliter adults and children 12 years and older 30 mL to 60 mL children 6 to 11 years 15 mL to 30 mL children under 6 years ask a doctor

Other information

- each 15 mL contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F) close cap tightly after use avoid freezing

Inactive ingredients anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments? 1-800-719-9260

01616 유 В

*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips'® Milk of Magnesia.

DISTRIBUTED BY FOODHOLD U.S.A., LLC LANDOVER, MD 20785 ©2017 S&S Brands, LLC



CAREONE MILK OF MAGNESIA

magnesium hydroxide suspension

Product Information

HUMAN OTC DRUG NDC:41520-949 Product Type Item Code (Source)

Route of Administration ORAL

Active Ingredient/Active Moiety

	Ingredient Name	Basis of Strength	Strength
l	MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6 V3LHY838,	MAGNESIUM	1200 mg
l	HYDRO XIDE ION - UNII:9159 UV381P)	HYDRO XIDE	in 15 mL

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)		
GLYCERIN (UNII: PDC6A3C0OX)		
WATER (UNII: 059QF0KO0R)		
SORBITOL (UNII: 506T60A25R)		
SUCRO SE (UNII: C151H8 M554)		
XANTHAN GUM (UNII: TTV12P4NEE)		
D&C RED NO. 28 (UNII: 767IP0 Y5NH)		

Product Characteristics			
Color	PINK (Opaque Light pink)	Score	
Shape		Size	
Flavor	CHERRY (wild cherry)	Imprint Code	
Contains			

ı	Packaging				
ı	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1 NI	DC:41520-949-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/22/2017	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	08/26/2012	

Labeler - American Sales Company (809183973)

Revised: 10/2017 American Sales Company