GOOD NEIGHBOR PHARMACY IBUPROFEN COLD AND SINUS- ibuprofen, pseudoephedrine hydrochloride tablet Dispensing Solutions, Inc.

Amerisource Bergen Ibuprofen Cold and Sinus Drug Facts

Active ingredient (in each caplet)

Ibuprofen, USP 200 mg (NSAID)*

Pseudoephedrine HCl, USP 30 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Nasal decongestant

Uses

temporarily relieves these symptoms associated with the common cold or flu:

- headache
- fever
- sinus pressure
- nasal congestion
- minor body aches and pains

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs, [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 12 years of age
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, or trouble urinating due to an enlarged prostate gland
- vou are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains pseudoephedrine, or any other nasal decongestant
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- symptoms continue or get worse
- any new symptoms appear
- you get nervous, dizzy, or sleepless
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- redness or swelling is present in the painful area

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

• do not take more than directed

- the smallest effective dose should be used
- adults and children 12 years of age and over:
- take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used.
- do not use more than 6 caplets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
- read all warnings and directions before use. Keep carton.

Inactive ingredients

acacia, calcium carbonate, carnauba wax, confectioner's sugar, corn starch, croscarmellose sodium, crospovidone, FD&C blue no. 2, FD&C red no. 40, FD&C yellow no. 6, gelatin, guar gum, hydrogenated vegetable oil, hydroxypropyl cellulose, kaolin, pharmaceutical ink, polyethylene glycol, powdered cellulose, povidone, pregelatinized starch, silicon dioxide, sodium benzoate, sucrose, talc, titanium dioxide, white wax

Questions or comments?

1-800-719-9260

Principal Display Panel

COMPARE TO ACTIVE INGREDIENTS IN ADVIL® COLD & SINUS

Non Drowsy

Ibuprofen Cold & Sinus

Ibuprofen 200 mg – Pain Reliever/Fever Reducer (NSAID)*

Pseudoephedrine HCl 30 mg – Nasal Decongestant

Tablets, USP

*nonsteroidal anti-inflammatory drug

Relieves Sinus Pressure, Nasal Congestion and Fever

See New Warnings Information

**OVAL-SHAPED TABLETS

NDC 68258-8902-02

BULK SOURCE DATA

DIST. BY: AMERISOURCEBERGEN
CHESTERBROOK, PA 19087
PRODUCT ID:
WHITE / BLUE / LIGHT BLUE / RED /
YELLOW / DARK RASPBERRY / PINK /
LIGHT BROWN / BROWN / BLACK
BOX PRINTED NDC 24385-465-60
BULK SOURCE NDC: 24385-0465-60
MFR. LOT: XXXXXX
PEDIGREE #: 712810569
DISPENSE IN THIS
TIGHT/LIGHT RESISTANT CONTAINER

TAKE___CAPLET(S)
ORALLY EVERY __HOURS
__TIMES A DAY. DO NOT
USE MORE THAN 6 CAPLETS
IN ANY 24-HOUR PERIOD
UNLESS DIRECTED BY A
DOCTOR. TAKE WITH FOOD.
AVOID ALCOHOL.



IBUPROFEN/PSE HCI COLD&SINUS

(NON-DROWSY)

20 (200 mg/30 mg) CAPLETS NDC 68258-8902-02 PRODUCT #810220

ACTIVE INGREDIENT (IN EACH COATED CAPLET):
IBUPROFEN USP....200 mg
(PURPOSE: PAIN RELIEVER/FEVER REDUCER)
PSEUDOEPHEDRINE HYDROCHLORIDE USP....30 mg
(PURPOSE: NASAL DECONGESTANT)
FD&C YELLOW No. 6
SEE BOX FOR WARNINGS INFORMATION.

COMPARE TO ACTIVE INGREDIENTS IN ADVIL® COLD & SINUS CAPLETS LOT#SAMPLE EXP: 00-00 Rx # 32692018

WARNING: KEEP OUT OF CHILDREN'S REACH STORE AT 68°- 77° F. SEE USP.

810220 NDC 68258-8902-02 IBUPROFEN/PSE HCI COLD&SINUS 20 (200 mg/30 mg) CAPLETS LOT # SAMPLE EXP: 00-00 MN 24385-0465-60 RX# 32692018

810220 NDC 68258-8902-02 IBUPROFEN/PSE HCI COLD&SINUS 20 (200 mg/30 mg) CAPLETS LOT # SAMPLE EXP: 00-00 MN 24385-0465-60 RX# 32692018

810220 NDC 68258-8902-02 IBUPROFEN/PSE HCI COLD&SINUS 20 (200 mg/30 mg) CAPLETS LOT # SAMPLE EXP: 00-00 MN 24385-0465-60 RX# 32692018





GOOD NEIGHBOR PHARMACY IBUPROFEN COLD AND SINUS

05/13

Date:

ibuprofen, pseudoephedrine hydrochloride tablet

Product Information

Product TypeHUMAN OTC DRUGItem Code (Source)NDC:68258-8902(NDC:24385-465)Route of AdministrationORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg	
PSEUDO EPHEDRINE HYDRO CHLO RIDE (UNII: 6 V9 V2RYJ8 N) (PSEUDO EPHEDRINE - UNII:7CUC9 DDI9 F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg	

Inactive Ingredients				
Ingredient Name	Strength			
ACACIA (UNII: 5C5403N26O)				
CALCIUM CARBONATE (UNII: H0 G9 379 FGK)				
CARNAUBA WAX (UNII: R12CBM0 EIZ)				
SUCROSE (UNII: C151H8M554)				
STARCH, CORN (UNII: O8232NY3SJ)				
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)				
CROSPOVIDONE (UNII: 68401960 MK)				
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
GELATIN (UNII: 2G86QN327L)				
GUAR GUM (UNII: E89 I16 37 KE)				
HYDRO XYPRO PYL CELLULO SE (UNII: RFW2ET671P)				
KAOLIN (UNII: 24H4NWX5CO)				

POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
PO VIDO NE (UNII: FZ989 GH94E)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	
WHITE WAX (UNII: 7G1J5DA97F)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	

Product Characteristics			
Color	BROWN	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	083
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68258-8902-2	2 in 1 CARTON		
1		10 in 1 BLISTER PACK		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074567	10/15/2001	

Labeler - Dispensing Solutions, Inc. (066070785)

Registrant - PSS World Medical, Inc. (101822682)

Establishment				
Name	Address	ID/FEI	Business Operations	
Dispensing Solutions, Inc.		066070785	relabel(68258-8902), repack(68258-8902)	

Revised: 6/2013 Dispensing Solutions, Inc.