

FIRST AID SHOT THERAPY HANGOVER RELIEF- choline salicylate and caffeine solution
First Aid Beverages, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

First Aid Shot Therapy® Hangover Relief

Drug Facts

Active ingredients	Purposes (in each bottle)
Choline salicylate 870mg (NSAID)*	Pain reliever
Caffeine 65mg	Pain reliever aid

* nonsteroidal anti-inflammatory drug

Uses

- for temporary relief of minor aches and pain associated with a hangover
- helps restore mental alertness or wakefulness when experiencing fatigue or drowsiness associated with a hangover
- also for temporary relief of headaches or body aches and pains alone

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Contains salicylate

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

When using this product limit use of caffeine containing drugs, foods, or drinks, because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat

- for occasional use only

Stop use and ask a doctor if

- an allergic reaction occurs (seek medical help right away)
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- you experience any of the following signs of stomach bleeding
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

For adults and children 12 years or older:

- swallow the contents of one bottle
- repeat every 4 hours while symptoms persist
- do not take more than 6 bottles in 24 hours
- children under 12 years: do not use

Other Information

- Keep from freezing
- Avoid excessive heat (over 104°F or 40°C)

Inactive Ingredients

Citric acid, natural flavors, purified water, sodium benzoate, sodium citrate, sucralose

Questions or comments?

1-877-958-8331 (Monday-Friday 9AM-5PM PST)

PRINCIPAL DISPLAY PANEL - 40 mL Bottle Label

•

**HANG
OVER**

**FAST
– LIQUID –
RELIEF®**

**FIRST AID
SHOT THERAPY®**

CITRUS

choline salicylate & caffeine
pain reliever
(NSAID)/pain reliever aid

1.35 FL OZ (40mL)

FDA
COMPLIANT

HANG OVER • HANG OVER • HANG OVER

Do not use if seal is broken

We hope F.A.S.T. relieved your hangover pain.

For more info visit:
www.firstaidshottherapy.com



FAST
-LIQUID-
RELIEF[®]



FIRST AID
SHOT THERAPY[®]

CITRUS

choline salicylate & caffeine
pain reliever
(NSAID)/pain reliever aid

FDA
COMPLIANT

1.35 FL OZ (40mL)

Single Dose Treats:

- Hangover Pain
- Headache
- Muscle Ache

1003-004-002

2456966-92

<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ an allergic reaction occurs (seek medical help right away) ■ ringing in the ears or loss of hearing occurs ■ pain gets worse or lasts for more than 10 days ■ fever lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <p>For adults and children 12 years and older:</p> <ul style="list-style-type: none"> ■ swallow the contents of one bottle ■ repeat every 4 hours while symptoms persist ■ do not take more than 6 bottles in 24 hours ■ children under 12 years: do not use <p>Other Information</p> <ul style="list-style-type: none"> ■ keep from freezing ■ avoid excessive heat (over 104°F or 40°C) <p>Inactive Ingredients</p> <p>citric acid, natural flavors, purified water, sodium benzoate, sodium citrate, sucralose</p> <p>Questions or comments? 1-877-958-8331 (Monday-Friday 9AM-5PM PST).</p>	<p>Drug Facts (continued)</p> <p>You may also report serious side effects to this number</p> <p>Distributed by First Aid Shot Therapy[®] Burlingame, CA 94010</p> <p>First Aid Shot Therapy[®] and cross symbol are registered trademarks of First Aid Shot Therapy[®]. www.firstaidshottherapy.com</p> <p><small>U.S. Patent #3,773,434</small></p>
--	--	---

2456966-92

Drug Facts

Active ingredients Purpose (in each bottle)

Choline salicylate 670mg (NSAID)*... Pain reliever

Caffeine 65mg... Pain reliever aid

*nonsteroidal anti-inflammatory drug

1004-004-000



8 57596 00413 0

LIFT PANEL FOR MORE DRUG FACTS INFORMATION. READ ALL WARNINGS UNDER LABEL BEFORE USE.

2456966-92

<p>Drug Facts (continued)</p> <p>Uses</p> <ul style="list-style-type: none"> ■ for temporary relief of minor aches and pain associated with a hangover ■ helps restore mental alertness or wakefulness when experiencing fatigue or drowsiness associated with a hangover ■ also for the temporary relief of headaches or body aches and pains alone <p>Warnings</p> <p>Reye's syndrome: Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Contains salicylate</p>	<p>Drug Facts (continued)</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed <p>Do not use if you are</p> <ul style="list-style-type: none"> ■ allergic to salicylates (including aspirin) ■ taking other salicylate products 	<p>Drug Facts (continued)</p> <p>Ask a doctor before use if:</p> <ul style="list-style-type: none"> ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for</p> <ul style="list-style-type: none"> ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis <p>When using this product limit use of caffeine containing drugs, foods, or drinks, because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat</p> <p>■ for occasional use only</p>
---	--	---

FIRST AID SHOT THERAPY HANGOVER RELIEF
choline salicylate and caffeine solution

Product Information	Product Type	Item Code (Source)	NDC:578 15-041
	HUMAN OTC DRUG		

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Choline Salicylate (UNII: KD510K1IQW) (Salicylic Acid - UNII:O414PZ4LPZ)	Choline Salicylate	870 mg in 40 mL
Caffeine (UNII: 3G6A5W338E) (Caffeine - UNII:3G6A5W338E)	Caffeine	65 mg in 40 mL

Inactive Ingredients

Ingredient Name	Strength
Anhydrous Citric Acid (UNII: XF417D3PSL)	
Sodium Benzoate (UNII: OJ245FE5EU)	
Sodium Citrate (UNII: 1Q73Q2JULR)	
Sucralose (UNII: 96K6UQ3ZD4)	
Water (UNII: 059QF0K00R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CITRUS	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57815-041-01	40 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	02/01/2015	

Labeler - First Aid Beverages, Inc. (078618157)

Revised: 6/2015

First Aid Beverages, Inc.