

NICOTINE POLACRILEX COATED CINNAMON- nicotine polacrilex gum, chewing
CVS Pharmacy

Nicotine Polacrilex Gum 2 mg and 4 mg Coated Cinnamon - CVS

Drug Facts

Active ingredient (in each chewing piece) - 2 mg

Nicotine Polacrilex 2 mg (nicotine)

Active ingredient (in each chewing piece) - 4 mg

Nicotine Polacrilex 4 mg (nicotine)

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may

need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- oral blistering occurs
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions - 2 mg

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Directions - 4 mg

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information - 2 mg

- **each piece contains:** calcium 115 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Other information - 4 mg

- **each piece contains:** calcium 115 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients - 2 mg

acacia, acesulfame potassium, butylated hydroxytoluene, carnauba wax, FD&C red #40, gum base, hydroxypropyl cellulose, hypromellose, natural and artificial cinnamon flavors, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide and xylitol.

Inactive ingredients - 4 mg

acacia, acesulfame potassium, butylated hydroxytoluene, carnauba wax, FD&C red #40, gum base, hydroxypropyl cellulose, hypromellose, natural and artificial cinnamon flavors, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide and xylitol.

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal display panel 2 mg

Compare to the active ingredient in Nicorette® Gum*

Sugar Free

Nicotine Gum

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARATTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg.

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING

1. You must be motivated to quit.
2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. Use with a support program as directed in the enclosed User's Guide.

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE INDIVIDUAL SEALS ARE OPEN OR TORN.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

Package label 2 mg

minute clinic
START TO STOP
CAN HELP YOU QUIT
 CVS.com/StartToStop

Visit MinuteClinic® for personal counseling with a licensed provider.

- Our program includes:**
- 1-on-1 consultation
 - Personalized plan & education
 - Ongoing coaching & support
 - Nicotine-replacement recommendations or prescriptions

CVS Health. Compare to the active ingredient in Nicorette® Gum*

CVS Health. Compare to the active ingredient in Nicorette® Gum*

NDC 69842-026-44

Sugar Free Nicotine Gum

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

2 mg

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Gum USP, 4 mg.

NDC 69842-026-44

Sugar Free Nicotine Gum

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

2 mg

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Gum USP, 4 mg.



Coated Cinnamon Flavor Actual Size

100 PIECES, 2 mg EACH

Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2021 CVS/pharmacy
 CVS.com® 1-800-SHOP CVS
 V-36037

CVS Quality
 Money Back Guarantee

#718198

0 50428 16534 8

PLD-8613A FDX06592

Drug Facts

Active ingredient (in each chewing piece)
 Nicotine polacrilex 2 mg (nicotine) Stop smoking aid

Use
 ■ reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings
 ■ If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Do not stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have
 ■ a sodium-sulfite drug
 ■ heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
 ■ high blood pressure not controlled with medication. Nicotine can increase blood pressure.
 ■ diabetes, ulcer or diabetes
 ■ history of seizures

Ask a doctor or pharmacist before use if you are
 ■ using a non-nicotine stop smoking drug
 ■ taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if
 ■ mouth, teeth, or jaw problems occur
 ■ irregular heartbeat or palpitations occur
 ■ you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
 ■ oral blistering occurs
 ■ you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. They used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222), right away.

Directions
 ■ If you are under 18 years of age, ask a doctor before use
 ■ before using this product, read the enclosed User's Guide for complete directions and other important information
 ■ begin using the gum on your quit day
 ■ if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
 ■ if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 8	Weeks 9 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

Drug Facts (continued)

■ nicotine gum is a medicine and must be used a certain way to get the best results
 ■ chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
 ■ repeat this process until most of the tingle is gone (about 30 minutes)
 ■ do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
 ■ to improve your chances of quitting, use at least 8 pieces per day for the first 6 weeks
 ■ if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you nausea, heartburn, rashes or other side effects.
 ■ do not use more than 24 pieces a day
 ■ it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information
 ■ each piece contains calcium 115 mg, sodium 11 mg
 ■ store at 20°-25°C (68°-77°F)
 ■ protect from light.

Inactive ingredients saccharin, xanthine potassium, bisphenol hydrophthalate, carmellose waxy, FD&C red #40, gum base, hydroxypropyl cellulose, hydroxyethylcellulose, natural and artificial cinnamon flavors, sodium bicarbonate, sodium carbonate, saccharin, lact, titanium dioxide, xylitol

Questions or comments?
 Call 1-877-763-3636 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

- TO INCREASE YOUR SUCCESS IN QUITTING:**
1. You must be motivated to quit.
 2. Use Enough – Chew at least 8 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
 3. Use Long Enough – Use Nicotine Polacrilex Gum for the full 12 weeks.
 4. Use with a support program as directed in the enclosed User's Guide.



To remove the gum, tear off single unit. Peel off backing starting at corner with loose edge. Push gum through foil.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

CVS HEALTH Sugar Free Nicotine Gum Cinnamon Flavor

Principal display panel 4 mg

Compare to the active ingredient in Nicorette® Gum*

Sugar Free

Nicotine Gum

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg.

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. Use with a support program as directed in the enclosed User's Guide.

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

Package label 4 mg

minute clinic
START TO STOP
CAN HELP YOU QUIT
CVS.com/StartToStop

Visit MinuteClinic® for personal counseling with a licensed provider.

Our program includes:

- 1-on-1 consultation
- Personalized plan & education
- Ongoing coaching & support
- Nicotine-replacement recommendations or prescriptions

CVS Health. Compare to the active ingredient in Nicorette® Gum*

CVS Health. Compare to the active ingredient in Nicorette® Gum*

Drug Facts

Active ingredient (in each chewing piece)	Purpose
Nicotine polacrilex.....4 mg (nicotine).....	Stop smoking aid

Use

- relieves withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

- If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.
- Ask a doctor before use if you have:
 - a sodium-restricted diet
 - heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
 - high blood pressure not controlled with medication. Nicotine can increase blood pressure.
 - stomach ulcer or diabetes
 - history of seizures

Talk to a doctor or pharmacist before use if you are:

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if:

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you get dizziness or lightheadedness
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to harm children and pets. Keep used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- If you are under 18 years of age, ask a doctor before use
- Before using this product, read the enclosed User's Guide for complete directions and other important information
- Begin using the gum on your quit day
- If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum
- If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 11	Weeks 12 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

Drug Facts (continued)

- Nicotine gum is a medicine and must be used a certain way to get the best results
- Chew the gum slowly until it begins to melt. Then push it between your cheek and gum. When the triangle is gone, begin chewing again, until the triangle returns
- Repeat this process until most of the triangle is gone (about 30 minutes)
- Do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- To improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- If you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you dizziness, headache, nausea or other side effects
- Do not use more than 24 pieces a day
- It is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- Each piece contains: calcium 115 mg, sodium 11 mg
- Store at 20°-25°C (68°-77°F)
- Protect from light

Inactive ingredients: xanthan gum, croscarmellose potassium, butylated hydroxytoluene, carboxymethyl cellulose, FD&C red #40, gum base, hydroxypropyl cellulose, hydroxypropyl methylcellulose, natural and artificial cinnamon flavor, sodium bicarbonate, sodium carbonate, saccharin, lact, titanium dioxide, xylitol

Questions or comments?
 Call 1-877-755-3888 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. Use with a support program as directed in the enclosed User's Guide.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

100 PIECES, 4 mg EACH

Coated Cinnamon Flavor Actual Size

CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2021 CVS/pharmacy
 CVS.com* 1-800-SHOP CVS
 V-36037

CVS Quality
 Money Back Guarantee

#718184

PLD-8620A F000665

CVSHEALTH Sugar Free Nicotine Gum Cinnamon Flavor

NICOTINE POLACRILEX COATED CINNAMON

nicotine polacrilex gum, chewing

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-026
Route of Administration	BUCCAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N260)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XYLITOL (UNII: VCQ006KQ1E)	

Product Characteristics

Color	pink	Score	no score
Shape	SQUARE	Size	14mm
Flavor	CINNAMON	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-026-44	100 in 1 CARTON	01/01/2014	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
---------------------------	---	-----------------------------	---------------------------

ANDA	ANDA079216	01/01/2014	
------	------------	------------	--

NICOTINE POLACRILEX COATED CINNAMON

nicotine polacrilex gum, chewing

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-028
Route of Administration	BUCCAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XYLITOL (UNII: VCQ006KQ1E)	

Product Characteristics

Color	pink	Score	no score
Shape	SQUARE	Size	14mm
Flavor	CINNAMON	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-028-44	100 in 1 CARTON	01/01/2014	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079219	01/01/2014	

Labeler - CVS Pharmacy (062312574)

Revised: 11/2022

CVS Pharmacy