

GENTAMED- gentamicin sulfate powder
Bimeda, Inc.

DESCRIPTION: Each gram of GentaMed™ Soluble Powder contains gentamicin sulfate equivalent to 333.33 mg gentamicin base.

RESIDUE WARNING: For use in swine drinking water only. Swine treated with the recommended dose of GentaMed™ Soluble Powder must not be slaughtered for food for at least ten (10) days following the last treatment.

INDICATIONS: GentaMed™ Soluble Powder is recommended for the control and treatment of colibacillosis in weanling swine caused by strains of *E. coli* sensitive to gentamicin and for the control and treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

DOSAGE AND ADMINISTRATION:

GentaMed Soluble Powder is available in a 360 gram jar. The 360 gram jar contains a scoop that provides approximately 18 grams of product when level full.

For Control and treatment of Colibacillosis-Administer GentaMed Soluble Powder in drinking water at the recommended level of 25 mg per gallon (1 scoop/240 gallons) for three consecutive days.

For Control and treatment of Swine Dysentery-Administer GentaMed Soluble Powder in drinking water at the recommended level of 50 mg per gallon (1 scoop/120 gallons) for three consecutive days.

The concentration of the medication should be adjusted in extremely hot or cold weather to ensure a gentamicin dosage of approximately 0.5 mg/lb/day for 3 days for colibacillosis or 1.0 mg/lb/day for 3 days for swine dysentery. A 50-lb pig will, under average conditions, consume 0.9 gallons of water per day.

For swine dysentery, if the condition recurs, the medication may be repeated. Because of the tendency for the disease to recur on a premise with a history of swine dysentery, a control program should be used following treatment.

DIRECTIONS FOR USE:

To prepare medicated water, mix GentaMed Soluble Powder and drinking water according to the following tables:

Proportioner Use:

For Control and Treatment of Colibacillosis

Add 18 g (1 scoop) GentaMed Soluble Powder to 2 gallons of water to make stock solution and dispense at the rate of one ounce per gallon of drinking water. This will result in drinking water that contains 25 mg/gallon.

For Control and Treatment of Swine Dysentery

Add 18 g (1 scoop) GentaMed Soluble Powder to 1 gallon of water to make stock solution and dispense at the rate of one ounce per gallon of drinking water. This will result in drinking water that contains 50 mg/gallon.

Bulk Preparation:

For Control and Treatment of Colibacillosis

Add 18 g (1 scoop) GentaMed Soluble Powder to 240 gallons of water.

For Control and Treatment of Swine Dysentery

Add 18 g (1 scoop) GentaMed Soluble powder to 120 gallons of water.

Medicated drinking water should not be stored or offered in rusty containers since the drug is quickly destroyed in such containers.

Medicated water should be prepared daily and be the sole source of drinking water for 3 consecutive days.

The daily water consumption figures shown below are approximations for mild temperatures and are presented as a guide only. Actual water consumption varies with environmental temperature, humidity, and diet.

Approximate Daily Water Consumption

Weight of Pig(lb)	Gallon for 5 Pigs	Gallons for 10 Pigs
25	3.0	6.0
50	4.5	9.0
75	6.0	12.0
100	7.5	15.0

SIDE EFFECTS: There have been no reported side effects following the recommended use of GentaMed™ Soluble Powder in swine.

CONTRAINDICATIONS: There are no known contraindications to this drug when used as directed.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

PRECAUTIONS: Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).

HOW SUPPLIED: GentaMed™ Soluble Powder is available in a 360 gram jar with 12 jars per shipper.

GentaMed™ Soluble Powder

(Gentamicin Sulfate)

Contains 120 grams of Gentamicin per jar

Antibacterial

FOR ANIMAL USE ONLY - KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California)-Use Only as Directed

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-494

Net weight: 360 g (12.7 oz)

INDICATIONS: GentaMed™ Soluble Powder is recommended for the control and treatment of colibacillosis in weaning swine caused by strains of *E. coli* sensitive to gentamicin, and for the control and treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

DOSE:
Colibacillosis – 1 level scoop per 240 gallons of drinking water for 3 consecutive days, 0.5 mg/lb/day. For proportioner use, add 1 level scoop to 2 gallons stock solution and dispense at the rate of 1 oz/gal drinking water (25 mg/gal).
Swine Dysentery – 1 level scoop per 120 gallons of drinking water for 3 consecutive days, 1.0 mg/lb/day. For proportioner use, add 1 level scoop to 1 gallon stock solution and dispense at the rate of 1 oz/gal drinking water (50 mg/gal).

GentaMed™ is a Trademark of Bimeda, Inc.

Bimeda
GentaMed™
Soluble Powder
(Gentamicin Sulfate)
 Contains 120 grams of Gentamicin per jar
Antibacterial

FOR ANIMAL USE ONLY - KEEP OUT OF REACH OF CHILDREN Restricted Drug (California) - Use Only as Directed

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
 Approved by FDA under ANADA # 200-494

Net Weight:
 360 g (12.7 oz)

Manufactured by:
 Bimeda, Inc.
 Le Sueur, MN 56058
 www.bimeda.com

EACH g CONTAINS: Gentamicin sulfate equivalent to 333.33 mg gentamicin base.

RESIDUE WARNING: Swine treated with the recommended dose of GentaMed™ Soluble Powder must not be slaughtered for food for at least ten (10) days following the last treatment.

Do not store or offer medicated water in rusty metal containers.

Read accompanying directions carefully. Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).

For use in swine drinking water only.

1GEN009 8GEN001 Rev. 03/20

8 99855 30064 5

PULL HERE TO OPEN

TAKE TIME OBSERVE LABEL DIRECTIONS

GENTAMED

gentamicin sulfate powder

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:61133-5135
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
gentamicin sulfate (UNII: 8X7386QLV) (GENTAMICIN - UNII:T6Z9V48IKG)	GENTAMICIN	120 g in 360 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61133-5135-1	360 g in 1 JAR		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200494	12/14/2011	

Labeler - Bimeda, Inc. (060492923)

Registrant - Bimeda, Inc. (060492923)

Establishment

Name	Address	ID/FEI	Business Operations
Bimeda, Inc.		060492923	manufacture

Revised: 4/2022

Bimeda, Inc.