DOUBLE DINE- iodine liquid IBA

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Double Dine NPE Free

ACTIVE INGREDIENTS

ACTIVE INGREDIENT:

CHLORHEXIDINE GLUCONATE	0.50%
IODINE	0.50%
(PROVIDES A Full .5% AVAILABLE IODINE AS USED	

Inactive Ingredients

INERT INGREDIENTS:	
GLYCERIN(AS A HUMECTANT AND SKIN CONDITIONER)	2.00%
OTHER INERTS	97.00%
(INCLUDES BUFFERING AGENTS AND SURFACTANTS)	
TOTAL:	100.00%

THIS PRODUCT CONTAINS HIGHLY EFFECTIVE LEVELS OF FREE IODINE FOR FAST KILL OF MASTITIS CAUSING ORGANISMS.

INDICATIONS AND USAGE

GENERAL RECOMMENDATIONS

- 1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
- 2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
- 3. Do not use this product for cleaning or sanitizing equipment
- 4. Wash Teat Dip cups after each milking.

RECOMMENDED PROCEDURES FOR TEAT SANITATION A. PRE-MILKING (PRE-DIPPING)

- 1. REMOVE VISIBLE SOILS FROM THE TEATS.
- 2. OBSERVE FOREMILK BY STRIPPING TWO OR THREE STREAMS OF MILK INTO A STRIP CUP.
- 3. DIP OR SPRAY THE COW'S TEATS FULL LENGTH WITH DOUBLE DINE TEAT DIP.
- 4 WAIT APPROXIMATELY 30 SECONDS.
- 5. REMOVE ALL TEAT DIP BY THOROUGHLY DRYING THE TEATS AND TEAT ORIFICES WITH A CLEAN, SINGLE SERVICE PAPER TOWEL(S).
- ATTACH MILKING UNIT.

B. POST-MILKING (POST-DIPPING)

Immediately after milking dip all teats full length with DOUBLE DINE Teat Dip. Ensure

good coverage of all sides of the teats and teat orifices.

WARNINGS

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS. PROTECT THIS PRODUCT FROM FREEZING.

CAUTION: KEEP OUT OF REACH OF CHILDREN.
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY TO USE PRODUCTS, WASH ANY SENSITIVE AREAS. **INGESTION:**DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

Double Dine



DOUBLE DINE

iodine liquid

Product Information			
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1057
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	0.5 kg in 100 kg		
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	0.5 kg in 100 kg		

P	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1057-8	213 kg in 1 DRUM		

2 NDC:29983-1057-4	19.36 kg in 1 PAIL		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug		01/15/2015	

Labeler - IBA (019494160)

Registrant - IBA (019494160)

Establishment			
Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc. DBA, Custom Chemical Formulators Inc.		028311595	manufacture, api manufacture, pack

Revised: 2/2024 IBA