

PAIN RELIEVER PLUS- acetaminophen, caffeine tablet
Rapid Care, Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

RapidCare - Pain Reliever Plus

Acetaminophen 500mg

Caffeine 65mg

Pain reliever/fever reducer

Pain reliever aid

Uses

temporarily relieves minor aches and pains due to:

- headache
- muscular aches

Warnings

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you

take

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Caffeine warning: The recommended dose of this product contains about as much caffeine as

a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while

taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Quick medical attention is critical for adults as well

as for children, even if you do not notice any signs or symptoms.

Do not use

- if you are allergic to acetaminophen
- with any other drug containing
- acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- any new symptoms occur
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- do not use more than directed
- adults and children 12 years and over: take 2 tablets every 6 hours; not more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

store at 20-25°C (68-77° F)

corn starch, croscarmellose sodium, crospovidone, D&C red #27 aluminum lake, FD&C red #40 aluminum lake,

FD&C yellow #6 aluminum lake, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, silicon dioxide, stearic acid, titanium dioxide



100ct Pain Reliever Plus	Die 2.5 wide x 4.25 deep x 3.5 height
	UPC: 731946898031
	Reorder # 903-100
	NDC #

100 Tablets - 2 per Pouch
 • Temporary relief from pain, fever and inflammation
 • Soothes muscle aches and minor arthritis pain
 Acetaminophen 500 Mg - Pain Reliever / Fever Reducer
 Caffeine 65 Mg - Pain Reliever Aid



Pain Reliever Plus

Acetaminophen 500 Mg - Pain Reliever / Fever Reducer
 Caffeine 65 Mg - Pain Reliever Aid

- Temporary relief from pain, fever and inflammation
- Soothes muscle aches and minor arthritis pain

Reorder # 903-100



Distributed by:
 Rapid Care First Aid
 California, CA 92011
 rapidcarefirstaid.com



Pain Reliever Plus

Acetaminophen 500 Mg - Pain Reliever / Fever Reducer
 Caffeine 65 Mg - Pain Reliever Aid

- Temporary relief from pain, fever and inflammation
- Soothes muscle aches and minor arthritis pain

100 Tablets - 2 per Pouch
 DO NOT USE IF POUCH IS TORN OR DAMAGED

Drug Facts

Active ingredients (in each tablet)
 Acetaminophen 500mg
 Caffeine 65mg

Purposes
 Pain reliever/fever reducer
 Pain reliever aid

Uses temporarily relieves minor aches and pains due to: **headache** **muscular aches**

Warnings
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: **skin redness** **itching** **rash**
 If a skin reaction occurs, stop use and seek medical help right away.
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take **more than 8 tablets in 24 hours, which is the maximum daily amount** **with other drugs containing acetaminophen** **3 or more alcoholic drinks every day while using this product**
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use **if you are allergic to acetaminophen** **with any other drug containing acetaminophen (prescription or non-prescription).** If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
Stop use and ask a doctor if **any new symptoms occur** **painful area is red or swollen** **pain gets worse or lasts for more than 10 days** **fever gets worse or lasts for more than 3**

Drug Facts (continued)
 3 days. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Directions
Do not use more than directed
adults and children 12 years and over: take 2 tablets every 6 hours; not more than 8 tablets in 24 hours
children under 12 years: ask a doctor

Other information
Store at 20-25°C (68-77°F)

Inactive ingredients corn starch, croscarmellose sodium, crospovidone, D&C red #27 aluminum lake, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hydroxypropyl methylcellulose, polyethylene glycol, povidone, pregelatinized starch, silicon dioxide, stearic acid, titanium dioxide

DO NOT USE IF POUCH IS TORN OR DAMAGED



PAIN RELIEVER PLUS

acetaminophen, caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73659-014
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	

Product Characteristics

Color	brown	Score	score with uneven pieces
Shape	ROUND	Size	17mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73659-014-01	50 in 1 BOX	04/18/2022	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:73659-014-02	125 in 1 BOX	04/18/2022	
2		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/18/2022	

Labeler - Rapid Care, Inc (877432778)