BISMUTH SUBSALICYLATE- bismuth subsalicylate capsule, gelatin coated CVS PHARMACY, INC.

Stomach Relief

Drug Facts

Active ingredient (in each softgel)

Purpose

Bismuth subsalicylate 262 mgUpset stomach reliever and anti-diarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including: heartburn, indigestion, nausea, gas, belching
- fullness

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.

When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout

arthritis

When using this product

• a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- swallow with water, do not chew
- adults and children 12 years and over:
- 2 softgels every ½ hour or 4 softgels every hour as needed for diarrhea
- 2 softgels every ½ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 softgels) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each softgel contains: salicylate 102.33 mg
- very low sodium
- do not exceed 25°C

Inactive ingredients

aerosil, beeswax, FD&C Red#33, FD&C Yellow#6, gelatin, glycerin, light liquid paraffin, purified water, soya lecithin, sodium carboxy methyl cellulose, sorbitol solution, titanium dioxide

Questions or comments?

1-888-333-9792

PRINCIPAL DISPLAY PANEL

Stomach Relief

BISMUTH SUBSALICYLATE 262 mg Upset stomach reliever/Antidiarrheal

Soothing relief for 5 symptoms: • Heartburn

- Indigestion
- Upset stomach
- Nausea
- Diarrhea

12 SOFTGELS



Stomach Relief

BISMUTH SUBSALICYLATE 262 mg

5

CVSHealth.

Compare to the active ingredient in Pepto-Bismol® Liquicaps™

Stomach Relief

BISMUTH SUBSALICYLATE 262 mg Upset stomach reliever/Antidiarrheal

Soothing relief for 5 symptoms:

- Heartburn
- Indigestion
- Upset stomach
- Nausea
- Diarrhea



12 SOFTGELS

Actual Size Pill on Bottom Panel

Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 © 2020 CVS/pharmacy CVS.com® 1-800-SHOP CVS

Made in India V-35257



*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademarks Pepto-Bismol®

Questions or comments? 1-888-938-9792

cellulose, sorbito solution, titanium dioxide Bquid paraffin, purfiled water, soys lecifinin, sodium carboxy methyf serosa, beeswax, FD&C Red#35, FD&C Yeacuaf6, gelatin, glycerin, agint inactive ingredients

O"62 beeaxs fon ob ■

mulibos wol ynev

m each soffgel contains: sancytate 102.33 mg Other information

mqujuk bjeut) at cleat Ilnigs to jueto breveut detrydration caused by

- use untă diarrhea stops but not more than 2 days ■chādren under 12 years; ask a doctor
- do not exceed & doses (16 softgets) in 24 hours spowsty' peauprin' jugidespou' usness)
- Teagu) sonsglubninsvo tof bebsen as tuori if of tuori št vrava alagifica S 🖪
 - 2 softgels every 3£ hour or 4 softgels every hour as needed for medults and children 12 years and over:

Directions = swallow with water, do not chew

contact a Poison Control Center right away. Keep out of reach of children. In case of overdose, get medical help or asu encled licroisselong rites ask a health professional before use.

migratures lasts more than 2 days multipling in the ears or loss of hearing occurs

Dung Facts (continued)

 ■ symptoms get worse or last more than 2 days Stop use and ask a doctor if

stool and/or tongue may occur

THIS PRODUCT IS PACKAGED IN A CHILO-RESISTANT AND TAMPEN-EVIDENT PACKAGE, USE ONLY IF BLISTER UNIT IS INTACT.

when using this product a temporary, but harmers, darkening of the ■ supcosôn@spou (griuning the blood) = dispetes = good = srthribs Ask a doctor or pharmacist before use if you are taking any drug for

> pogs aug ui snamu 1846) aved voy li eau enoted notech a stat.

man ulcer ma bleeding problem m bloody or black shool Do not use if you have

ataluborq efial/ciles rento gnisist ■

early sign of Reys's syndrome, a rare but serious illness. Allergy alent: Contains salicytate. Do not take if you are an alergic to salicytance (including aspirin) voluigiudi occini, couzrig si quetor pecenze girese symbioms contiq pe au When using this product, if changes in behavior with nauses and from chicken pox or flu-flixe symptoms should not use this product. Reye's syndrome: Chadren and teenagers who have or are recovering

бицирая веб вазпеи иодзабрии шподначи upset stomach due to overhodugence in food and drink, including:

m pravance distribes maistribes

earlmeib-itns bns Upset stomach resever Purpose

Active ingredient (in each softgel)





bismuth subsalicylate capsule, gelatin coated

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69842-799

Route of Administration ORAL

Active Ingredient/Active Moiety Ingredient Name BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII: 0414PZ4LPZ) BISMUTH SUBSALICYLATE BISMUTH SUBSALICYLATE BISMUTH SUBSALICYLATE

Inactive Ingredients			
Ingredient Name	Strength		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)			
YELLOW WAX (UNII: 2ZA36H0S2V)			
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)			
LIGHT MINERAL OIL (UNII: N6K5787QVP)			
GELATIN (UNII: 2G86QN327L)			
GLYCERIN (UNII: PDC6A3C0OX)			
D&C RED NO. 33 (UNII: 9DBA0SBB0L)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
WATER (UNII: 059QF0KO0R)			

Product Characteristics							
Color	pink	Score	no score				
Shape	OVAL	Size	21mm				
Flavor		Imprint Code	262				
Contains							

Packaging						
# Item Code	Package Description	Marketing Start Date	Marketing End Date			
NDC:69842- 799-12	1 in 1 CARTON	07/09/2020				
1	12 in 1 BLISTER PACK; Type 0: Not a Combination Product					

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M008	07/09/2020			

Labeler - CVS PHARMACY, INC. (062312574)

Revised: 12/2023 CVS PHARMACY, INC.