

ANTACID CHERRY FLAVORED- aluminum hydroxide, magnesium hydroxide, simethicone liquid

Rij Pharmaceutical Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 5 mL) (One teaspoonful)

Aluminum Hydroxide (Equi. to Dried Gel, USP) 200 mg

Magnesium Hydroxide 200 mg

Simethicone 20 mg

Purpose

Antacid

Antigas

Uses

relieves

- heartburn
- acid indigestion
- sour stomach
- associated symptoms of gas and upset stomach
- overindulgence of food and drink

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

symptoms last for more than 2 weeks

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

Directions

- **shake well before each use**
- adults and children over 12 years, take 2 - 4 teaspoonsful between meals, at bedtime or as directed by a physician
- do not take more than 12 teaspoonfuls in a 24 hour period
- do not use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.
- Store at room temperature 15° - 30°C (59° - 86°F)
- keep tightly closed
- protect from freezing
- ** considered dietetically sodium free
- each teaspoon (5 mL) contains: magnesium 83 mg
- each teaspoon (5 mL) contains: sodium 1 mg
- does not meet USP requirements for preservative effectiveness

Inactive ingredients

flavors, hypromellose, parabens, saccharin, sodium hypochlorite, sorbitol, water and xanthan gum

Principal Display Panel

NDC 53807-405-12

Compare to the active ingredient in Mylanta®†



ANTACID

ANTACID • ANTI-GAS

Magnesia & Alumina
.....*Oral Suspension*
with Simethicone

- *Non-Constipating*
- *Pleasant Tasting*
- *Sodium Free***



12 FL. OZ. (355 mL)

ANTACID

Drug Facts

Active ingredients

(in each 5 mL) (One teaspoonful)

Purpose

Aluminum Hydroxide

(Equi. to Dried Gel, USP) 200 mgAntacid

Magnesium Hydroxide 200 mgAntacid

Simethicone 20 mgAntigas

Uses relieves ■ heartburn ■ acid indigestion ■ sour stomach
■ associated symptoms of gas and upset stomach ■ overindulgence
of food and drink

Warnings

Ask a doctor before use if you have ■ kidney disease
■ a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are
taking a prescription drug. Antacids may interact with certain
prescription drugs.

Stop use and ask a doctor if ■ symptoms last more than 2
weeks

Keep out of reach of children. In case of accidental overdose,
get medical help or contact a Poison Control Center immediately.

Directions ■ shake well before each use

- adults and children over 12 years: take 2-4 teaspoonsful between
meals, at bedtime or as directed by a physician
- do not take more than 12 teaspoonfuls in a 24 hour period
- do not use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON CAP
IS BROKEN OR MISSING.** ■ store at room temperature 15°-30°C
(59°-86°F) ■ keep tightly closed ■ protect from freezing

** considered dietetically sodium free

- each teaspoon (5 mL) contains: magnesium 83 mg
- each teaspoon (5 mL) contains: sodium 1 mg
- does not meet USP requirements for preservative effectiveness.

Inactive ingredients: flavors, hypromellose, parabens
saccharin, sodium hypochlorite, sorbitol, water and xanthan gum.

†This product is not manufactured or distributed by Johnson & Johnson • Merck, owner of
the registered trademark Mylanta®.

▲ Preferred Plus Pharmacy products are satisfaction guaranteed or your money back from
retailer where purchased.

Manufactured By:
RLJ PHARMACEUTICAL CORP.
40 Commercial Avenue,
Middletown, NY 10941

Distributed By:
Kinray Inc.,
Whitestone NY 11357



aluminum hydroxide, magnesium hydroxide, simethicone liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53807-405
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDRO XIDE	200 mg in 5 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDRO XIDE ION - UNII:9159UV381P)	MAGNESIUM HYDRO XIDE	200 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	20 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
BUTYL PARABEN (UNII: 3QP1IU3FV8)	
SACCHARIN (UNII: FST467XS7D)	
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)	
SORBITOL (UNII: 506T60A25R)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	WHITE (opaque)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53807-405-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part332	03/16/1999	

Labeler - Rij Pharmaceutical Corporation (144679156)

Establishment

Name	Address	ID/FEI	Business Operations
Rij Pharmaceutical Corporation		144679156	manufacture(53807-405)

Revised: 4/2018

Rij Pharmaceutical Corporation