

**NON-ASPIRIN - acetaminophen tablet**  
**Etex Pharm Co., Ltd**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active Ingredients (in each tablet)**

Acetaminophen 500mg

**Purpose**

Pain Reliever/fever reducer

**Warnings**

**Liver Warning:**

this product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days

These could be sign of a serious condition

**If pregnant or breast-feeding**

ask a health professional before use.

**Keep out of reach of children.**

Enter section text here

### **Overdose warning:**

Taking more than the recommended dose(overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- do not take more than directed (see overdose warning)

---

adults and children 12 years and over	<ul style="list-style-type: none"><li>• take 2 tablets every 4 to 6 hours as needed</li><li>• do not take more than 8 tablets in 24 hours</li><li>• do not use for more than 10 days unless directed by a doctor</li></ul>
---	--

children under 12 years	do not use this adult Extra Strength product in children under 6 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage
----------------------------	---

---

### **Other Information**

- **do not use if imprinted safety seal under cap is broken or missing.**
- store between 20-25°C (68-77°F)
- see end panel for lot number and expiration date

### **Inactive Ingredients**

benzalkonium chloride, boric acid, edetate disodium, purified water, sodium borate, and sodium chloride

non-aspirin 50 tablet

**Active ingredient in each tablet** Acetaminophen 500mg. **Purpose** Pain reliever/fever reducer  
**Uses** temporarily relieves minor aches and pains due to:  
 ■ headache ■ muscular aches ■ backache ■ arthritis ■ the common cold ■ toothache ■ menstrual cramps ■ temporarily reduces fever  
**Warnings Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 8 tablets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product  
**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription) If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product  
**Ask a doctor before use if you have liver disease** Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin  
**Stop use and ask a doctor if** ■ new symptoms occur ■ redness or swelling is present ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days  
 These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.**

Pure-Aid™ NDC No. 43459-0152-6  
 \*Compare to the active ingredient of Extra Strength Tylenol®

Non-Aspirin  
**Pain Relief**  
 Contains Acetaminophen **EXTRA STRENGTH**  
 Pain Reliever / Fever Reducer

50 TABLETS  
 500mg each

Tablets

← READ THE LABEL →

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  
**Directions** ■ do not take more than directed (see overdose warning) **Adults and children 12 years and over** ■ take 2 tablets every 4 to 6 hours as needed ■ do not take more than 8 tablets in 24 hours ■ do not use for more than 10 days unless directed by a doctor **Children under 12 years** ■ do not use this adult Extra Strength product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage  
**Other information** ■ do not use if imprinted safety seal under cap is broken or missing ■ store between 20-25°C (68-77°F) ■ see end panel for lot number and expiration date  
**Inactive ingredients** Dextrin, Pregelatinized starch, Lauryl Sodium Sulfate, Hydroxypropyl Cellulose, Dry starch  
 \*This product is not manufactured or distributed by McNeil Consumer & Specialty Pharmaceuticals, distributor of Tylenol® Extra Strength.  
**Exclusively distributed by:**  
 Kareway Product Inc., Compton, CA 90220  
 Made in Korea

LOT 9001  
 EXP. 06/2013

LOT 9001  
 EXP. 06/2013

8 56023 00152 0

Pure-Aid™ NDC No. 43459-0152-6  
 \*Compare to the active ingredient of Extra Strength Tylenol®

Non-Aspirin  
**Pain Relief**  
 Contains Acetaminophen **EXTRA STRENGTH**  
 Pain Reliever / Fever Reducer

50 TABLETS  
 500mg each

Tablets

**Drug Facts**  
**Active ingredient (in each tablet)** Acetaminophen 500mg. **Purpose** Pain reliever/fever reducer  
**Uses** temporarily relieves minor aches and pains due to:  
 ■ headache ■ muscular aches ■ backache ■ arthritis ■ the common cold ■ toothache ■ menstrual cramps ■ temporarily reduces fever  
**Warnings Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 8 tablets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product  
**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription) If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product  
**Ask a doctor before use if you have liver disease** Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin  
**Stop use and ask a doctor if** ■ new symptoms occur ■ redness or swelling is present ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days  
 These could be signs of a serious condition.

Pure-Aid™ NDC No. 43459-0152-6  
 \*Compare to the active ingredient of Extra Strength Tylenol®

Non-Aspirin  
**Pain Relief**  
 Contains Acetaminophen **EXTRA STRENGTH**  
 Pain Reliever / Fever Reducer

50 TABLETS  
 500mg each

Tablets

**Drug Facts (continued)**  
**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.**  
**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  
**Directions** ■ do not take more than directed (see overdose warning) **Adults and children 12 years and over** ■ take 2 tablets every 4 to 6 hours as needed ■ do not take more than 8 tablets in 24 hours ■ do not use for more than 10 days unless directed by a doctor **Children under 12 years** ■ do not use this adult Extra Strength product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage  
**Other information** ■ do not use if imprinted safety seal under cap is missing ■ store between 20-25°C (68-77°F) ■ see end panel for lot number and expiration date  
**Inactive ingredients** Dextrin, Pregelatinized starch, Lauryl Sodium Sulfate, Hydroxypropyl Cellulose, Dry starch

Pure-Aid™  
**Pain Relief**  
**EXTRA STRENGTH**

\*This product is not manufactured or distributed by McNeil Consumer & Specialty Pharmaceuticals, distributor of Tylenol® Extra Strength.  
**Exclusively distributed by:**  
 Kareway Product Inc.,  
 Compton, CA 90220  
**Non-Aspirin**  
 Made in Korea  
 NDC No. 43459-0152-6

**Uses**

temporarily relieves minor aches and pains due to:

- headache
- muscular aches

- backache
- arthritis
- the common cold
- toothache
- menstrual cramps
- temporarily reduces fever

## NON-ASPIRIN

acetaminophen tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43459-0153
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	500 mg

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	13mm
<b>Flavor</b>		<b>Imprint Code</b>	A500
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43459-0153-6	1 in 1 BOX		
1		50 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	08/01/2010	

## NON-ASPIRIN

acetaminophen tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43459-0152
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	500 mg

**Product Characteristics**

Color	white	Score	no score
Shape	CAPSULE	Size	17mm
Flavor		Imprint Code	A500
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43459-0152-6	1 in 1 BOX		
1		40 in 1 BOTTLE		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/15/2010	

**Labeler** - Etex Pharm Co., Ltd (631034449)**Registrant** - Etex Pharm Co., Ltd (631034449)**Establishment**

Name	Address	ID/FEI	Business Operations
Etex Pharm Co., Ltd		631034449	manufacture

Revised: 6/2010

Etex Pharm Co., Ltd