

**ORIGINAL NIGHT-TIME MULTI-SYMPTOM COLD/FLU RELIEF- acetaminophen,  
dextromethorphan hydrobromide, doxylamine succinate liquid  
Discount Drug Mart**

-----

**Drug Facts**

**Active ingredients (in each 30 mL dose cup)**

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine Succinate 12.5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

**Keep out of reach of children**

**Keep out of reach of children.**

**Uses**

temporarily relieves cold/flu symptoms

- sore throat
- headache
- minor aches and pain
- fever
- runny nose and sneezing
- cough due to minor throat and bronchial irritation

**DO NOT USE IF IMPRINTED SHRINK BRAND IS MISSING OR BROKEN**

Failure to follow these warnings could result in serious consequences.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if adult/child takes: ● more than 4 doses in 24 hours, which is the maximum daily amount for this product ● with other drugs containing acetaminophen ● 3 or more alcoholic drinks everyday while using this product.

**Allergy Alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: ● skin reddening ● blisters ● rash

If skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash,

Nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription) If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are now taking a prescription monamine oxidase inhibitor (MAOI) (certain drugs for depression psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

### **Ask a doctor before use if you have**

- a sodium restricted diet
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland.

### **Ask a doctor or pharmacist before use**

- if you are taking sedatives or tranquilizers
- if you are taking the blood thinning drug warfarin

### **When using this product**

- **do not use more than directed**
- avoid alcoholic drinks
- excitability may occur, especially in children
- marked drowsiness may occur
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- redness or swelling is present
- symptoms do not get better within 7 days or are accompanied by a fever
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Overdose warning:**

Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help, or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms

### **Directions**

- take only as recommended-see Overdose warning
- Use dose cup or tablespoon (TBSP)
- do not exceed 4 doses per 24 hours
- if taking Night Time at night and Day Time during the day, limit total to 4 doses per 24 hours.

---

adults and children 12 years and over	30 mL (2 TBSP) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

---

### **Other information**

- **each 30 mL dose cup contains:** sodium 45 mg
- store at room temperature

### **Inactive ingredients**

citric acid, D and C yellow # 10, FD and C Green No. 3, FD and C Yellow No. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sucrose

### **Questions?**

Call weekdays **1-877-798-5944**

**DISCOUNT drug mart FOOD FAIR Original Night-Time Multi-Symptom Cold/Flu Relief product label**

**Drug Facts (continued)**

**Stop use and ask a doctor if:**

- redness or swelling is present
- symptoms do not get better within 7 days or are accompanied by a fever
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause serious health problems. In case of overdose get medical help or contact a Poison Control Center rightaway. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as recommended—see Overdose warning
- Use dose cup or tablespoon (TBSPI) • do not exceed 4 doses per 24 hours
- If taking Night Time at night and Day Time during the day, limit total to 4 doses per 24 hours.

adults & children 12 years and over	30 mL (2 TBSPI) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

**Other information** • each 30 mL dose cup contains: sodium 45 mg  
• store at room temperature

**Inactive ingredients** citric acid, D&C yellow #10, FD&C Green No. 3, FD&C Yellow No. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sucrose

**Questions?** Call weekdays 1-877-796-5944  
\*This product is not manufactured or distributed by Procter & Gamble owner of the registered trademark Vicks® Nyquil®

**SATISFACTION GUARANTEED**  
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

**DISCOUNT**  
**drug mart**  
FOOD FAIR

\*COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® NYQUIL®

**Original**  
**Night-Time**  
**MULTI-SYMP TOM**  
**COLD/FLU RELIEF**

Acetaminophen..... Pain Reliever/Fever Reducer  
Dextromethorphan HBr..... Cough Suppressant  
Doxylamine Succinate..... Antihistamine

**12 FL OZ (354 mL)**

**DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN**  
Failure to follow these warnings could result in serious consequences

**Drug Facts**

**Active ingredients (in each 30 mL dose cup)**

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine Succinate 12.5 mg	Antihistamine

**Uses** temporarily relieve cold/flu symptoms: • sore throat • headache • minor aches and pain • fever • runny nose and sneezing • cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount; • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash

If skin reaction occurs, stop use and seek medical help right away.

**Some throat warnings:** If more throat is severe, persists for more than two days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product • to make a child sleepy

**Ask a doctor before use if you have:** • a certain restricted diet • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or its occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use** • if you are taking sedatives or tranquilizers • if you are taking the blood thinning drug warfarin

**When using this product:** • do not use more than directed • avoid alcoholic drinks • excitability may occur, especially in children • masked drowsiness may occur • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness

\*COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® NYQUIL®

**DISCOUNT**  
**drug mart**  
**FOOD FAIR**

**Original**  
**Night-Time Multi-Symptom Cold/Flu Relief**

Acetaminophen ..... Pain Reliever/Fever Reducer  
Dextromethorphan HBR ..... Cough Suppressant  
Doxylamine Succinate ..... Antihistamine  
12 FL OZ (354 mL)

\* This product is not manufactured or distributed by Procter and Gamble owner of the registered trademark Vicks® Nyquil®

LR-065 REV 01

**SATISFACTION GUARANTEED**

IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO THE STORE WHERE PURCHASED. IF UNABLE TO RETURN TO THE STORE, SEND REASON FOR DISSATISFACTION , NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE MEDINNA, OHIO 44256

<b>ORIGINAL NIGHT-TIME MULTI-SYMP TOM COLD/FLU RELIEF</b>			
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53943-501
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 355W5USQ3G)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SUCROSE</b> (UNII: C151H8M554)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-501-28	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/21/2014	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/21/2014	

**Labeler** - Discount Drug Mart (047741335)