

**ADULT LOW DOSE ENTERIC COATED ASPIRIN- aspirin tablet, coated**  
**Geri-Care Pharmaceutical Corp**

-----  
**gc 981**

**Active ingredient (in each tablet)**

Aspirin 81 mg(NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- for the temporary relief of minor aches and pains
- ask your doctor about other uses for aspirin

**Warnings**

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy Alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer.
- if you ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have: -high blood pressure -heart disease -liver cirrhosis -kidney disease -asthma

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout or arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding: -feel faint -vomit blood -have bloody or black stools -have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- any new symptoms occur
- ringing in the ears or loss of hearing occurs

These can be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not exceed recommended dose
- drink a full glass of water with each dose
- swallow whole, do not chew or crush
- **adults and children 12 years and older:** take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor
- **children under 12:** ask a doctor

**Other information**

- store at 20-25°C (68-77°F); excursions permitted between 15°C - 30°C (59°F - 86°F)

**Inactive ingredients**

cellulose, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, PEG, polydextrose, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, starch, talc, titanium dioxide, triacetin, triethyl citrate, wax.

Questions or comments?

1-800-540-3765

package label

NDC 57896-981-01



**ADULT LOW DOSE ENTERIC COATED ASPIRIN 81 mg PAIN RELIEVER (NSAID)**

Compare to active ingredient in BAYER® Low Dose Aspirin™

**100 Tablets**

**Tamper Evident:** Do not use if imprinted seal under cap is missing or broken.

**Drug Facts**

**Active ingredient (in each tablet)** Aspirin 81 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ for the temporary relief of minor aches and pains ■ ask your doctor about other uses for aspirin

**Warnings**  
**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy Alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if** ■ the stomach bleeding warning applies to you ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have: □ high blood pressure □ heart disease □ liver cirrhosis □ kidney disease □ asthma

**\*This product is not manufactured or distributed by the owner of the registered trademark Bayer®.**  
 Dist. by: Gerl-Care Pharmaceuticals Corp.  
 1295 Towbin Avenue, Lakewood, NJ 08701

REV 1023L



3 57896 98101 3

**PEEL HERE FOR MORE DRUG FACTS**

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout or arthritis.

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away.  
 ■ you experience any of the following signs of stomach bleeding: □ feel faint □ vomit blood □ have bloody or black stools  
 ■ pain gets worse or lasts more than 10 days  
 ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur ■ ringing in the ears or loss of hearing occurs  
 These can be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose get medical help or contact a Poison Control Center right away.

**Directions** ■ do not exceed recommended dose ■ drink a full glass of water with each dose ■ swallow whole, do not chew or crush

**Adults and children 12 years and older:** take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor

**Children under 12:** ask a doctor

**Other information** ■ store at 20-25°C (68-77°F); excursions permitted between 15°C-30° (59°F - 86°F)

**Inactive ingredients:** cellulose, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, PEG, polydextrose, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, starch, talc, titanium dioxide, triacetin, triethyl citrate, wax

**Questions or comments?** 1-800-540-3765

**STOP PEELING**

## ADULT LOW DOSE ENTERIC COATED ASPIRIN

aspirin tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:57896-981
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

### Inactive Ingredients

Ingredient Name	Strength

<b>SHELLAC</b> (UNII: 46N107B71O)
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>TRIACETIN</b> (UNII: XHX3C3X673)
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)
<b>METHACRYLIC ACID</b> (UNII: 1CS02G8656)
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	L
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-981-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2012	
2	NDC:57896-981-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2013	
3	NDC:57896-981-30	300 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2013	
4	NDC:57896-981-12	120 in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2012	
5	NDC:57896-981-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2000	

**Labeler** - Geri-Care Pharmaceutical Corp (611196254)

**Registrant** - Geri-Care Pharmaceutical Corp (611196254)

Revised: 11/2023

Geri-Care Pharmaceutical Corp