HEAD CONGESTION MUCUS PE- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated GOODSENSE

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1173-GDS-2022-0803

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever
J	reducer
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
 - sinus congestion and pressure
 - headache
 - minor aches and pains
 - nasal congestion
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	 take 2 caplets every 4 hours do not take more than 10 caplets in 24 hours
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

GOODSENSE®

NDC 50804-173-02

Non-Drowsy

Head Congestion + Mucus PE

Acetaminophen, Guaifenesin, Phenylephrine HCl

Pain Reliever / Fever Reducer, Expectorant, Nasal Decongestant

For Relief of

- Sinus Pressure
- Headache
- Chest Congestion

Actual Size

24 CAPLETS

Compare to active ingredients of Sudafed PE® Head Congestion + Mucus†

corn starch, croscarmellose sodium, crospovidone, colloidal silicon dioxide, Inactive ingredients

■ retain carton for complete product information and warnings store between 20-25°C (68-77°F) in a dry place Other information

Drug Facts (continued)

stearic acid, talc, titanium dioxide polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, magnesium stearate, maltodextrin, microcrystalline cellulose,

Dung Facts (continued)

A	9	12 years
	■ ask a doctor	children under
	24 hours	
	 do not take more than 10 caplets in 	12 years and over

■ do not take more than directed (see overdose warning) Directions

notice any signs or symptoms. attention is critical for adults as well as for children even if you do not Poison Control Center right away (1-800-222-1222). Quick medical Overdose warning: In case of overdose, get medical help or contact a Keep out of reach of children. If pregnant or breast-feeding, ask a health professional before use.

L nese could be signs of a senous condition.

- cough comes back or occurs with rash or headache that lasts
 - uew symptoms occur
 - redness or swelling is present
- fever gets worse or lasts more than 3 days ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
 - Dervousness, dizziness, or sleeplessness occur Stop use and ask a doctor if

When using this product do not exceed recommended dosage Trinning arug warrarin

Ask a doctor or pharmacist before use if you are taking the blood

Drug Facts (continued)

 condu tust occurs with too much phiedm (mucus) quonic pronchitis, or emphysema

 bersistent or chronic cough such as occurs with smoking, asthma, ■ trouble unnating due to an enlarged prostate gland

■ thyroid disease ■ diabetes uiðu pjoog blessnie Ask a doctor before use if you have I her disease I heart disease

 if you have ever had an allergic reaction to this product or any of its ingredients an MAOI, ask a doctor or pharmacist before taking this product. the MAOI drug. If you do not know if your prescription drug contains conditions, or Parkinson's disease), or for 2 weeks after stopping (MAOI) (certain drugs for depression, psychiatric or emotional it you are now taking a prescription monoamine oxidase inhibitor

acetaminophen, ask a doctor or pharmacist. nonprescription). If you are not sure whether a drug contains

■ with any other drug containing acetaminophen (prescription or

If a skin reaction occurs, stop use and seek medical help right away. Symptoms may include:

skin reddening

blisters

rash Allergy alert. Acetaminophen may cause severe skin reactions.

3 or more alcoholic drinks every day while using this product ■ with other drugs containing acetaminophen

■ more than 4,000 mg of acetaminophen in 24 hours Severe liver damage may occur if you take dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours.

Liver warning: This product contains acetaminophen. The maximum daily

Drug Facts (continued)



NSGS = temporarily relieves these symptoms associated with

Nasal decongestant Роспутербите нСт 5 mg. Guaifenesin 200 mg Expectorant

Purpose Active ingredients (in each caplet)

Drug Facts

 remporanty reduces tever bassageways of bothersome mucus and make coughs more productive helps loosen pitlegm (mucus) and thin bronchial secretions to rid the bronchial minor aches and pains

■ usasi congeston siuns coudestion and bressure hay fever or other upper respiratory allergies, and the common cold:

Drug Facts (continued)

GOODSENSE.

NDC 50804-173-02

Non-Drowsy

Head Congestion + Mucus PE

Pain Reliever / Fever Reducer, Expectorant, Nasal Decongestant

cetaminophen, Guaifenesin, Phenylephrine HCl

For Relief of

- Sinus Pressure

Headache

Chest Congestion

24 CAPLETS

Compare to active ingredients of Sudafed PE® Head Congestion + Mucus†

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Sudafed PE Head Congestion + Mucus.

PE®

Distributed by:
Perrigo Direct, Inc.
Perrigo Direct, Inc.
Peachtree City, GA 30269
www.PerrigoDirect.com
1-844-705-4384
GoodSense® is a registered
trademark of L Perrigo Company.

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

HEAD CONGESTION MUCUS PE

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50804-173
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE	5 mg

Inactive Ingredients	
Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
CROSPOVIDONE (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: 08232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	white	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	AAA;1173
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:50804- 173-02	2 in 1 CARTON	08/03/2022		
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			

rmation		
pplication Number or Monograph Citation	Marketing Start Date	Marketing End Date
341	08/03/2022	
•	Citation	Citation Date

Labeler - GOODSENSE (076059836)

Revised: 8/2022 GOODSENSE