**MOUNTAIN SERIES WEEKENDER MEDICAL - benzalkonium chloride, povidone-iodine, acetaminophen, aspirin, diphenhydramine hydrochloride, ibuprofen, bacitracin zinc, neomycin sulfate, polymyxin b sulfate**
**Tender Corporation dba Adventure Medical Kits**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

---

**Mountain Series Weekender Medical Kit**

## Active Ingredient

Active Ingredient:

Benzalkonium Chloride 0.40%

## Purpose

Antiseptic

## Use

For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

## Warnings

Warning: For external use only.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.

Do not use in the eyes or over large areas of the body.

## Directions

Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

## Inactive Ingredients

Inactive Ingredient: Purified water

LOT/EXP: Made in CHINA

20130301

Antiseptic Towelette

Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID

**Active Ingredients**

Active Ingredient: .....Bacitracin Zinc 400 units

Neomycin Sulfate 5mg ( equivalent to 3.5 mg Neomycin base)

Polymyxin B Sulfate 5000 units

**Purpose**

Triple Antibiotic

Uses: To help prevent infection in:
minor cuts; scrapes; burns

**Warnings**

For external use only.

Do not use: in eyes; over large areas of the body;

If allergic to any of the ingredients; for more than one week unless directed by a physician.

Stop use and consult a doctor:

if the condition persists or gets worse; a rash or other allergic reaction develops

Keep out of reach of children.

If ingested, contact a Poison

Control Center right away.

**Directions**

Directions: clean affected area; apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily; may be covered with a sterile bandage

Other information:

Store at room temperature.

**Inactive Ingredient**
water

Genuine Triple Antibiotic

First Aid Ointment

To Help Prevent Infection

Each Gram Contains:

Bacitracin Zinc 400 units

Neomycin Sulfate 5 mg

(equivalent to 3.5 mg

Neomycin base)

Polymyxin B Sulfate 5000 units

Net Wt. 0.5g ; (1/64 oz)

Manufactured in CHINA for

GENUINE FIRST AID.

Triple Antibiotic Ointment 10pcs

Net wt. 0.9g (1/32oz)

100
Triple Antibiotic

**Active ingredient (in each tablet)**

Aspirin (NSAID*) 325 mg

*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

**Temporarily relieves minor aches and pains associated with**

- headache
- minor arthritis
- common cold
- menstrual cramps
- muscular aches
- backache
- toothache

**Temporarily reduces fever.**

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

**Do not use**
- if you have ever had an allergic reaction to any other pain reliever/ fever reducer
- right before or after heart surgery
- if you are taking prescription drugs for gout, diabetes or arthritis

**Ask a doctor before use if**
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**
- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- if ringing in the ears or loss of hearing occurs redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**
- **do not take more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- drink a full glass of water with each dose

**Adults and children: (12 years and older)**

Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

**Children under 12 years:**

Do not give to children under 12 years of age.

**Other information**

- read all product information before using
- store at room temperature 59º-86ºF (15º-30ºC)
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**

croscarmellose sodium*, hypromellose, microcrystalline cellulose*, mineral oil*, polyethylene glycol*, propylene glycol*, starch, titanium dioxide*

**may contain*

**Questions or comments?**

1-800-634-7680

**116R Medique APAP 325 mg Label**

Medique®

Aspirin

5 Grain (325 mg)

Pain Reliever/Fever Reducer (NSAID)

See new warnings information

East To Swallow

Film Coated Tablets

24 Tablets

(12 x 2)

Compare active ingredients to Bayer®

Registered Trademark Bayer Consumer

Tamper Evident Unit Dose Packets

**Active ingredients**

(in each capsule)

Diphenhydramine Hydrochloride 25 mg

**Purpose**

Antihistamine

## Uses

Temporarily relieves runny nose and decreases sneezing, itching of the nose and throat, itchy-watery eyes due to hay fever or other respiratory allergies.

## Warnings

**Do not use**

- with any other product containing diphenhydramine, even one that is used on skin.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- glaucoma

**Ask a doctor or pharmacist before use if you are**

- taking any drugs for asthma
- taking sedatives or tranquilizers

**When Using This Product**

- marked drowsiness may occur
- avoid alcohol beverages
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of the reach of children**. In case of overdose, contact a physician or poison control center immediately.

## Directions

- **do not use more than directed**

**Adults and children: (12 years and older):** Take 1 to 2 capsules every 4 to 6 hours as needed. Do not take more than 12 capsules in 24 hours.

**Children under 12 years:** Do not give to children under 12 years of age.

## Other information

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

## Inactive ingredients

D&C Red #28, FD&C Blue #1, FD&C Red #40, gelatin, starch

**Questions or comments? 1-800-634-7680**

**182R Medique Diphen Label**

Collect Medi-Bucks

See inside flap for more details

Medique®

Diphen

Diphenhydramine HCl 25 mg

Hay Fever/Allergies

Fiebre del Heno/Alergias

Pull to Open

TiraParaAbrir

Easy To Swallow Capsules

Capsulas Faciles de Tragar

200 Capsules

(200 x 1)

Tamper Evident Unit Dose Packets

Empaquetado con Sellado Evidente en Dosis Untarias

***Active ingredient***
***(in each tablet)***

Ibuprofen (NSAID*) 200 mg

*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

**Temporarily relieves minor aches and pains associated with**

- headache
- backache
- common cold
- minor arthritis pain
- toothache
- menstrual cramps
- muscular aches

**Temporarily reduces fever.**

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include:

- hives
- asthma (wheezing)
- rash
- blisters
- skin reddening

- facial swelling
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**
- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts for more than 10 days
  - fever gets worse or lasts for more than 3 days
  - redness or swelling is present in the painful area
- any new or unexpected symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***
- **do not use more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children: (12 years and older)**

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:**

Do not give to children under 12 years of age.

***Other information***

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 40°C (above 104°F)
- tamper evident sealed packets
- do not use any opened or torn packets

***Inactive ingredients***

carnauba wax*, cellulose*, colloidal silicon dioxide, corn starch*, hypromellose, iron oxide red*, lactose, magnesium stearate, microcrystalline cellulose*, polydextrose, polyethylene glycol, povidone, silica*, sodium lauryl sulfate*, sodium starch glycolate, stearic acid*, titanium dioxide, triacetin*

*may contain

***Questions or comments?*** 1-800-634-7680

**Principal Display Panel**

Medi-First®
Ibuprofen 200 mg
Pain Reliever/Fever Reducer (NSAID)
Easy To Swallow Film Coated Tablets
See new warnings information
100 tablets (50 x 2)
Compare active ingredient to: Genuine Advil®
Registered trademark Wyeth Consumer
Tamper Evident Unit Dose Packets

**Active ingredient (in each tablet)**

Acetaminophen 325 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

**For the temporary relief of minor aches and pains associated with**

- headache
- common cold
- muscular aches
- toothache
- minor arthritis pain
- menstrual cramps

**For the reduction of fever.**

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

**Ask a doctor before use if you have**

- liver disease

**Ask a doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin

**Stop using and ask a doctor if**

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- redness or swelling is present

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**If pregnant or breast-feeding,** ask a health professional before use.

## Directions

- **do not use more than directed**

**Adults and children: (12 years and older)**

Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.

**Children under 12 years:**

Do not give to children under 12 years of age.

## Other information

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

## Inactive ingredients

corn starch, hypromellose, maltodextrin*, microcrystalline cellulose*, polyethylene glycol, povidone, pregelatinized starch*, sodium starch glycolate, stearic acid

** may contain*

**Questions or comments? 1-800-634-7680**

**Principal Display Panel**

**145R Medique APAP 325 mg Label**

Collect MediBucks

See inside flap for more details

Medique®

APAP

Acetaminophen 325 mg

Pain Reliever/Fever Reducer

Alivia el Dolor/Reduce la Fiebre

Easy To Swallow

Film Coated Tablets

Facil de Tragar Tabletas con Cubierta Pelicular

Pull to Open

Tire Para Abrir

See new warnings information

500 Tablets

(250 x 2)

Tamper Evident Unit Dose Packets

Empaquetado con Sellado Evidente en Dosis Unitarias

**Weekender Kit Label**

Mountain Series Medical Kit Weekender

Adventure Medical Kits BE SAFE

1-6 People up to 7 days

Povidone-iodine 10%

Antiseptic

Warnings

Do not use

- if allergic to iodine
- in the eyes

For external use only

Ask a doctor before use if injuries are

- deep or puncture wounds
- serious burns

Stop use and ask a doctor if

- redness, irritation, swelling or pain persists or increases
- infection occurs

Avoid pooling beneath patient

Keep out of reach of children. In case of accidental ingestion, seek professionalassistance or consult a poison control center immediately.



Medique®

2 Tablets

Made in USA

# APAP

| Active ingredient (in each tablet) | Purpose |
|---|---|
| Acetaminophen 325 mg. | Pain reliever/ fever reducer |

## Uses

**For the temporary relief of minor aches and pains associated with**

- headache
- muscular aches
- minor arthritis pain
- toothache
- common cold
- menstrual cramps

**For the reduction of fever.**

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Store at 59°-86°F (15-30°C) • Tamper-evident sealed packets • Do not use any opened or torn packets

*Mfd. for MEDIQUE PRODUCTS • Fort Myers, FL 33967*

Lot 1831 Exp. 01-2013

## Warnings (continued)

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

**Ask a doctor before use if you have**

- liver disease

**Ask a doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin

**Stop using and ask a doctor if**

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- redness or swelling is present

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**If pregnant or breast-feeding,** ask a health professional before use.

## Directions

- do not use more than directed

**Adults and children: (12 years and older)**
Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.

**Children under 12 years:**
Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

Made in USA

2 Tablets

# IBUPROFEN

**Active ingredient (in each tablet)** **Purpose**

Ibuprofen (NSAID*) 200mg ..... Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

**Uses** **Temporarily relieves minor aches and pains due to**

■ headache ■ toothache ■ backache ■ menstrual cramps ■ common cold ■ muscular aches ■ minor arthritis pain

**Temporarily reduces fever.**

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash ■ asthma (wheezing) ■ blisters ■ shock If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Store at 68-77°F (20-25°C) • Avoid excessive heat, above 104°F (40°C) • Tamper-evident sealed packets • Do not use any opened or torn packets

*Mfd. for* MEDIQUE PRODUCTS • Fort Myers, FL 33967

Lot 1922 Exp. 02-2012

**Warnings** (continued)

**Do not use**

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if**

■ stomach bleeding warning applies to you
■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product**

■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area
■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ **do not use more than directed**
■ **the smallest effective dose should be used**

**Adults and Children (12 years and older):** Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INGREDIENTS

2 Tablets

# Medique®

Made in USA

# Aspirin

| Active ingredient (in each tablet) | Purpose |
|---|---|
| Aspirin (NSAID*) 325 mg..... | Pain reliever/fever reducer |

*nonsteroidal anti-inflammatory drug

**Uses** Temporarily relieves minor aches and pains due to
■ headache ■ muscular aches ■ minor arthritis pain ■ backache
■ common cold ■ toothache ■ menstrual cramps
Temporarily reduces fever.

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash ■ asthma (wheezing) ■ blisters ■ shock
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Store at 59°-86°F (15°-30°C) • Tamper-Evident Sealed Packets • Do Not Use any Opened or Torn Packets

*Mfd. for MEDIQUE PRODUCTS • Fort Myers, FL 33967*

Lot 1856 Exp. 11-2012

## Warnings (continued)

**Do not use**
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery ■ if you are taking a prescription drug for gout, diabetes or arthritis

**Ask a doctor before use if**
■ stomach bleeding warning applies to you
■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**
■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product**
■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing ■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions: Do not use more than directed

**Adults and children: (12 years and older)** Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

**Children: (under 12 years)** Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

# Medique®

NDC# 47682-182-46

Made in USA

# Diphen

1 Capsule

| Active Ingredient (In each capsule) | Purpose |
|---|---|
| Diphenhydramine Hydrochloride 25mg | Antihistamine |

**Uses:**

Temporarily relieves the following symptoms associated with hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

Store at room temperature • Tamper-Evident Sealed Packets • Do Not Use any Opened or Torn Packets

**Mfd. for MEDIQUE PRODUCTS**
**Fort Myers, FL 33967**

**Warnings**

**Ask a Doctor Before Use**

**If you have:**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

**If you are:**

- taking any drugs for asthma
- taking sedatives or tranquilizers

**When Using this product:**

- drowsiness may occur
- avoid alcohol beverages

alcohol, sedatives and tranquilizers may increase the drowsiness effect

use caution when driving a motor vehicle or operating machinery

excitability may occur, especially in children.

**o not exceed recommended dosage. Keep this and all drugs ut of the reach of children.** In case of accidental overdose, ntact a physician or poison control center immediately. As with y drug, if you are pregnant or nursing a baby, seek the advice a health professional before using this product.

**rections:** Do not use more than directed

ults and children: (12 years and older) Take 1 capsule ry 4 to 6 hours as needed. Do not take more than 12 capsules 4 hours, or as directed by a doctor.

dren under 12 years: Do not give to children under 12 years ge unless directed by a doctor.

**REORDER** AST-001

# Antiseptic Towelette

# Toallitas Antisepticas

Genuine First Aid LLC, Clearwater FL 33755
www.GenuineFirstAid.com

ISO 9001:2010

1 /pouch

GENUINE FIRST AID®

2

DRUG FACTS - Antiseptic Towelette

Active Ingredient: Purpose:
Benzalkonium Chloride 0.40%..First Aid Antiseptic
Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.
Warnings: For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
If unusual redness, swelling or other symptoms occur, consult a physician immediately.
Do not use: In the eyes, or over large areas of the body.
Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.
Inactive ingredient: Purified water

Made in CHINA

- - - - TEAR HERE - - - -

LOT/EXP:
20130301

Reorder TAO-001

# Genuine Triple Antibiotic

**First Aid Ointment**
**To Help Prevent Infection**

**Each Gram Contains:**
**Bacitracin Zinc 400 units**
**Neomycin Sulfate 5 mg**
**(equivalent to 3.5 mg Neomycin base)**
**Polymyxin B Sulfate 5000 units**
**Net Wt 0.5g · (1/64 oz)**

Manufactured in CHINA for
GENUINE FIRST AID®

Drug Facts
Uses: To help prevent infection in • minor cuts • scrapes • burns
Warnings:
For external use only
Do not use • in eyes • over large areas of the body • if allergic to any of the ingredients • for more than one week unless directed by a physician
Stop use and consult a doctor • if the condition persists or gets worse • a rash or other allergic reaction develops
Keep out of reach of children.
If ingested, contact a Poison Control Center right away.
Directions: • clean affected area • apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily • may be covered with a sterile bandage
Other information:
Store at room temperature
LOT/EXP:
20130118

ACTIVITY
1-6 PEOPLE
UP TO 7 DAYS
Adventure Medical Kits
BE SAFE
MOUNTAIN SERIES
MEDICAL KIT
WEEKENDER




# MOUNTAIN SERIES WEEKENDER MEDICAL

benzalkonium chloride, povidone-iodine, acetaminophen, aspirin, diphenhydramine hydrochloride, ibuprofen, bacitracin zinc, neomycin sulfate, polymyxin b sulfate kit

## Product Information

| | | | |
|---|---|---|---|
| **Product Type** | HUMAN OTC DRUG | **Item Code (Source)** | NDC:44224-0118 |

## Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:44224-0118-1 | 1 in 1 KIT | | |

## Quantity of Parts

| Part # | Package Quantity | Total Product Quantity |
|---|---|---|
| **Part 1** | 6 PACKAGE | 4.8 mL |
| **Part 2** | 3 TUBE | 1.5 g |
| **Part 3** | 2 PACKET | 4 |

| | | |
|---|---|---|
| **Part 4** | 2 PACKET | 2 |
| **Part 5** | 1 PACKET | 2 |
| **Part 6** | 2 PACKET | 4 |
| **Part 7** | 1 PACKET | 22 g |

## Part 1 of 7

## ANTISEPTIC TOWELETTE

benzalkonium chloride swab

### Product Information

| | | | |
|---|---|---|---|
| **Item Code (Source)** | NDC:52124-0001 | | |
| **Route of Administration** | TOPICAL | **DEA Schedule** | |

### Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|---|---|
| **BENZALKONIUM CHLORIDE** (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y) | BENZALKONIUM CHLORIDE | 0.40 mL in 100 mL |

### Inactive Ingredients

| Ingredient Name | Strength |
|---|---|
| **WATER** (UNII: 059QF0KO0R) | |

### Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:52124-0001-1 | 0.8 mL in 1 PACKAGE | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC monograph not final | part333A | 04/23/2010 | |

## Part 2 of 7

## GENUINE TRIPLE ANTIBIOTIC

bacitracin zinc,neomycin sulfate,polymyxin b sulfate ointment

### Product Information

| | | |
|---|---|---|
| **Item Code (Source)** | NDC:52124-0003 | |

| Route of Administration | TOPICAL | DEA Schedule | |
|---|---|---|---|

### Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|---|---|
| **BACITRACIN ZINC** (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I) | BACITRACIN ZINC | 400 [iU] in 1 g |
| **NEOMYCIN SULFATE** (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297) | NEOMYCIN SULFATE | 5 mg in 1 g |
| **POLYMYXIN B SULFATE** (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K) | POLYMYXIN B SULFATE | 5000 [iU] in 1 g |

### Inactive Ingredients

| Ingredient Name | Strength |
|---|---|
| **WATER** (UNII: 059QF0KO0R) | |

### Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:52124-0003-1 | 0.5 g in 1 TUBE | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC monograph final | part333B | 02/16/2010 | |

## Part 3 of 7

## MEDIQUE ASPIRIN

aspirin tablet, film coated

### Product Information

| Item Code (Source) | NDC:47682-116 | | |
|---|---|---|---|
| Route of Administration | ORAL | DEA Schedule | |

### Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|---|---|
| **ASPIRIN** (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN | 325 mg |

### Inactive Ingredients

| Ingredient Name | Strength |
|---|---|
| **HYPROMELLOSES** (UNII: 3NXW29V3WO) | |
| **POLYETHYLENE GLYCOL** (UNII: 3WJQ0SDW1A) | |
| **PROPYLENE GLYCOL** (UNII: 6DC9Q167V3) | |

| | |
|---|---|
| **STARCH, CORN** (UNII: O8232NY3SJ) | |

### Product Characteristics

| | | | |
|---|---|---|---|
| **Color** | white (white) | **Score** | no score |
| **Shape** | ROUND (ROUND) | **Size** | 10mm |
| **Flavor** | | **Imprint Code** | 44;157;aspirin |
| **Contains** | | | |

### Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:47682-116-99 | 2 in 1 PACKET | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC monograph final | part343 | 12/30/2008 | |

## Part 4 of 7

## MEDIQUE DIPHEN

diphenhydramine hydrochloride capsule

### Product Information

| | | | |
|---|---|---|---|
| **Item Code (Source)** | NDC:47682-182 | | |
| **Route of Administration** | ORAL | **DEA Schedule** | |

### Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|---|---|
| **DIPHENHYDRAMINE HYDROCHLORIDE** (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M) | DIPHENHYDRAMINE HYDROCHLORIDE | 25 mg |

### Inactive Ingredients

| Ingredient Name | Strength |
|---|---|
| **D&C RED NO. 28** (UNII: 767IP0Y5NH) | |
| **FD&C BLUE NO. 1** (UNII: H3R47K3TBD) | |
| **FD&C RED NO. 40** (UNII: WZB9127XOA) | |
| **GELATIN** (UNII: 2G86QN327L) | |
| **STARCH, CORN** (UNII: O8232NY3SJ) | |

### Product Characteristics

| | | | |
|---|---|---|---|
| **Color** | pink (pink) , white (white) | **Score** | no score |
| **Shape** | CAPSULE (CAPSULE) | **Size** | 14mm |
| **Flavor** | | **Imprint Code** | CPC;835 |
| **Contains** | | | |

## Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:47682-182-46 | 1 in 1 PACKET | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC monograph final | part341 | 12/30/2008 | |

## Part 5 of 7

# MEDI-FIRST IBUPROFEN

ibuprofen tablet, film coated

## Product Information

| | | | |
|---|---|---|---|
| **Item Code (Source)** | NDC:47682-808 | | |
| **Route of Administration** | ORAL | **DEA Schedule** | |

## Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|---|---|
| **IBUPROFEN** (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN | 200 mg |

## Inactive Ingredients

| Ingredient Name | Strength |
|---|---|
| **CELLULOSE, MICROCRYSTALLINE** (UNII: OP1R32D61U) | |
| **HYPROMELLOSES** (UNII: 3NXW29V3WO) | |
| **LACTOSE** (UNII: J2B2A4N98G) | |
| **MAGNESIUM STEARATE** (UNII: 70097M6I30) | |
| **POLYDEXTROSE** (UNII: VH2XOU12IE) | |
| **POLYETHYLENE GLYCOL** (UNII: 3WJQ0SDW1A) | |
| **POVIDONE** (UNII: FZ989GH94E) | |
| **SILICON DIOXIDE** (UNII: ETJ7Z6XBU4) | |
| **STARCH, CORN** (UNII: O8232NY3SJ) | |
| **STEARIC ACID** (UNII: 4ELV7Z65AP) | |
| **TITANIUM DIOXIDE** (UNII: 15FIX9V2JP) | |
| **TRIACETIN** (UNII: XHX3C3X673) | |

| Product Characteristics | | | |
|---|---|---|---|
| **Color** | white (white) | **Score** | no score |
| **Shape** | ROUND (ROUND) | **Size** | 10mm |
| **Flavor** | | **Imprint Code** | 44;352 |
| **Contains** | | | |

| Packaging | | | | |
|---|---|---|---|---|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:47682-808-99 | 2 in 1 PACKET | | |

| Marketing Information | | | |
|---|---|---|---|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| ANDA | ANDA075139 | 12/30/2008 | |

## Part 6 of 7

## MEDIQUE APAP

acetaminophen tablet, film coated

| Product Information | | | |
|---|---|---|---|
| **Item Code (Source)** | NDC:47682-145 | | |
| **Route of Administration** | ORAL | **DEA Schedule** | |

| Active Ingredient/Active Moiety | | |
|---|---|---|
| Ingredient Name | Basis of Strength | Strength |
| **ACETAMINOPHEN** (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |

| Inactive Ingredients | |
|---|---|
| Ingredient Name | Strength |
| **HYPROMELLOSES** (UNII: 3NXW29V3WO) | |
| **POLYETHYLENE GLYCOL** (UNII: 3WJQ0SDW1A) | |
| **POVIDONE** (UNII: FZ989GH94E) | |
| **SODIUM STARCH GLYCOLATE TYPE A POTATO** (UNII: 5856J3G2A2) | |
| **STARCH, CORN** (UNII: O8232NY3SJ) | |
| **STEARIC ACID** (UNII: 4ELV7Z65AP) | |

| Product Characteristics | | | |
|---|---|---|---|
| **Color** | white (white) | **Score** | no score |

| | | | |
|---|---|---|---|
| **Shape** | ROUND (ROUND) | **Size** | 10mm |
| **Flavor** | | **Imprint Code** | AZ;234 |
| **Contains** | | | |

### Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:47682-145-99 | 2 in 1 PACKET | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC monograph not final | part343 | 12/30/2008 | |

## Part 7 of 7

## APLICARE POVIDONE-IODINE SOLUTION

povidone-iodine solution solution

### Product Information

| | | | |
|---|---|---|---|
| **Item Code (Source)** | NDC:52380-0001 | | |
| **Route of Administration** | TOPICAL | **DEA Schedule** | |

### Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|---|---|
| **POVIDONE-IODINE** (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4) | POVIDONE-IODINE | 9.8 g in 100 g |

### Inactive Ingredients

| Ingredient Name | Strength |
|---|---|
| **CITRIC ACID MONOHYDRATE** (UNII: 2968PHW8QP) | |
| **SODIUM PHOSPHATE, DIBASIC** (UNII: GR686LBA74) | |
| **SODIUM HYDROXIDE** (UNII: 55X04QC32I) | |
| **NONOXYNOL-9** (UNII: 48Q180SH9T) | |
| **WATER** (UNII: 059QF0KO0R) | |

### Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:52380-0001-3 | 22 g in 1 PACKET | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC monograph not final | part333A | 03/01/1984 | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC monograph final | part333A | 08/01/2011 | |

## Labeler - Tender Corporation dba Adventure Medical Kits (064437304)

Revised: 8/2011 Tender Corporation dba Adventure Medical Kits