

**IBUPROFEN- ibuprofen tablet, film coated**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Ibuprofen Tablets, 200 mg Drug Facts**

**Active ingredient (in each tablet)**

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- **Adults and children 12 years and older:**
  - take 1 tablet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 tablet, 2 tablets may be used
  - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- **Children under 12 years:** ask a doctor

### **Other information**

- read all warnings and directions before use
- store between 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- see end panel for lot number and expiration date

### **Inactive ingredients**

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

### **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

meijer®

Compare to Advil® active ingredient

ibuprofen Tablets, 200 mg

SEE REVISED WARNING

Pain Reliever | Fever Reducer (NSAID)

50 Tablets | actual size

60471 6E C15



PID 467466

**DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.**  
Important: Read all product information before using.  
Keep this package for important information.

**Drug Facts**

**Active ingredient (in each tablet)**  
Ibuprofen 200 mg (NSAID).....Pain reliever/fever reducer  
nonsteroidal anti-inflammatory drug

**Purposes**  
Ibuprofen temporarily releases minor aches and pains due to:  
■ muscle aches ■ minor pain of arthritis ■ headache  
■ the common cold ■ menstrual cramps ■ temporarily reduces fever

**Warnings**

Always alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters ■ If an allergic reaction occurs, stop use and seek medical help right away.  
Some stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding, the chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for longer time than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers/fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as

**Drug Facts (cont. from front)**

asthma ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug  
When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the genital area ■ any new symptoms appear  
If pregnant or breast-feeding, ask a health professional before use. It is especially important to use ibuprofen at 20 weeks or later in pregnancy unless explicitly directed to do so by a doctor because it may cause problems for the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away (1-800-222-1222).

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used ■ Adults and children 12 years and older: ■ take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor Children under 12 years: ask a doctor

**Drug Facts (continued)**

Other list of contents ■ read all warnings and directions before use ■ store between 20° to 25°C (68° to 77°F) ■ avoid high humidity ■ use and expiration date ■ inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, hydroxypropylcellulose, polyethylene glycol, polybutene-2, croscarmellose sodium, titanium dioxide, polyethylene glycol, polybutene-2, croscarmellose sodium, hydroxypropylcellulose, polyethylene glycol, polybutene-2, croscarmellose sodium, titanium dioxide  
**Questions or comments?** 1-800-779-9260

\*\*This product is not manufactured or distributed by PF Consumer HealthCare, LLC, owner of the registered trademark Advil®.  
DISTR BY MEIJER DISTRIBUTION, N.C.  
MI 49544  
www.meijer.com



actual size

meijer

NDC 41250-604-71

Compare to Advil® active ingredient\*\*

**ibuprofen** Tablets, 200 mg

Pain Reliever | Fever Reducer (NSAID)

**50 Tablets**

actual size

SEE REVISED WARNING

**IBUPROFEN**  
ibuprofen tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-604
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	BROWN	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	I2
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:41250-604-82	2 in 1 CARTON	05/13/2011	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-604-71	1 in 1 CARTON	02/15/1988	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41250-604-78	1 in 1 CARTON	12/15/1990	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:41250-604-90	500 in 1 BOTTLE; Type 0: Not a Combination Product	05/22/1992	

5	NDC:41250-604-93	1000 in 1 BOTTLE; Type 0: Not a Combination Product	01/17/2007	12/26/2016
6	NDC:41250-604-62	1 in 1 CARTON	09/07/1994	
6		24 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:41250-604-76	1 in 1 CARTON	03/20/1992	03/31/2021
7		120 in 1 BOTTLE; Type 0: Not a Combination Product		
8	NDC:41250-604-85	250 in 1 BOTTLE; Type 0: Not a Combination Product	04/30/2018	04/30/2018

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA072096	02/15/1988	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 10/2023

Meijer Distribution Inc