

# FESOTERODINE FUMARATE- fesoterodine fumarate tablet, extended release Golden State Medical Supply, Inc.

-----

## HIGHLIGHTS OF PRESCRIBING INFORMATION FESOTERODINE FUMARATE

**These highlights do not include all the information needed to use FESOTERODINE FUMARATE EXTENDED-RELEASE TABLETS safely and effectively. See full prescribing information for FESOTERODINE FUMARATE EXTENDED-RELEASE TABLETS.**

### FESOTERODINE FUMARATE extended-release tablets, for oral use

**Initial U.S. Approval: 2008**

#### ----- RECENT MAJOR CHANGES -----

Indications and Usage ( 1.1) 06/2021  
Dosage and Administration ( 2.1, 2.3, 2.5, 2.6) 06/2021  
Contraindications ( 4) 06/2021  
Warnings and Precautions ( 5.1, 5.2, 5.3, 5.4, 5.6) 06/2021

#### ----- INDICATIONS AND USAGE -----

Fesoterodine fumarate is indicated for the treatment of:

- Overactive bladder(OAB) in adults with symptoms of urge urinary incontinence, urgency, and frequency. ( 1.1)

#### ----- DOSAGE AND ADMINISTRATION -----

- OAB in Adults: The recommended starting dosage is 4 mg orally once daily. Based upon individual response and tolerability, increase the maximum dosage of 8 mg once daily. ( 2.1)
- Adult Patients with Renal or Hepatic Impairment: Refer to the full prescribing information for recommended dosage. ( 2.3)
- Dosage Modifications Due to Strong CYP3A4 Inhibitors: Refer to the full prescribing information for recommended dosage. ( 2.5)
- Administration: Swallow whole with liquid. Do not chew, divide, or crush. Take with or without food. ( 2.6)

#### ----- CONTRAINDICATIONS -----

- Known or suspected hypersensitivity to fesoterodine fumarate extended-release tablets or any of its ingredients or to tolterodine tartrate tablets or tolterodine tartrate extended-release capsules. ( 4)
- Urinary retention ( 4)
- Gastric retention ( 4)
- Uncontrolled narrow-angle glaucoma. ( 4)

#### ----- WARNINGS AND PRECAUTIONS -----

- Angioedema: Promptly discontinue fesoterodine fumarate extended-release tablets and provide appropriate therapy. ( 5.1).
- Urinary Retention: Fesoterodine fumarate extended-release tablets are not recommended in patients with clinically significant bladder outlet obstruction because of the risk of urinary retention. ( 5.2)
- Decreased Gastrointestinal Motility: Fesoterodine fumarate extended-release tablets are not recommended for use in patients with decreased gastrointestinal motility, such as those with severe constipation. ( 5.3)
- Decreased Gastrointestinal Motility: Fesoterodine fumarate extended-release tablets are not recommended for use in patients with decreased gastrointestinal motility, such as those with severe constipation. ( 5.3)
- Worsening of Narrow Angle Glaucoma: Use fesoterodine fumarate extended-release tablets with caution in patients being treated for narrow-angle glaucoma. ( 5.4)
- Central Nervous System Effects: Somnolence has been reported with fesoterodine fumarate extended-release tablets. Advise patients not to drive or operate heavy machinery until they know how fesoterodine fumarate extended-release tablets affects them ( 5.5)
- Worsening of Myasthenia Gravis Symptoms: Use fesoterodine fumarate extended-release tablets with caution in patients with myasthenia gravis. ( 5.6)

#### ----- ADVERSE REACTIONS -----

Most frequently reported adverse events with fesoterodine fumarate extended-release tablets in adult patients with OAB ( $\geq 4\%$ ) were: dry mouth (placebo, 7%; fesoterodine fumarate extended-release tablets 4 mg, 19%; fesoterodine fumarate extended-release tablets 8 mg, 35%) and constipation (placebo, 2%; fesoterodine fumarate extended-release tablets 4 mg, 4%; fesoterodine fumarate extended-release

tablets 8 mg, 6%). ( 6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Dr. Reddy's Laboratories, Inc. at 1-888-375-3784 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

**Revised: 3/2022**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **RECENT MAJOR CHANGES**

#### **1 INDICATIONS AND USAGE**

##### **1.1 Adult Overactive Bladder**

#### **2 DOSAGE AND ADMINISTRATION**

##### **2.1 Recommended Dosage for Adult Patients with OAB**

##### **2.3 Recommended Dosage in Adult Patients with Renal Impairment**

##### **2.5 Fesoterodine fumarate Dosage Modifications Due to Strong CYP3A4 Inhibitors**

##### **2.6 Administration Instructions**

#### **3 DOSAGE FORMS AND STRENGTHS**

#### **4 CONTRAINDICATIONS**

#### **5 WARNINGS AND PRECAUTIONS**

##### **5.1 Angioedema**

##### **5.2 Urinary Retention in Adult Patients with Bladder Outlet Obstruction**

##### **5.3 Decreased Gastrointestinal Motility**

##### **5.4 Worsening of Narrow-Angle Glaucoma**

##### **5.5 Central Nervous Systems Effects**

##### **5.6 Worsening of Myasthenia Gravis Symptoms**

#### **6 ADVERSE REACTIONS**

##### **6.1 Clinical Trials Experience**

##### **6.2 Post-marketing Experience**

#### **7 DRUG INTERACTIONS**

##### **7.1 Antimuscarinic Drugs**

##### **7.2 CYP3A4 Inhibitors**

##### **7.3 CYP3A4 Inducers**

##### **7.4 CYP2D6 Inhibitors**

##### **7.5 Drugs Metabolized by Cytochrome P450**

##### **7.6 Oral Contraceptives**

##### **7.7 Warfarin**

##### **7.8 Drug-Laboratory Test Interactions**

#### **8 USE IN SPECIFIC POPULATIONS**

##### **8.1 Pregnancy**

##### **8.2 Lactation**

##### **8.4 Pediatric Use**

##### **8.5 Geriatric Use**

##### **8.6 Renal Impairment**

##### **8.7 Hepatic Impairment**

#### **10 OVERDOSAGE**

#### **11 DESCRIPTION**

#### **12 CLINICAL PHARMACOLOGY**

##### **12.1 Mechanism of Action**

12.2 Pharmacodynamics

12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

**13.1 Carcinogenesis, Mutagenesis, Impairment Of Fertility**

### **14 CLINICAL STUDIES**

**14.1 Adult Overactive Bladder**

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

#### **1.1 Adult Overactive Bladder**

Fesoterodine fumarate extended-release tablets are indicated for the treatment of overactive bladder (OAB) in adults with symptoms of urge urinary incontinence, urgency, and frequency.

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Recommended Dosage for Adult Patients with OAB**

The recommended starting dosage of fesoterodine fumarate extended-release tablets in adults is 4 mg orally once daily. Based upon individual response and tolerability, increase to the maximum dosage of fesoterodine fumarate extended-release tablets 8 mg once daily. For administration instructions, see Dosage and Administration ( 2.6).

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

#### **2.3 Recommended Dosage in Adult Patients with Renal Impairment**

The recommended dosage of fesoterodine fumarate extended-release tablets in adult patients with renal impairment is described in Table 1 [see Use in Specific Populations ( 8.6)]. For administration instructions, see Dosage and Administration ( 2.6).

**Table 1: Fesoterodine Fumarate Extended-Release Tablets Recommended Dose in Adult Patients with Renal Impairment (Administered Orally Once Daily)**

<b>Estimated Creatinine Clearance <sup>1</sup></b>	<b>Recommended Dose</b>
CLcr 30 to 89 mL/min	8 mg

CLcr 15 to 29 mL/min	4 mg
CLcr <sup>c</sup> 15 mL/min	4 mg

<sup>1</sup> Calculate CLcr using the Cockcroft-Gault formula

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

## **2.5 Fesoterodine fumarate Dosage Modifications Due to Strong CYP3A4 Inhibitors**

### Adult Patients with OAB

The maximum recommended dosage is fesoterodine fumarate extended-release tablets 4 mg orally once daily in adult patients taking strong CYP3A4 inhibitors [see Drug Interactions ( 7.2) and Clinical Pharmacology ( 12.3)]. For administration instructions, see Dosage and Administration ( 2.6).

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

## **2.6 Administration Instructions**

Swallow fesoterodine fumarate extended-release tablets whole with liquid. Do not chew, divide, or crush. Take with or without food [see *Clinical Pharmacology* ( 12.3) ].

## **3 DOSAGE FORMS AND STRENGTHS**

Fesoterodine fumarate extended-release tablets, 4 mg are light blue colored, oval shaped, biconvex, film coated, debossed with ' ˆY ' on one side and 'FS' on the other side.

Fesoterodine fumarate extended-release tablets, 8 mg are blue colored, oval shaped, biconvex, film coated, debossed with ' ˆY ' on one side and 'FT' on the other side.

## **4 CONTRAINDICATIONS**

Fesoterodine fumarate extended-release tablets are contraindicated in patients with any of the following,

- known or suspected hypersensitivity to fesoterodine fumarate extended-release tablets or any of its ingredients, or to tolterodine tartrate tablets or tolterodine tartrate extended-release capsules [ see *Clinical Pharmacology* ( 12.1) ].
- Reactions have included angioedema [ see *Warnings and Precautions* ( 5.1) ].
- urinary retention [ see *Warnings and Precautions* ( 5.2) ]
- gastric retention [ see *Warnings and Precautions* ( 5.3) ]
- uncontrolled narrow-angle glaucoma [ see *Warnings and Precautions* ( 5.4) ]

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Angioedema**

Angioedema of the face, lips, tongue, and/or larynx has been reported with fesoterodine fumarate extended-release tablets . In some cases, angioedema occurred after the first dose; however, cases have been reported to occur hours after the first dose or after multiple doses. Angioedema associated with upper airway swelling may be life-threatening.

Fesoterodine fumarate extended-release tablets contraindicated in patients with a known or suspected hypersensitivity to fesoterodine fumarate extended-release tablets or any of its ingredients [ *see Contraindications ( 4)*  ]. If involvement of the tongue, hypopharynx, or larynx occurs, fesoterodine fumarate should be promptly discontinued and appropriate therapy and/or measures to ensure a patent airway should be promptly provided.

## **5.2 Urinary Retention in Adult Patients with Bladder Outlet Obstruction**

The use of fesoterodine fumarate extended-release tablets, like other antimuscarinic drugs, in patients with clinically significant bladder outlet obstruction, including patients with urinary retention, may result in further urinary retention and kidney injury. The use of fesoterodine fumarate is not recommended in patients with clinically significant bladder outlet obstruction, and is contraindicated in patients with urinary retention [ *see Contraindications ( 4) and Adverse Reactions ( 6.1)* ].

## **5.3 Decreased Gastrointestinal Motility**

Fesoterodine fumarate extended-release tablets are associated with decreased gastric motility. Fesoterodine fumarate extended-release tablets are contraindicated in patients with gastric retention [ *see Contraindications ( 4)*  ]. The use of Fesoterodine fumarate extended-release tablets are not recommended in patients with decreased gastrointestinal motility, such as those with severe constipation.

## **5.4 Worsening of Narrow-Angle Glaucoma**

Fesoterodine fumarate extended-release tablets can worsen controlled narrow-angle glaucoma. Fesoterodine fumarate extended-release tablets are contraindicated in patients with uncontrolled narrow-angle glaucoma [ *see Contraindications (4)* ]. Fesoterodine fumarate should be used with caution in patients being treated for narrow-angle glaucoma..

## **5.5 Central Nervous Systems Effects**

Fesoterodine fumarate is associated with anticholinergic central nervous system (CNS) adverse reactions [ *see Adverse Reactions (6.1)* ]. A variety of CNS anticholinergic effects have been reported, including headache, dizziness, and somnolence. Patients should be monitored for signs of anticholinergic CNS effects, particularly after beginning treatment or increasing the dose. Advise patients not to drive or operate heavy machinery until they know how fesoterodine fumarate affects them. If a patient experiences anticholinergic CNS effects, fesoterodine fumarate extended-release tablets dose reduction or discontinuation should be considered.

## **5.6 Worsening of Myasthenia Gravis Symptoms**

Fesoterodine fumarate extended-release tablets should be used with caution in patients with myasthenia gravis due to the risk of worsening of symptoms of the disease.

# **6 ADVERSE REACTIONS**

The following clinically significant adverse reactions are described elsewhere in labeling:

- Angioedema [ see *Warnings and Precautions ( 5.1)*  ]
- Urinary Retention [ see *Warnings and Precautions ( 5.2)*  ]
- Decreased Gastrointestinal Motility [ see *Warnings and Precautions ( 5.3)*  ]

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

### Adult Overactive Bladder (OAB)

The safety of fesoterodine fumarate extended-release tablets was evaluated in Phase 2 and 3 controlled trials in a total of 2,859 patients with overactive bladder, of which 2,288 were treated with fesoterodine fumarate extended-release tablets. Of this total, 782 received fesoterodine fumarate extended-release tablets 4 mg/day, and 785 received fesoterodine fumarate extended-release tablets 8 mg/day with treatment periods of 8- or 12- weeks. Approximately 80% of these patients had greater than 10-weeks of exposure to fesoterodine fumarate extended-release tablets in these trials.

A total of 1,964 patients participated in two 12-week, Phase 3 efficacy and safety studies and subsequent open-label extension studies. In these two studies combined, 554 patients received fesoterodine fumarate extended-release tablets 4 mg/day and 566 patients received fesoterodine fumarate extended-release tablets 8 mg/day.

In Phase 2 and 3 placebo-controlled trials combined, the incidences of serious adverse events in patients receiving placebo, fesoterodine fumarate extended-release tablets 4 mg, and fesoterodine fumarate extended-release tablets 8 mg were 1.9%, 3.5%, and 2.9%, respectively. All serious adverse events were judged to be not related or unlikely to be related to study medication by the investigator, except for four patients receiving fesoterodine fumarate who reported one serious adverse reaction each: angina, chest pain, gastroenteritis, and QT prolongation on ECG.

The most commonly reported adverse event in patients treated with fesoterodine fumarate was dry mouth. The incidence of dry mouth was higher in those taking 8 mg/day (35%) and in those taking 4 mg/day (19%), as compared to placebo (7%). Dry mouth led to discontinuation in 0.4%, 0.4%, and 0.8% of patients receiving placebo, fesoterodine fumarate extended-release tablets 4 mg, and fesoterodine fumarate extended-release tablets 8 mg, respectively. For those patients who reported dry mouth, most had their first occurrence of the event within the first month of treatment.

The second most commonly reported adverse event was constipation. The incidence of constipation was 2% in those taking placebo, 4% in those taking 4 mg/day, and 6% in those taking 8 mg/day.

Table 4 lists adverse events, regardless of causality, that were reported in the combined Phase 3, randomized, placebo-controlled trials at an incidence greater than placebo and in 1% or more of patients treated with fesoterodine fumarate extended-release tablets 4 or 8 mg once daily for up to 12 weeks.

**Table 4: Adverse Events with an Incidence Exceeding the Placebo Rate and Reported by  $\geq$ 1% of Patients from Double-Blind, Placebo-Controlled Phase 3 Trials of 12-weeks Treatment Duration**

	Placebo	Fesoterodine Fumarate Extended-Release	Fesoterodine Fumarate Extended-Release
System organ			

<b>System organ class/Preferred term</b>	<b>Placebo N=554 %</b>	<b>Tablets 4 mg/day N=554 %</b>	<b>Tablets 8 mg/day N=566 %</b>
Gastrointestinal disorders			
Dry mouth	7.0	18.8	34.6
Constipation	2.0	4.2	6.0
Dyspepsia	0.5	1.6	2.3
Nausea	1.3	0.7	1.9
Abdominal pain upper	0.5	1.1	0.5
Infections			
Urinary tract infection	3.1	3.2	4.2
Upper respiratory tract infection	2.2	2.5	1.8
Eye disorders			
Dry eyes	0	1.4	3.7
Renal and urinary disorders			
Dysuria	0.7	1.3	1.6
Urinary retention	0.2	1.1	1.4
Respiratory disorders			
Cough	0.5	1.6	0.9
Dry throat	0.4	0.9	2.3
General disorders			
Edema peripheral	0.7	0.7	1.2
Musculoskeletal disorders			
Back pain	0.4	2.0	0.9
Psychiatric disorders			
Insomnia	0.5	1.3	0.4
Investigations			
ALT increased	0.9	0.5	1.2
GGT increased	0.4	0.4	1.2
Skin disorders			
Rash	0.5	0.7	1.1

ALT = alanine aminotransferase; GGT = gamma glutamyltransferase

Patients also received fesoterodine fumarate extended-release tablets for up to three years in open-label extension phases of one Phase 2 and two Phase 3 controlled trials. In all open-label trials combined, 857, 701, 529, and 105 patients received fesoterodine fumarate extended-release tablets for at least 6 months, 1 year, 2 years, and 3 years, respectively. The adverse events observed during long-term, open-label studies were similar to those observed in the 12-week, placebo-controlled studies, and included dry mouth, constipation, dry eyes, dyspepsia, and abdominal pain. Similar to the controlled studies, most adverse events of dry mouth and constipation were mild to moderate in intensity. Serious adverse events, judged to be at least possibly related to study medication by the investigator and reported more than once during the open-label treatment period of up to 3 years, included urinary retention (3 cases), diverticulitis (3

cases), constipation (2 cases), irritable bowel syndrome (2 cases), and electrocardiogram QT corrected interval prolongation (2 cases).

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

## **6.2 Post-marketing Experience**

The following adverse reactions have been identified during post-approval use of fesoterodine extended-release tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Cardiac disorders:* Palpitations

*Central nervous system disorders:* Dizziness, headache, somnolence

*Eye disorders:* Blurred vision

*General disorders and administrative site conditions:* Hypersensitivity reactions, including angioedema with airway obstruction, face edema

*Skin and subcutaneous tissue disorders:* Urticaria, pruritus

## **7 DRUG INTERACTIONS**

### **7.1 Antimuscarinic Drugs**

Coadministration of fesoterodine fumarate with other antimuscarinic agents that produce dry mouth, constipation, urinary retention, and other anticholinergic pharmacological effects may increase the frequency and/or severity of such effects. Anticholinergic agents may potentially alter the absorption of some concomitantly administered drugs due to anticholinergic effects on gastrointestinal motility.

### **7.2 CYP3A4 Inhibitors**

Doses of fesoterodine fumarate extended-release tablets greater than 4 mg are not recommended in adult patients taking strong CYP3A4 inhibitors, such as ketoconazole, itraconazole, and clarithromycin [ *see Dosage and Administration ( 2.5)*  ].

In a study in adults, coadministration of the strong CYP3A4 inhibitor ketoconazole with fesoterodine led to approximately a doubling of the maximum concentration ( $C_{max}$ ) and area under the concentration versus time curve (AUC) of 5-hydroxymethyl tolterodine (5-HMT), the active metabolite of fesoterodine. Compared with CYP2D6 extensive metabolizers not taking ketoconazole, further increases in the exposure to 5-HMT were observed in subjects who were CYP2D6 poor metabolizers taking ketoconazole [ *see Clinical Pharmacology ( 12.3)*  ].

There is no clinically relevant effect of moderate CYP3A4 inhibitors on the pharmacokinetics of fesoterodine. Following blockade of CYP3A4 by coadministration of the moderate CYP3A4 inhibitor fluconazole 200 mg twice a day for 2 days, the average (90% confidence interval) increase in  $C_{max}$  and AUC of the active metabolite of fesoterodine was approximately 19% (11% to 28%) and 27% (18% to 36%) respectively. No dosing adjustments are recommended in the presence of moderate CYP3A4 inhibitors (e.g., erythromycin, fluconazole, diltiazem, verapamil and grapefruit juice).

The effect of weak CYP3A4 inhibitors (e.g. cimetidine) was not examined; it is not

expected to be in excess of the effect of moderate inhibitors [ see *Clinical Pharmacology ( 12.3)*  ].

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

### **7.3 CYP3A4 Inducers**

No dosing adjustments are recommended in the presence of CYP3A4 inducers, such as rifampin and carbamazepine. Following induction of CYP3A4 by coadministration of rifampin 600 mg once a day,  $C_{max}$  and AUC of the active metabolite of fesoterodine decreased by approximately 70% and 75%, respectively, after oral administration of fesoterodine fumarate extended-release tablets 8 mg. The terminal half-life of the active metabolite was not changed.

### **7.4 CYP2D6 Inhibitors**

The interaction with CYP2D6 inhibitors was not tested clinically. In poor metabolizers for CYP2D6, representing a maximum CYP2D6 inhibition,  $C_{max}$  and AUC of the active metabolite are increased 1.7- and 2-fold, respectively.

No dosing adjustments are recommended in the presence of CYP2D6 inhibitors.

### **7.5 Drugs Metabolized by Cytochrome P450**

*In vitro* data indicate that at therapeutic concentrations, the active metabolite of fesoterodine does not have the potential to inhibit or induce Cytochrome P450 enzyme systems [ see *Clinical Pharmacology ( 12.3)*  ].

### **7.6 Oral Contraceptives**

In the presence of fesoterodine, there are no clinically significant changes in the plasma concentrations of combined oral contraceptives containing ethinyl estradiol and levonorgestrel [ see *Clinical Pharmacology ( 12.3)*  ].

### **7.7 Warfarin**

A clinical study has shown that fesoterodine extended-release tablets 8 mg once daily has no significant effect on the pharmacokinetics or the anticoagulant activity (PT/INR) of warfarin 25 mg. Standard therapeutic monitoring for warfarin should be continued [see *Clinical Pharmacology ( 12.3)*].

### **7.8 Drug-Laboratory Test Interactions**

Interactions between fesoterodine fumarate and laboratory tests have not been studied

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

There are no available data with the use of fesoterodine fumarate extended-release tablets in pregnant women and adolescents to evaluate for a drug associated risk of major birth defects or miscarriage, adverse maternal or fetal outcomes. In animal reproduction studies, oral administration of fesoterodine to pregnant mice and rabbits

during organogenesis resulted in fetotoxicity at maternal exposures that were 6 and 3 times, respectively, the maximum recommended human dose (MRHD) of 8 mg/day based on AUC (see Data). The background risk of major birth defects and miscarriage for the indicated population are unknown. However, in the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

## Data

### *Animal Data*

No dose-related teratogenicity was observed in reproduction studies performed in mice and rabbits. In mice at 6 to 27 times the expected exposure at the maximum recommended human dose (MRHD) of 8 mg based on AUC (75 mg/kg/day, oral), increased resorptions and decreased live fetuses were observed. One fetus with cleft palate was observed at each dose (15, 45, and 75 mg/kg/day), at an incidence within the background historical range. In rabbits treated at 3 to 11 times the MRHD (27 mg/kg/day, oral), incompletely ossified sternebrae (retardation of bone development) and reduced survival were observed in fetuses. In rabbits at 9 to 11 times the MRHD (4.5 mg/kg/day, subcutaneous), maternal toxicity and incompletely ossified sternebrae were observed in fetuses (at an incidence within the background historical range). In rabbits at 3 times the MRHD (1.5 mg/kg/day, subcutaneous), decreased maternal food consumption in the absence of any fetal effects was observed. Oral administration of 30 mg/kg/day fesoterodine to mice in a pre- and post-natal development study resulted in decreased body weight of the dams and delayed ear opening of the pups. No effects were noted on mating and reproduction of the F<sub>1</sub> dams or on the F<sub>2</sub> offspring.

## **8.2 Lactation**

### Risk Summary

There is no information on the presence of fesoterodine fumarate in human milk, the effects on the breastfed child, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for fesoterodine fumarate and any potential adverse effects on the breastfed child from fesoterodine fumarate or from the underlying maternal condition.

## **8.4 Pediatric Use**

The safety and effectiveness of fesoterodine fumarate have not been established in pediatric patients younger than 6 years of age or weighing 25 kg or less.

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

## **8.5 Geriatric Use**

No dose adjustment is recommended for the elderly. The pharmacokinetics of fesoterodine are not significantly influenced by age.

Of the 1,567 patients who received fesoterodine fumarate 4 mg or 8 mg orally once daily in Phase 2 and 3, placebo-controlled, efficacy and safety studies for OAB, 515 (33%) were 65 years of age or older, and 140 (9%) were 75 years of age or older. No overall difference in effectiveness was observed between patients younger than 65 years of age and those 65 years of age or older in these studies; However, the incidence of antimuscarinic adverse reactions, including dry mouth, constipation, dyspepsia, increase in residual urine, dizziness (at 8 mg only) and urinary tract infection, was higher

in patients 75 years of age and older as compared to younger patients [ see *Clinical Studies (14.1) and Adverse Reactions ( 6)*].

## **8.6 Renal Impairment**

In adult patients with severe renal impairment (CLCR <30 mL/min), C<sub>max</sub> and AUC are increased 2- and 2.3-fold, respectively. Doses of fesoterodine fumarate extended-release tablets greater than 4 mg are not recommended in adult patients with severe renal impairment. In patients with mild or moderate renal impairment (CL<sub>CR</sub> ranging from 30 to 80 mL/min), C<sub>max</sub> and AUC of the active metabolite are increased up to 1.5- and 1.8-fold, respectively, as compared to healthy subjects. No dose adjustment is recommended in patients with mild or moderate renal impairment [ see *Clinical Pharmacology (12.3), and Dosage and Administration ( 2.3)* ].

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

## **8.7 Hepatic Impairment**

Patients with severe hepatic impairment (Child-Pugh C) have not been studied; therefore fesoterodine fumarate is not recommended for use in these patients. In patients with moderate (Child-Pugh B) hepatic impairment, C<sub>max</sub> and AUC of the active metabolite are increased 1.4- and 2.1-fold, respectively, as compared to healthy subjects. No dose adjustment is recommended in patients with mild or moderate hepatic impairment [ see *Clinical Pharmacology ( 12.3)*].


## **10 OVERDOSAGE**

Overdosage with fesoterodine fumarate can result in severe anticholinergic effects. Treatment should be symptomatic and supportive. In the event of overdosage, ECG monitoring is recommended.

## **11 DESCRIPTION**

Fesoterodine fumarate extended-release tablets contains fesoterodine fumarate and is an extended-release tablet. Fesoterodine is rapidly de-esterified to its active metabolite (R)-2-(3-diisopropylamino-1-phenylpropyl)-4-hydroxymethyl-phenol, or 5-hydroxymethyl tolterodine, which is a muscarinic receptor antagonist.

Chemically, fesoterodine fumarate is designated as isobutyric acid 2-((R)-3-diisopropylammonium-1-phenylpropyl)-4-(hydroxymethyl) phenyl ester hydrogen fumarate. The molecular formula is C<sub>30</sub>H<sub>41</sub>NO<sub>7</sub> and its molecular weight is 527.4. The structural formula is:



The asterisk (\*) indicates the chiral carbon.

Fesoterodine fumarate is a white crystalline powder, which is freely soluble in water. Each Fesoterodine fumarate extended-release tablet contains either 4 mg or 8 mg of fesoterodine fumarate and the following inactive ingredients: FD&C Blue #2, fructose, glyceryl behenate, hypromellose, lactose anhydrous, lecithin soya, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc and titanium dioxide.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Fesoterodine is a competitive muscarinic receptor antagonist. After oral administration, fesoterodine is rapidly and extensively hydrolyzed by nonspecific esterases to its active metabolite, 5-hydroxymethyl tolterodine, which is responsible for the antimuscarinic activity of fesoterodine.

Muscarinic receptors play a role in contractions of urinary bladder smooth muscle. Inhibition of these receptors in the bladder is presumed to be the mechanism by which fesoterodine produces its effects.

### 12.2 Pharmacodynamics

In a urodynamic study involving patients with involuntary detrusor contractions, the effects after the administration of fesoterodine on the volume at first detrusor contraction and bladder capacity were assessed. Administration of fesoterodine increased the volume at first detrusor contraction and bladder capacity in a dose-dependent manner. These findings are consistent with an antimuscarinic effect on the bladder.

**Cardiac Electrophysiology:** The effect of fesoterodine 4 mg and 28 mg on the QT interval was evaluated in a double-blind, randomized, placebo- and positive-controlled (moxifloxacin 400 mg once a day) parallel trial with once-daily treatment over a period of 3 days in 261 male and female subjects aged 44 to 65 years. Electrocardiographic parameters were measured over a 24-hour period at pre-dose, after the first administration, and after the third administration of study medication. Fesoterodine 28 mg was chosen because this dose, when administered to CYP2D6 extensive metabolizers, results in an exposure to the active metabolite that is similar to the exposure in a CYP2D6 poor metabolizer receiving fesoterodine 8 mg together with CYP3A4 blockade. Corrected QT intervals (QTc) were calculated using Fridericia's correction and a linear individual correction method. Analyses of 24-hour average QTc, time-matched baseline-corrected QTc, and time-matched placebo-subtracted QTc intervals indicate that fesoterodine at doses of 4 and 28 mg/day did not prolong the QT

interval. The sensitivity of the study was confirmed by positive QTc prolongation by moxifloxacin.

In this study, conducted in subjects aged 44 to 65 years, fesoterodine fumarate was associated with an increase in heart rate that correlates with increasing dose. When compared to placebo, the mean increase in heart rate associated with fesoterodine 4 mg/day and fesoterodine 28 mg/day was 3 beats/minute and 11 beats/minute, respectively.

In the two, phase 3, placebo-controlled studies in adult in patients with overactive bladder, the mean increases in heart rate compared to placebo were 3 to 4 beats/minute in the fesoterodine 4 mg/day group and 3 to 5 beats/minute in the fesoterodine 8 mg/day group.

### 12.3 Pharmacokinetics

**Absorption:** After oral administration, fesoterodine is well absorbed. Due to rapid and extensive hydrolysis by nonspecific esterases to its active metabolite 5-hydroxymethyl tolterodine, fesoterodine cannot be detected in plasma. Bioavailability of the active metabolite is 52%. After single or multiple-dose oral administration of fesoterodine in doses from 4 mg to 28 mg, plasma concentrations of the active metabolite are proportional to the dose. Maximum plasma levels are reached after approximately 5 hours. No accumulation occurs after multiple-dose administration.

A summary of pharmacokinetic parameters for the active metabolite after a single dose of fesoterodine fumarate 4 mg and 8 mg in extensive and poor metabolizers of CYP2D6 is provided in Table 8.

**Table 8: Summary of geometric mean [CV] pharmacokinetic parameters for the active metabolite after a single dose of fesoterodine fumarate 4 mg and 8 mg in extensive and poor CYP2D6 metabolizers**

Parameter	Fesoterodine Fumarate 4 mg		Fesoterodine Fumarate 8 mg	
	EM (N=16)	PM (N=8)	EM (N=16)	PM (N=8)
C <sub>max</sub> (ng/mL)	1.89 [43%]	3.45 [54%]	3.98 [28%]	6.90 [39%]
AUC <sub>0-tz</sub> (ng*h/mL)	21.2 [38%]	40.5 [31%]	45.3 [32%]	88.7 [36%]
t <sub>max</sub> (h) <sup>a</sup>	5 [2 to 6]	5 [5 to 6]	5 [3 to 6]	5 [5 to 6]
t <sub>1/2</sub> (h)	7.31 [27%]	7.31 [30%]	8.59 [41%]	7.66 [21%]

EM = extensive CYP2D6 metabolizer, PM = poor CYP2D6 metabolizer, CV = coefficient of variation C<sub>max</sub> = maximum plasma concentration, AUC<sub>0-tz</sub> = area under the concentration time curve from zero up to the last measurable plasma concentration, t<sub>max</sub> = time to reach C<sub>max</sub>, t<sub>1/2</sub> = terminal half-life<sup>a</sup> Data presented as median (range)

**Effect of Food:** There is no clinically relevant effect of food on the pharmacokinetics of fesoterodine. In a study of the effects of food on the pharmacokinetics of fesoterodine in 16 healthy male volunteers, concomitant food intake increased the active metabolite of fesoterodine AUC by approximately 19% and C<sub>max</sub> by 18% [ see *Dosage and Administration* ( 2.1)].

**Distribution:** Plasma protein binding of the active metabolite is low (approximately 50%) and is primarily bound to albumin and alpha-1-acid glycoprotein. The mean steady-state volume of distribution following intravenous infusion of the active metabolite is 169 L.

**Metabolism:** After oral administration, fesoterodine is rapidly and extensively

hydrolyzed to its active metabolite. The active metabolite is further metabolized in the liver to its carboxy, carboxy-N-desisopropyl, and N-desisopropyl metabolites via two major pathways involving CYP2D6 and CYP3A4. None of these metabolites contribute significantly to the antimuscarinic activity of fesoterodine.

**Variability in CYP2D6 Metabolism:** A subset of individuals (approximately 7% of Caucasians and approximately 2% of African Americans) are poor metabolizers for CYP2D6.  $C_{max}$  and AUC of the active metabolite are increased 1.7- and 2-fold, respectively, in CYP2D6 poor metabolizers, as compared to extensive metabolizers.

**Excretion:** Hepatic metabolism and renal excretion contribute significantly to the elimination of the active metabolite. After oral administration of fesoterodine, approximately 70% of the administered dose was recovered in urine as the active metabolite (16%), carboxy metabolite (34%), carboxy-N-desisopropyl metabolite (18%), or N-desisopropyl metabolite (1%), and a smaller amount (7%) was recovered in feces.

The terminal half-life of the active metabolite is approximately 4 hours following an intravenous administration. The apparent terminal half-life following oral administration is approximately 7 hours.

### **Pharmacokinetics in Specific Populations Geriatric Patients:\**

**Geriatric Patients:** Following a single 8 mg oral dose of fesoterodine, the mean ( $\pm$ SD) AUC and  $C_{max}$  for the active metabolite 5-hydroxymethyl tolterodine in 12 elderly men (mean age 67 years) were  $51.8 \pm 26.1$  h\*ng/mL and  $3.8 \pm 1.7$  ng/mL, respectively. In the same study, the mean ( $\pm$ SD) AUC and  $C_{max}$  in 12 young men (mean age 30 years) were  $52 \pm 31.5$  h\*ng/mL and  $4.1 \pm 2.1$  ng/mL, respectively. The pharmacokinetics of fesoterodine were not significantly influenced by age [ see *Use in Specific Populations ( 8.5)*  ].

**Gender:** Following a single 8 mg oral dose of fesoterodine, the mean ( $\pm$ SD) AUC and  $C_{max}$  for the active metabolite 5-hydroxymethyl tolterodine in 12 elderly men (mean age 67 years) were  $51.8 \pm 26.1$  h\*ng/mL and  $3.8 \pm 1.7$  ng/mL, respectively. In the same study, the mean ( $\pm$ SD) AUC and  $C_{max}$  in 12 elderly women (mean age 68 years) were  $56 \pm 28.8$  h\*ng/mL and  $4.6 \pm 2.3$  ng/mL, respectively. The pharmacokinetics of fesoterodine were not significantly influenced by gender.

**Race:** The effects of Caucasian or Black race on the pharmacokinetics of fesoterodine were examined in a study of 12 Caucasian and 12 Black African young male volunteers. Each subject received a single oral dose of 8 mg fesoterodine. The mean ( $\pm$ SD) AUC and  $C_{max}$  for the active metabolite 5-hydroxymethyl tolterodine in Caucasian males were  $73 \pm 27.8$  h\*ng/mL and  $6.1 \pm 2.7$  ng/mL, respectively. The mean ( $\pm$ SD) AUC and  $C_{max}$  in Black males were  $65.8 \pm 23.2$  h\*ng/mL and  $5.5 \pm 1.9$  ng/mL, respectively . The pharmacokinetics of fesoterodine were not significantly influenced by race.

**Renal Impairment:** In patients with mild or moderate renal impairment ( $CL_{CR}$  ranging from 30 to 80 mL/min),  $C_{max}$  and AUC of the active metabolite are increased up to 1.5- and 1.8-fold, respectively, as compared to healthy subjects. In patients with severe renal impairment ( $CL_{CR} < 30$  mL/min),  $C_{max}$  and AUC are increased 2- and 2.3-fold, respectively. [see *Use in Specific Populations (8.6), Warnings and Precautions (5), and Dosage and Administration (2.3)*].

**Hepatic Impairment:** In patients with moderate (Child-Pugh B) hepatic impairment,  $C_{max}$  and AUC of the active metabolite are increased 1.4- and 2.1-fold, respectively, as compared to healthy subjects.

Subjects with severe hepatic impairment (Child-Pugh C) have not been studied [ see *Use in Specific Populations (8.7)*].

## **Drug-Drug Interactions**

**Drugs Metabolized by Cytochrome P450:** At therapeutic concentrations, the active metabolite of fesoterodine does not inhibit CYP1A2, 2B6, 2C8, 2C9, 2C19, 2D6, 2E1, or 3A4, or induce CYP1A2, 2B6, 2C9, 2C19, or 3A4 in vitro [ see *Drug Interactions* ( 7.5) ].

**CYP3A4 Inhibitors:** Following blockade of CYP3A4 by coadministration of the strong CYP3A4 inhibitor ketoconazole 200 mg twice a day for 5 days,  $C_{max}$  and AUC of the active metabolite of fesoterodine increased 2- and 2.3-fold, respectively, after oral administration of fesoterodine fumarate 8 mg to CYP2D6 extensive metabolizers. In CYP2D6 poor metabolizers,  $C_{max}$  and AUC of the active metabolite of fesoterodine increased 2.1- and 2.5-fold, respectively, during coadministration of ketoconazole 200 mg twice a day for 5 days.  $C_{max}$  and AUC were 4.5- and 5.7-fold higher, respectively, in subjects who were CYP2D6 poor metabolizers and taking ketoconazole compared to subjects who were CYP2D6 extensive metabolizers and not taking ketoconazole. In a separate study coadministering fesoterodine with ketoconazole 200 mg once a day for 5 days, the  $C_{max}$  and AUC values of the active metabolite of fesoterodine were increased 2.2-fold in CYP2D6 extensive metabolizers and 1.5- and 1.9-fold, respectively, in CYP2D6 poor metabolizers.  $C_{max}$  and AUC were 3.4- and 4.2-fold higher, respectively, in subjects who were CYP2D6 poor metabolizers and taking ketoconazole compared to subjects who were CYP2D6 extensive metabolizers and not taking ketoconazole.

There is no clinically relevant effect of moderate CYP3A4 inhibitors on the pharmacokinetics of fesoterodine. In a drug-drug interaction study evaluating the coadministration of the moderate CYP3A4 inhibitor fluconazole 200 mg twice a day for 2 days, a single 8 mg dose of fesoterodine was administered 1 hour following the first dose of fluconazole on day 1 of the study. The average (90% confidence interval) for the increase in  $C_{max}$  and AUC of the active metabolite of fesoterodine was approximately 19% (11% to 28%) and 27% (18% to 36%) respectively.

The effect of weak CYP3A4 inhibitors (e.g. cimetidine) was not examined; it is not expected to be in excess of the effect of moderate inhibitors [ see *Drug Interactions* (7.2)], *Warnings and Precautions* (5), and *Dosage and Administration* (2.3)].

**CYP3A4 Inducers:** Following induction of CYP3A4 by coadministration of rifampicin 600 mg once a day,  $C_{max}$  and AUC of the active metabolite of fesoterodine decreased by approximately 70% and 75%, respectively, after oral administration of fesoterodine fumarate 8 mg. The terminal half-life of the active metabolite was not changed.

Induction of CYP3A4 may lead to reduced plasma levels. No dosing adjustments are recommended in the presence of CYP3A4 inducers [ see *Drug Interactions* ( 7.3) ].

**CYP2D6 Inhibitors:** The interaction with CYP2D6 inhibitors was not studied. In poor metabolizers for CYP2D6, representing a maximum CYP2D6 inhibition,  $C_{max}$  and AUC of the active metabolite are increased 1.7- and 2-fold, respectively. [ see *Drug Interactions* ( 7.4) ].

**Oral Contraceptives:** Thirty healthy female subjects taking an oral contraceptive containing 0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel were evaluated in a 2-period crossover study. Each subject was randomized to receive concomitant administration of either placebo or fesoterodine 8 mg once daily on days 1 to 14 of hormone cycle for 2 consecutive cycles. Pharmacokinetics of ethinyl estradiol and levonorgestrel were assessed on day 13 of each cycle. Fesoterodine increased the AUC and  $C_{max}$  of ethinyl estradiol by 1 - 3% and decreased the AUC and  $C_{max}$  of levonorgestrel by 11 to 13% [ see *Drug Interactions* (7.6)].

**Warfarin:** In a cross-over study in 14 healthy male volunteers (18 to 55 years), a single oral dose of warfarin 25 mg was given either alone or on day 3 of once daily dosing for

9 days with fesoterodine 8 mg. Compared to warfarin alone dosing, the  $C_{max}$  and AUC of S-warfarin were lower by ~ 4 %, while the  $C_{max}$  and AUC of R-warfarin were lower by approximately 8 % and 6% for the co-administration, suggesting absence of a significant pharmacokinetic interaction. There were no statistically significant changes in the measured pharmacodynamic parameters for anti-coagulant activity of warfarin ( $INR_{max}$ ,  $AUC_{INR}$ ), with only a small decrease noted in  $INR_{max}$  of ~ 3 % with the co-administration relative to warfarin alone.  $INR$  versus time profiles across individual subjects in the study suggested some differences following co-administration with fesoterodine, although there was no definite trend with regard to the changes noted [ see *Drug Interactions* ( 7.7) ].

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment Of Fertility**

#### Carcinogenicity

No evidence of drug-related carcinogenicity was found in 24 month studies with oral administration to mice and rats. The highest tolerated doses in mice (females 45 to 60 mg/kg/day, males 30 to 45 mg/kg/day) correspond to 11 to 19 times (females) and 4 to 9 times (males) the estimated human AUC values reached with fesoterodine 8 mg, which is the Maximum Recommended Human Dose (MRHD). In rats, the highest tolerated dose (45 to 60 mg/kg/day) corresponds to 3 to 8 times (females) and 3 to 14 times (males) the estimated human AUC at the MRHD.

#### Mutagenesis

Fesoterodine was not mutagenic or genotoxic *in vitro* (Ames tests, chromosome aberration tests) or *in vivo* (mouse micronucleus test).

#### Impairment of Fertility

Fesoterodine had no effect on male reproductive function or fertility at doses up to 45 mg/kg/day in mice. At 45 mg/kg/day, a lower number of corpora lutea, implantation sites and viable fetuses was observed in female mice administered fesoterodine for 2 weeks prior to mating and continuing through day 7 of gestation. The maternal No-Observed-Effect Level (NOEL) and the NOEL for effects on reproduction and early embryonic development were both 15 mg/kg/day. At the NOEL, the systemic exposure, based on AUC, was 0.6 to 1.5 times higher in mice than in humans at the MRHD, whereas based on peak plasma concentrations, the exposure in mice was 5 to 9 times higher.

## **14 CLINICAL STUDIES**

### **14.1 Adult Overactive Bladder**

The efficacy of fesoterodine fumarate extended-release tablets was evaluated in two, Phase 3, randomized, double-blind, placebo-controlled, 12-week studies for the treatment of overactive bladder with symptoms of urge urinary incontinence, urgency, and urinary frequency. Entry criteria required that patients have symptoms of overactive bladder for  $\geq$  6-months duration, at least 8 micturitions per day, and at least 6 urinary urgency episodes or 3 urge incontinence episodes per 3-day diary period.

Patients were randomized to a fixed dose of fesoterodine fumarate 4 or 8 mg/day or placebo. In one of these studies, 290 patients were randomized to an active control arm (an oral antimuscarinic agent). For the combined studies, a total of 554 patients received placebo, 554 patients received fesoterodine fumarate 4 mg/day, and 566 patients received fesoterodine fumarate 8 mg/day. The majority of patients were Caucasian (91%) and female (79%) with a mean age of 58 years (range 19 to 91 years).

The primary efficacy endpoints were the mean change in the number of urge urinary incontinence episodes per 24 hours and the mean change in the number of micturitions (frequency) per 24 hours. An important secondary endpoint was the mean change in the voided volume per micturition.

Results for the primary endpoints and for mean change in voided volume per micturition from the two 12 week clinical studies of fesoterodine fumarate are reported in Table 10.

**Table 10: Mean baseline and change from baseline to Week 12 for urge urinary incontinence episodes, number of micturitions, and volume voided per micturition**

Study 1				Study 2		
Parameter	Placebo N=279	Fesoterodine Fumarate 4 mg/day N=265	Fesoterodine Fumarate 8 mg/day N=276	Placebo N=266	Fesoterodine Fumarate 4 mg/day N=267	Fesoterodine Fumarate 8 mg/dayN=267
Number of urge incontinence episodes per 24 hours <sup>a</sup>						
Baseline	3.7	3.8	3.7	3.7	3.9	3.9
Change from baseline	-1.20	-2.06	-2.27	-1	-1.77	-2.42
p-value vs. placebo	-	0.001	<0.001	-	<0.003	<0.001
Number of micturitions per 24 hours						
Baseline	12	11.6	11.9	12.2	12.9	12
Change from baseline	-1.02	-1.74	-1.94	-1.02	-1.86	-1.94
p-value vs. placebo	-	<0.001	<0.001	-	0.032	<0.001
Voided volume per micturition (mL)						
Baseline	150	160	154	159	152	156
Change from baseline	10	27	33	8	17	33
p-value vs. placebo	-	<0.001	<0.001	-	0.150	<0.001

vs. = versus

<sup>a</sup> Only those patients who were urge incontinent at baseline were included for the

analysis of number of urge incontinence episodes per 24 hours: In Study 1, the number of these patients was 211, 199, and 223 in the placebo, fesoterodine fumarate 4 mg/day and fesoterodine fumarate 8 mg/day groups, respectively. In Study 2, the number of these patients was 205, 228, and 218, respectively.

Figures 1-4: The following figures show change from baseline over time in number of micturitions and urge urinary incontinence episodes per 24 h in the two studies.

Figure 1: Change in Number of Micturitions per 24 h (Study 1)




Figure 2: Change in Urge Incontinence Episodes per 24 h (Study 1)




Figure 3: Change in Number of Micturations per 24 h (Study 2)





Figure 4: Change in Urge Incontinence Episodes per 24 h (Study 2)



A reduction in number of urge urinary incontinence episodes per 24 hours was observed for both doses as compared to placebo as early as two weeks after starting fesoterodine fumarate therapy.

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Fesoterodine fumarate extended-release tablets, 4 mg are light blue colored, oval shaped, biconvex, film-coated, debossed with '4' on one side and 'FS' on the other side and are supplied in bottles of 30's.

Bottles of 30 NDC 51407-661-30

Fesoterodine fumarate extended-release tablets, 8 mg are blue colored, oval shaped, biconvex, film-coated, debossed with '8' on one side and 'FT' on the other side and are supplied in bottles of 30's.

Bottles of 30 NDC 51407-662-30

Store fesoterodine fumarate extended-release tablets, at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from moisture.

## **17 PATIENT COUNSELING INFORMATION**

Advise the patient to read the FDA-Approved Patient Labeling (Patient Information)

### **Angioedema**

Inform patients and/or their caregivers that fesoterodine fumarate extended-release tablets may cause angioedema, which could result in life-threatening airway obstruction. Advise patients and/or their caregivers to promptly discontinue fesoterodine fumarate extended-release tablets and seek immediate medical attention if they experience edema of the lips, tongue or laryngopharynx, or difficulty breathing.

### **Antimuscarinic Effects**

Inform patients that fesoterodine fumarate, like other antimuscarinic agents, may produce clinically significant adverse effects related to antimuscarinic pharmacological activity including constipation and urinary retention. Fesoterodine fumarate, like other antimuscarinics, may be associated with blurred vision, therefore, patients should be advised to exercise caution in decisions to engage in potentially dangerous activities until the drug's effects on the patient have been determined. Heat prostration (due to decreased sweating) can occur when fesoterodine fumarate, like other antimuscarinic drugs, is used in a hot environment.

### **Alcohol**

Patients should also be informed that alcohol may enhance the drowsiness caused by fesoterodine fumarate, like other anticholinergic agents. Patients should read the patient leaflet entitled "Patient Information Fesoterodine Fumarate Extended-release Tablets" before starting therapy with fesoterodine fumarate.

R x only

Distributor:

**Dr. Reddy's Laboratories Inc.,**

Princeton, NJ 08540

**Made in India**

Issued: 03/2022

Marketed/Packaged by:

GSMS, Inc.

## **Patient Information**

### **Fesoterodine Fumarate Extended-release Tablets, for oral use**

(FES oh TER oh deen)

Read the Patient Information that comes with fesoterodine fumarate extended-release tablets before you start taking it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your healthcare provider about your medical condition or your treatment.

#### **What are fesoterodine fumarate extended-release tablets?**

Fesoterodine fumarate extended-release tablets are a prescription medicine used:

- in adults to treat symptoms of a condition called overactive bladder (OAB), including urge urinary incontinence (leaking or wetting accidents due to a strong need to urinate), urinary urgency (having a strong need to urinate right away), or urinary frequency (having to urinate too often).

It is not known if fesoterodine fumarate is safe and effective in children younger than 6 years of age or with a body weight 55 pounds (25 kg) or less.

#### **Who should not take fesoterodine fumarate extended-release tablets?**

Do not take fesoterodine fumarate extended-release tablets if you:

- are allergic to fesoterodine fumarate or any of its ingredients. See the end of this leaflet for a complete list of ingredients.
- are allergic to tolterodine tartrate tablets or tolterodine tartrate extended-release capsules.
- are not able to empty your bladder (urinary retention).
- have delayed or slow emptying of your stomach (gastric retention).
- have an eye problem called uncontrolled narrow-angle glaucoma.

#### **Before you take fesoterodine fumarate extended-release tablets, tell your healthcare provider about all your medical conditions, including if you:**

- have problems emptying your bladder or you have a weak urine stream.
- have any stomach or intestinal problems, or problems with constipation.
- are receiving treatment for an eye problem called narrow-angle glaucoma.
- have a condition called Myasthenia Gravis.
- have kidney problems
- have liver problems
- are pregnant or plan to become pregnant. It is not known if fesoterodine fumarate extended-release tablets will harm your unborn baby. Talk to your healthcare provider if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if fesoterodine fumarate passes into your breast milk. You should talk to your healthcare provider about the best way to feed your baby while taking fesoterodine fumarate extended-release tablets.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal products. Fesoterodine fumarate extended-release tablets may affect the way other medicines work, and other medicines may affect how fesoterodine fumarate extended-release

tablets work. Especially tell your healthcare provider if you are taking antimucarinc, antibiotics, or antifungal medicines.

Know all the medicines you take. Keep a list of them with you to show your healthcare provider and pharmacist each time you get a new medicine.

### **How should I take fesoterodine fumarate extended-release tablets?**

- Take fesoterodine fumarate extended-release tablets exactly as your healthcare provider tells you to take it.
- Your healthcare provider may lower your dose of fesoterodine fumarate extended-release tablets if you are an adult with severe kidney problems.
- Take fesoterodine fumarate extended-release tablets with liquid and swallow the tablet whole. Do not chew, divide, or crush the tablet.
- Take fesoterodine fumarate extended-release tablets with or without food.
- If you miss a dose of fesoterodine fumarate extended-release tablets, begin taking fesoterodine fumarate extended-release tablets again the next day. Do not take 2 doses of fesoterodine fumarate extended-release tablets in the same day.
- If you take too much fesoterodine fumarate extended-release tablets, call your healthcare provider or go to an emergency department right away.

### **What should I avoid while taking Fesoterodine fumarate extended-release tablets?**

- Fesoterodine fumarate extended-release tablets can cause blurred vision, dizziness, and drowsiness. Do not drive, operate machinery, or do other dangerous activities until you know how fesoterodine fumarate affects you.
- Use caution in hot environments. Decreased sweating and severe heat illness can happen when medicines such as fesoterodine fumarate extended-release tablets are used in a hot environment.
- Drinking alcohol while taking medicines such as fesoterodine fumarate extended-release tablets may cause increased drowsiness.

### **What are the possible side effects of fesoterodine fumarate extended-release tablets?**

#### **Fesoterodine fumarate extended-release tablets may cause serious side effects, including:**

- **Serious allergic reactions.** Symptoms of a serious allergic reaction may include swelling of the face, lips, throat or tongue. If you have these symptoms, you should stop taking fesoterodine fumarate extended-release tablets and get emergency medical help right away.
- **inability to empty bladder (urinary retention).** Fesoterodine fumarate extended-release tablets may increase your chances of not being able to empty your bladder if you have bladder outlet obstruction. Tell your healthcare provider right away if you are unable to empty your bladder.
- **central nervous system (CNS) effects.** Talk to your healthcare provider right away if you get any of these side effects: headache, dizziness, and drowsiness.
- **worsening of Myasthenia Gravis symptoms.** The most common side effects of fesoterodine fumarate extended-release tablets in adults include:

- dry mouth
- constipation

Talk to your healthcare provider about any side effect that bothers you or that does not go away. These are not all the possible side effects of fesoterodine fumarate extended-release tablets. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store fesoterodine fumarate extended-release tablets?**

- Store fesoterodine fumarate extended-release tablets between 68°F to 77°F (20°C to 25°C).
- Protect the medicine from moisture by keeping the bottle closed tightly.
- **Keep fesoterodine fumarate extended-release tablets and all medicines out of the reach of children.**

### **General information about the safe and effective use of fesoterodine fumarate extended-release tablets**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use fumarate extended-release tablets for a condition for which it was not prescribed. Do not give fesoterodine fumarate extended-release tablets to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about fesoterodine fumarate extended-release tablets that is written for health professionals.

### **What are the ingredients in fesoterodine fumarate extended-release tablets?**

**Active ingredient:** fesoterodine fumarate

**Inactive ingredients:** FD&C Blue #2, fructose, glyceryl behenate, hypromellose, lactose anhydrous, lecithin soya, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc and titanium dioxide.

*Pediatric use information is approved for Pfizer Inc.'s TOVIAZ® (fesoterodine fumarate) extended-release tablets. However, due to Pfizer Inc.'s marketing exclusivity rights, this drug product is not labeled with that information.*

Trademarks are the property of their respective owners.

To reorder additional Patient Information Leaflets, please contact Dr. Reddy's Customer Service at 1-866-733-3952.

R x only

Distributor:

**Dr. Reddy's Laboratories Inc.,**

Princeton, NJ 08540

**Made in India**

Issued: 03/2022

Marketed/Packaged by:

GSMS, Inc.

Camarillo, CA USA 93012

Fesoterodine fumarate extended-release tablets, 4mg

Container Label - 30's Count

GTIN :  
LOT :  
EXP :  
SN :

NDC 51407-**661**-30

**Fesoterodine Fumarate**  
Extended-Release Tablets

**4 mg**

PHARMACIST: Dispense the Patient Information to each patient.  
Print Patient Information at: <https://gsms.us/>

**GSMS**<sup>®</sup> **30 tablets**  
incorporated Rx Only

5140766130

Each film-coated extended-release tablet contains: fesoterodine fumarate 4 mg.

**DOSAGE:** See accompanying Prescribing Information.

Protect from moisture.

Swallow whole. Do not chew, divide, or crush the tablet.

**Keep this and all medication out of the reach of children.**

Store at 20° - 25°C (68° - 77°F); [See USP Controlled Room Temperature].

Manufactured by:  
Dr. Reddy's Laboratories Limited  
Marketed by: GSMS, Incorporated  
Camarillo, CA 93012 USA Rev. 07/22

Fesoterodine fumarate extended-release tablets, 8 mg

Container Label - 30's Count

GTIN :  
LOT :  
EXP :  
SN :

NDC 51407-**662**-30

**Fesoterodine Fumarate**  
Extended-Release Tablets

**8 mg**

PHARMACIST: Dispense the Patient Information to each patient.  
Print Patient Information at: <https://gsms.us/>

**GSMS**<sup>®</sup> **30 tablets**  
incorporated Rx Only

5140766230

Each film-coated extended-release tablet contains: fesoterodine fumarate 8 mg.

**DOSAGE:** See accompanying Prescribing Information.

Protect from moisture.

Swallow whole. Do not chew, divide, or crush the tablet.

**Keep this and all medication out of the reach of children.**

Store at 20° - 25°C (68° - 77°F); [See USP Controlled Room Temperature].

Manufactured by:  
Dr. Reddy's Laboratories Limited  
Marketed by: GSMS, Incorporated  
Camarillo, CA 93012 USA Rev. 07/22

**FESOTERODINE FUMARATE**

fesoterodine fumarate tablet, extended release

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:51407-661(NDC:43598-247)
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>FESOTERODINE FUMARATE</b> (UNII: EOS72165S7) (FESOTERODINE - UNII:621G617227)	FESOTERODINE FUMARATE	4 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>FRUCTOSE</b> (UNII: 6YSS42VSEV)	
<b>GLYCERYL DIBEHENATE</b> (UNII: R8WTH25YS2)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	

<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)

Product Characteristics			
<b>Color</b>	blue (light blue)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	13mm
<b>Flavor</b>		<b>Imprint Code</b>	Drreddyslogo;FS
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51407-661-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	07/07/2022	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204975	08/13/2019	

## FESOTERODINE FUMARATE

fesoterodine fumarate tablet, extended release

Product Information			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:51407-662(NDC:43598-248)
<b>Route of Administration</b>	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>FESOTERODINE FUMARATE</b> (UNII: EOS72165S7) (FESOTERODINE - UNII:621G617227)	FESOTERODINE FUMARATE	8 mg

Inactive Ingredients	
Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>FRUCTOSE</b> (UNII: 6YSS42VSEV)	
<b>GLYCERYL DIBEHENATE</b> (UNII: R8WTH25YS2)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	

<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	13mm
<b>Flavor</b>		<b>Imprint Code</b>	Drreddyslogo;FT
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51407-662-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	07/07/2022	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204975	08/13/2019	

**Labeler - Golden State Medical Supply, Inc. (603184490)**

**Establishment**

Name	Address	ID/FEI	Business Operations
Golden State Medical Supply, Inc.		603184490	repack(51407-661, 51407-662) , relabel(51407-662, 51407-661)