

**REGENCY PRE AND POST UDDER WASH- phenol udder wash solution**  
**CLSolutions**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----

**Regency Pre and Post Udder Wash**

**ACTIVE INGREDIENTS:**

<b>Active Ingredients</b>	<b>Purpose</b>
PHENOL (0.78% w/v)	Disinfectant
PHENOLATE SODIUM (0.03% w/v)	Disinfectant

**PURPOSE:**

<b>Active Ingredients</b>	<b>Purpose</b>
PHENOL (0.78% w/v)	Disinfectant
PHENOLATE SODIUM (0.03% w/v)	Disinfectant

**Keep Out of Reach of Children.**

If swallowed, get medical help or contact Poison Control Center right away.

**USES:**

**FOR ANIMAL USE ONLY**

Aids in controlling mastitis organisms with a softening effect on teats

**Directions for Use:**

Wash udder and teats with damp cloth to remove foreign debris.

Apply 0.5X Pre-Milking Teat Dip.

Allow product to disinfect teats during the time it takes to prep or milk preceding cows.

Wash teats by hand with clean damp cloth to remove product prior to milking.

**CAUTION:**

*Avoid eye contact. For external use only.*

**FIRST AID:**

*For Ingestion - Immediately give castor oil or vegetable oil (15-30cc). Get medical attention immediately.*

*Be ready to induce vomiting at the advice of a physician or poison control center.*

*For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.*

*When using this product:*

Do Not Store in Direct Sunlight.

Avoid Excessive Heat.

### User Safety Warnings

**Do Not Allow Product To Freeze.** If frozen, allow the product to completely thaw in a warm room.

Shake and thoroughly mix product before using.

Rinse teat dip cup with water after each milking and invert to dry.

Never pour used teat dip into original container.

Keep cap on container when not in use.

### INACTIVE INGREDIENTS:

Water, Glycerin, Linear Alkylbenzene Sulfonate, Sodium Borate, Fragrance, FD&C Blue #1



**Regency**  
Masticide®  
"Ready To Use Udder Wash"

with   
Sporicidin®  
Technology

**Pre & Post Udder Wash**

Aids in controlling mastitis organisms with a softening effect on teats.  
**Actives: 0.78% (w/v) Phenol with 0.03% (w/v) Sodium Phenate**  
Udder Wash Will Not Burn, Dry, Irritate Or Chap Teats.  
Contains a special blend of emollients and phenol.

**NOTE:** Store at room temperature away from direct sunlight.  
**NET CONTENTS: 270 U.S. Gallons**

**FOR ANIMAL USE ONLY**  
**KEEP OUT OF REACH OF CHILDREN**  
**DO NOT STORE IN DIRECT SUNLIGHT**  
**AVOID EXCESSIVE HEAT**



**FIRST AID:** For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

**CAUTION:** Avoid eye contact. For external use only.

**DIRECTIONS:** Wash udder and teats with damp cloth to remove foreign debris. Apply Regency® Masticide® Pre & Post Udder Wash. Let the product disinfect teats during the time it takes to prep or milk preceding cows. **Wash teats by hand with clean damp cloth to remove product prior to milking.**

If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. **Shake and thoroughly mix product before using.**

**INGREDIENTS:** Water, Emollients, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Fragrance, FD&C Blue #1.

**Distributed Exclusively By:**  
CLSolutions, LLC™  
7031 North 16th St.  
Omaha, NE 68112  
(402) 502-3928  
www.clsag.com  
CLS-PPW-1

Sporicidin® is a registered trademark of Contec, Inc.  
United States Patent # U56,586,477 B1

**LOT NO.** \_\_\_\_\_

**Regency Masticide Pre and Post Udder Wash - Ready To Use Udder Wash with Sporidicin Technology**

Aids in controlling mastitis organisms with a softening effect on teats.

**0.78% (w/v) Phenol with 0.03% (w/v) Sodium Phenate**

Teat Dip Will Not Burn, Dry, Irritate or Chap Teats. Contains special blend of emollients and phenol.

**NET CONTENTS: 270 U.S. Gallons**

# Regency<sup>®</sup> Masticide<sup>®</sup>

"Ready To Use Udder Wash"

## Pre & Post Udder Wash

Aids in controlling mastitis organisms with a softening effect on teats.

**Actives: 0.78% (w/v) Phenol with 0.03% (w/v) Sodium Phenate**

Udder Wash Will Not Burn, Dry, Irritate Or Chap Teats.

Contains a special blend of emollients and phenol.

**NOTE:** Store at room temperature away from direct sunlight.

**NET CONTENTS: 55 U.S. Gallons (208.2 L)**

with



**FOR ANIMAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN  
DO NOT STORE IN DIRECT SUNLIGHT  
AVOID EXCESSIVE HEAT**



**FIRST AID:** For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

**CAUTION:** Avoid eye contact. For external use only.

**DIRECTIONS:** Wash udder and teats with damp cloth to remove foreign debris. Apply Regency<sup>®</sup> Masticide<sup>®</sup> Pre & Post Udder Wash. Let the product disinfect teats during the time it takes to prep or milk preceding cows. **Wash teats by hand with clean damp cloth to remove product prior to milking.**

If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. **Shake and thoroughly mix product before using.**

**INGREDIENTS:** Water, Emollients, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Fragrance, FD&C Blue #1.

**Distributed Exclusively By:**  
CLSolutions, LLC<sup>™</sup>  
7031 North 16th St.  
Omaha, NE 68112  
(402) 502-3928  
www.clsag.com

Sporicidin<sup>®</sup> is a registered trademark of Contec, Inc.  
United States Patent # US6,586,477 B1  
CLS-PPW-55GAL-1



## Regency Masticide Pre and Post Udder Wash - Ready To Use Udder Wash with Sporicidin Technology

Aids in controlling mastitis organisms with a softening effect on teats.

**0.78% (w/v) Phenol with 0.03% (w/v) Sodium Phenate**

Teat Dip Will Not Burn, Dry, Irritate or Chap Teats. Contains special blend of emollients and phenol.

**NET CONTENTS: 55 Gallons (208.2 L)**

# Regency<sup>®</sup> Masticide<sup>®</sup>

"Ready To Use Udder Wash"

## Pre & Post Udder Wash

Aids in controlling mastitis organisms with a softening effect on teats.

**Actives: 0.78% (w/v) Phenol with 0.03% (w/v) Sodium Phenate**

Udder Wash Will Not Burn, Dry, Irritate Or Chap Teats.

Contains a special blend of emollients and phenol.

**NOTE:** Store at room temperature away from direct sunlight.

**NET CONTENTS: 5 U.S. Gallons (18.927 L)**

with



**FOR ANIMAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN  
DO NOT STORE IN DIRECT SUNLIGHT  
AVOID EXCESSIVE HEAT**



**FIRST AID:** For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

**CAUTION:** Avoid eye contact. For external use only.

**DIRECTIONS:** Wash udder and teats with damp cloth to remove foreign debris. Apply Regency<sup>®</sup> Masticide<sup>®</sup> Pre & Post Udder Wash. Let the product disinfect teats during the time it takes to prep or milk preceding cows. **Wash teats by hand with clean damp cloth to remove product prior to milking.**

If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. **Shake and thoroughly mix product before using.**

**INGREDIENTS:** Water, Emollients, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Fragrance, FD&C Blue #1.

**Distributed Exclusively By:**  
CLSolutions, LLC<sup>™</sup>  
7031 North 16th St.  
Omaha, NE 68112  
(402) 502-3928  
www.clsag.com

Sporicidin<sup>®</sup> is a registered trademark of Contec, Inc.  
United States Patent # US6,586,477 B1  
CLS-PPW-5GAL-1



## Regency Masticide Pre and Post Udder Wash - Ready To Use Udder Wash with Sporicidin Technology

Aids in controlling mastitis organisms with a softening effect on teats.

**0.78% (w/v) Phenol with 0.03% (w/v) Sodium Phenate**

Teat Dip Will Not burn, Dry, Irritate or Chap Teats, Contains a special blend of emollients and phenol.



**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86018-001-04	1021.95 L in 1 CONTAINER		
2	NDC:86018-001-03	208.2 L in 1 DRUM		
3	NDC:86018-001-02	18.927 L in 1 PAIL		
4	NDC:86018-001-01	3.785 L in 1 JUG		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/18/2017	

**Labeler** - CLSolutions (079338118)**Registrant** - ABS Corporation (048022727)**Establishment**

Name	Address	ID/FEI	Business Operations
ABS Corporation		048022727	api manufacture, pack, label, manufacture

Revised: 12/2017

CLSolutions