

GOOD SENSE ASPIRIN- aspirin tablet, chewable
L. Perrigo Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Perrigo Aspirin Drug Facts

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

- children under 12 years: consult a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

Questions or comments?

1-800-719-9260

Principal Display Panel

GOODSENSE®

Chewable

See New Warning

Aspirin Regimen

Talk to your doctor or other healthcare provider before using this product for your heart.

Aspirin 81 mg

Actual Size

Pain Reliever (NSAID)

Orange Flavor

Low Dose

36 Tablets

Compare to active ingredient of Bayer® Chewable Aspirin

46768 C2 C8



Drug Facts

Active ingredient (in each chewable tablet)
Aspirin 81 mg (NSAID)¹..... Pain reliever
¹nonsteroidal anti-inflammatory drug

Purpose
Pain reliever
Uses for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if

- you are taking a prescription drug for
 - gout
 - diabetes
 - arthritis
- Stop use and ask a doctor if**
 - an allergic reaction occurs. Seek medical help right away.
 - you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs
- These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store at 20-25°C (68-77°F)

GOODSENSE[®]

NDC 0113-0467-68

Chewable

See New Warning

Aspirin Regimen
Talk to your doctor or other healthcare provider before using this product for your heart.

Aspirin 81mg

Actual Size
Orange Flavor

Pain Reliever (NSAID)
Low Dose

Compare to active ingredient of Bayer[®] Chewable Aspirin

36 Tablets



how2recycle.info

Gluten Free
Distributed By
Perrigo[®]
Allergan, MI 49010

Questions or comments?

1-800-719-9260

Drug Facts (continue d)
Inactive ingredients: corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

GOOD SENSE ASPIRIN

aspirin tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-0467	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	81 mg	
Inactive Ingredients				
Ingredient Name			Strength	
STARCH, CORN (UNII: O8232NY3SJ)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
Product Characteristics				
Color	ORANGE	Score	no score	
Shape	ROUND	Size	8mm	
Flavor	ORANGE	Imprint Code	L467	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0467-68	1 in 1 CARTON	11/04/1992	
1		36 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0113-0467-08	3 in 1 CARTON	11/04/1992	
2		1 in 1 CARTON		
2		36 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part343	11/04/1992		

Labeler - L. Perrigo Company (006013346)