

ASPIRIN- aspirin tablet, film coated
CHAIN DRUG MARKETING ASSOCIATION INC

Quality Choice 44-157

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 63868-352-10

**QC®
QUALITY
CHOICE**

**†Compare to the
active ingredient in
Genuine BAYER® Aspirin**

**Aspirin
(NSAID)**

**Pain Reliever
Fever Reducer 325 mg**

Coated Tablets

100 Tablets

actual size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured
or distributed by Bayer AG, owner of
the registered trademark Genuine
Bayer® Aspirin.

50844 REV0122C15712

Distributed by CDMA, Inc.

43157 W Nine Mile

Novi, MI 48375

www.qualitychoice.com

Questions: 800-935-2362

NDC 63868-352-10



[†]Compare to the
active ingredient in
Genuine BAYER® Aspirin

Aspirin[®]

(NSAID)

Pain Reliever
Fever Reducer **325 mg**

Coated Tablets



actual size

No print/no varnish
Lot & Exp date

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin.

Distributed by CDMA, Inc.
43157 W Nine Mile
Novi, MI 48375
www.qualitychoice.com
Questions: 800-352-2562

SALE OF THIS PRODUCT IS GUARANTEED



6 35515 90431 6

TAMPER EVIDENCE: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

<i>Drug Facts</i>
Active ingredient (in each tablet) Aspirin 325 mg (NSAID)* Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug
Uses temporarily relieves: <ul style="list-style-type: none"> ■ minor pain of arthritis ■ headache ■ muscle pain ■ toothache ■ pain and fever of colds
Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ facial swelling ■ shock ■ hives ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take a blood thinning (anticoagulant) or steroid drug ■ have 3 or more alcoholic drinks every day while using this product ■ have had stomach ulcers or bleeding problems ■ take more or for a longer time than directed ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

PEEL HERE FOR MORE DRUG FACTS

<p>Drug Facts (continued)</p> <p>Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer</p> <p>■ if you have ever had an allergic reaction to this product or any of its ingredients</p> <hr/> <p>Ask a doctor before use if ■ you are taking a diuretic</p> <p>■ stomach bleeding warning applies to you</p> <p>■ you have a history of stomach problems, such as heartburn</p> <p>■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease</p> <p>■ you have asthma</p> <hr/> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis</p> <hr/> <p>Stop use and ask a doctor if ■ an allergic reaction occurs.</p> <p>Seek medical help right away:</p> <p>■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood</p> <p>■ have bloody or black stools</p> <p>■ have stomach pain that does not get better</p> <p>■ ringing in the ears or a loss of hearing occurs</p> <p>■ pain gets worse or lasts more than 10 days</p> <p>■ fever gets worse or lasts more than 3 days</p> <p>■ redness or swelling is present ■ new symptoms occur</p> <p>These could be signs of a serious condition.</p> <hr/> <p>If pregnant or breast-feeding, ask a health professional before use. It's especially important not to use aspirin at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <hr/> <p>Directions ■ do not take more than directed</p> <p>■ drink a full glass of water with each dose</p> <p>■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor</p> <p>■ children under 12 years: do not use unless directed by a doctor</p> <hr/> <p>Other information</p> <p>■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package</p> <hr/> <p>Inactive ingredients corn starch, hypromellose, polyethylene glycol, propylene glycol</p> <hr/> <p>Questions or comments? 1-800-426-9391</p>	<p>--- STOP PEELING ---</p> <p>Adhesive Only</p>
---	--

Quality Choice 44-157

ASPIRIN

aspirin tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-352
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;157;ASPIRIN	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-352-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	
2	NDC:63868-352-03	300 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013		02/10/1996	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(63868-352)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63868-352)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63868-352)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(63868-352)

Establishment			
Name	Address	ID/FEI	Business Operations

LNK International, Inc.		967626305	pack(63868-352)
-------------------------	--	-----------	-----------------

Revised: 11/2023

CHAIN DRUG MARKETING ASSOCIATION INC