PROPYLENE GLYCOL - propylene glycol liquid United Agricultural & Veterinary Products, Inc/Centaur, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

PROPYLENE GLYCOL

Feed Grade

For treatment and prevention of Acetonemia (Ketosis) in dairy cows.

For animal use only.

Keep out of reach of children.

INDICATIONS

An aid in the treatment of bovine ketosis (Acetonemia).

Propylene glycol is very palatable and up to 10% in feed is acceptable to dairy cows. Propylene glycol does not destroy rumen micro-organisms or leave undesirable residues after oxidation.

When administering propylene glycol to cows suffering from milk fever, mastitis or metritus, consult your veterinarian.

TAKE TIME OBSERVE LABEL DIRECTIONS

ADMINISTRATION

PREVENTION: As aid in prevention of Acetonemia (Ketosis) in dairy cattle, give 6 to 8 fluid ounces per head per day beginning 2 weeks before calving and continuing for 6 weeks after calving.

TREATMENT: As aid in treatment of Acetonemia (Ketosis) in dairy cattle, give 6 to 16 fluid ounces (dependent of severity) per head per day for 10 days.

Use as drench or mix with feed.

Store Between 59-86°F (15-30°C)

ACTIVE INGREDIENTS: Propylene Glycol ..



Propylene Glycol

INDICATIONS: An aid in the treatment of bovine ketosis

Propylene glycol is very palatable and up to 10% in feed is acceptable to dairy cows. Propylene glycol does not destroy rumen micro-organisms or leave undesirable residues after oxidation.

When administering propylene glycol to cows suffering from milk fever, mastitis or metritus, consult your





Feed Grade

For treatment and prevention of Acetonemia (Ketosis) in dairy cows.

For animal use only. Keep out of reach of children.

NET CONTENTS: 1 GALLON (128 FL. OZ.) 3.785 L

ADMINISTRATION:

PREVENTION: As aid in prevention of Acetonemia (Ketosis) in dairy cattle, give 6 to 8 fluid ounces per head per day. Beginning 2 weeks before calving and continuing for 6 weeks after calving.

TREATMENT: As aid in treatment of Acetonemia (Ketosis) in dairy cattle, give 6 to 16 fluid ounces (dependant of severity) per head per day for 10 days. Use as drench or mix with feed.

Store Between 59-86°F (15-30°C)

Manufactured By Centaur, Inc. Olathe, KS 66061 1-800-236-6180



Rev. 04-14

Lot #:

PROPYLENE GLYCOL

propylene glycol liquid

Product Information

Product Type OTC ANIMAL DRUG Item Code (Source) NDC:58305-055

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength PROPYLENE GLYCOL (UNII: 6DC9Q167V3) (PROPYLENE GLYCOL -**PROPYLENE** 3.785 L in 3.785 L UNII:6DC9Q167V3) GLYCOL

Packaging

Revised: 6/2014

# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:58305-055-80	3.785 L in 1 JUG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/12/2011	

Labeler - United Agricultural & Veterinary Products, Inc/Centaur, Inc. (174239087)