GOOD SENSE MIGRAINE FORMULA- acetaminophen, aspirin, caffeine tablet, film coated Proficient Rx LP

Migraine Formula Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID*)

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever aid

Use

• treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- · you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- · you have liver disease
- · stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever

- reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for:
- diabetes
- gout
- arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- · vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- close cap tightly after use

• read all product information before using. Keep this box for important information.

Inactive ingredients

carnauba wax, crospovidone, FD&C blue #1 aluminum lake, hypromellose, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, saccharin sodium, stearic acid, titanium dioxide

Questions comments?

1-800-719-9260

Principal Display Panel

GOOD SENSE®

Treats Migraine

Migraine Formula

Acetaminophen, Aspirin (NSAID) and Caffeine Tablets

Pain Reliever/Pain Reliever Aid

60 Coated Caplets**

**Capsule-Shaped Tablets

Compare to active ingredients of Excedrin® Migraine





NDC 82804-002-60

Packaged By: Proficient Rx LP Thousand Oaks, CA 91320

Migraine Formula 250mg /250mg /65mg #60 Caplets Lot #:00000 SN# MASTER NDC 82804-002-60 Exp:00/00/00

Migraine Formula 250mg /250mg /65mg #60 Caplets Lot #.00000 SN# MASTER NDC 82804-002-60 Exp:00/00/00

Migraine Formula 250mg /250mg /65mg #60 Caplets Lot #:00000 SN# MASTER NDC 82804-002-60 Exp:00/00/00



GTIN: 00382804002603 SN# MASTER Exp. 00/00/00 Lot #:00000

Migraine Formula 250mg /250mg /65mg

#60

Caplets

Each caplet contains: Acetaminophen 250 mg Pain reliever / Aspirin 250 mg (NSAID*) Pain reliever / Caffeine 65 mg Pain reliever aid *nonsteroidal anti-inflammatory drug

White, oval shaped, unscored caplet, with imprint code L374

Product ID: ZM000260 Dist. By: Perrigo Allegan, MI 49010 Store at 20°-25°C (68°-77°F)

Keep medication out of the reach of children

GOOD SENSE MIGRAINE FORMULA

acetaminophen, aspirin, caffeine tablet, film coated

Product	Information
----------------	--------------------

Product Type HUMAN OTC DRUG Item Code (Source) NDC:82804-002(NDC:0113-0374)

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	250 mg	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg	
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg	

Inactive Ingredients		
Ingredient Name	Strength	
CARNAUBA WAX (UNII: R12CBM0EIZ)		
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		

Product Characteristics			
Color	WHITE	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	L374
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82804-002- 30	30 in 1 BOTTLE; Type 0: Not a Combination Product	08/24/2023	
2	NDC:82804-002- 60	60 in 1 BOTTLE; Type 0: Not a Combination Product	08/24/2023	
3	NDC:82804-002- 90	90 in 1 BOTTLE; Type 0: Not a Combination Product	08/24/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075794	04/17/2002	

Labeler - Proficient Rx LP (079196022)

Establishment			
Name	Address	ID/FEI	Business Operations
Proficient Rx LP		079196022	REPACK(82804-002), RELABEL(82804-002)

Revised: 8/2023 Proficient Rx LP