RAY DOL- pain relieving cream cream Genomma Lab USA Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredients

Menthol - 10%

Methyl salicylate - 15%

Purpose

Topical Analgesic

Topical Analgesic

Uses

Temporary relief of minor aches and pains of muscle and joints associated with

- simple back aches
- arthritis
- strains
- bruises
- sprains

Warnings

For external use only

Do not use

- on wounds, damaged, broken or irritated skin
- with a heating pad

When using this product

- use only as directed
- avoid contact with eyes or mucous membranes
- Do not apply bandage tightly

Stop use and ask doctor if

- Condition worsens, or symtoms persist for more than 7 days
- symptoms clear up and occur again within a few days
- excessive skin irritation occurs
- **If pregnant or breast feeding**, ask a health professional before use.

Keep this and all drugs out of the reach of children. If swallowed, get medical help or contact a Poison Control Center immediately.

Directions

- **Adults and children 2 years of age and older**; Apply to affected area not more than 3 to 4 times daily.
- Children under 2 years of age; consult a doctor.

Other information

- Store at controlled room temperature 20°- 25°C (68° to 77°F)
- Close cap tightly after use.

Inactive ingredients

Carbomer, Cetyl Alcohol, Oleth-3 Phosphate, Purified Water, Stearic acid, Triethanolamine

Principal Display Panel - 3.0 oz Carton

RAY DOL

Topical Analgesic

Greaselees Stainless

Temporary Relief

Penetrating temporary relief of minor muscle aches

For Arthritis, Muscle & Joint Pain

10% Menthol and

15% Methyl Salicylate


Net WT. 3.0 oz (85g)



Principal Display Panel - 3.0 oz Tube RAY DOL

Pain Relieving Cream

Topical Analgesic



DRUG FACTS	
Active ingredients	Purpose
Menthol - 10%, Methyl Salicylate - 15%	Topical Analgesic
Hone: Tomporany rollief of minor achee and pains of muscoles	and joints associated with:

Uses: Temporary relief of minor aches and pains of muscles and joints associated with simple backache arthritis strains bruises sprains.

Warnings: For external use only. Do not use - on wounds or damaged skin, broken (open) or irritated skin - with a heating pad. When using this product - use only as directed - avoid contact with eyes or mucous membranes - do not bandage tightly. Stop use and ask a doctor if condition worsen's or symptoms persist for more than 7 days - symptoms clear up and occur again within a few days - excessive skin irritation occurs. If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children to avoid accidental ingestion. If swallowed, get medical help or contact a Poison Control Center immediately.

Directions • use only as directed. Adults and children 12 years of age and older: Apply to affected area not more than 3 to 4 times daily. Children under 12 years of age: Consult a doctor. Questions or comments? Call to

INFORMACIÓN DEL MEDICAMENTO

Ingredientes activos Propósito
Mentol - 10%, Silicato de Metilo - 15% Analgésico Tópico

Usos: Para el alivio temporal de dolores menores y dolores musculares y de articulaciones asociados com • dolor de espalda simple • artiritis • torceduras • moretones • esquinces

Advertencias: Para uso externo solamente. No se utilice • sobre heridas, piel dañada, rasgada o irritada • en conjunto con una almohadila térmica. Cuando use este producto • Utilice únicamente según las indicaciones • evitar el contacto con los ojos y membranas mucosa • no aplicar un vendaje muy ajustado Suspenda su uso y consulte a su médico si • la condición empeora o el dolor persiste por más de 7 días • los síntomas desaparecen y vuelven a aparecer dentro de algunos días • se presenta irritación excesiva de la piel. En caso de embarazo o lactancia, consulte a un profesional médico antes de su uso. Manténgase fuera del alcance de los niños. En caso de ingerirlo, obtenga asistencia médica o contacte a un Centro de Control de Envenenamiento de manera inmediata.

Indicaciones • utilizar según indicaciones. Adultos y niños mayores de 12 años: Aplicar en el área afectada no mas de 3 a 4 veces por dia. Niños menores de 12 años: Consulte a su médico.

¿Preguntas o comentarios? Llame lada sin costo 1 877 99 GENOM (43666)

For further details look at the box / Para más detailes ver la caja
See tube crimp for Lot No./Lote No. and Exp.

Made in the USA
Distributed by / Distribution por: Genomma Lab USA, Inc., Houston, TX 77027

RAY DOL

pain relieving cream cream

Product Information

Route of Administration

Product TypeHUMAN OTC DRUGItem Code (Source)NDC:50066-057

TOPICAL

Active Ingredient/Active Majety

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	100 mg in 1 g	
METHYL SALICYLATE (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII:O414PZ4LPZ)	METHYL SALICYLATE	150 mg in 1 g	

Inactive Ingredients	
Ingredient Name	Strength
CARBOMER 940 (UNII: 4Q93RCW27E)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
TRO LAMINE (UNII: 903K93S3TK)	
WATER (UNII: 059QF0KO0R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

OLETH-3 PHO SPHATE (UNII: 8Q0Z18J1VL)

F	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50066-057-73	1 in 1 CARTON		
1		85 g in 1 TUBE		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part348	12/15/2013		

Labeler - Genomma Lab USA Inc. (832323534)

Registrant - Faria LLC dba Sheffield Pharmaceuticals (151177797)

Establishment			
Name	Address	ID/FEI	Business Operations
Faria LLC dba Sheffield Pharmaceuticals		151177797	manufacture (50066-057)

Revised: 1/2014 Genomma Lab USA Inc.