

**ASPIRIN - aspirin tablet, extended release**  
**Gemini Pharmaceuticals, Inc. dba Plus Pharmaceuticals**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID\*)

\* nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- temporarily relieves minor aches and pains
- for other uses, see your doctor, but do not use for more than 10 days without consulting your doctor because serious side effects may occur

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu- like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

- you have asthma
- you are taking a diuretic
- you have not been drinking fluids
- you have lost a lot of fluid due to vomiting or diarrhea

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug
- under a doctor's care for any serious condition

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleedin

feel faint

vomit blood

have bloody or black stools

- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions:**

- Drink a full glass of water with each dose
- **adults and children 12 years and over:** take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- **children under 12 years:** consult a doctor

**Other information**

- **Do not use if imprinted safety seal under cap is broken or missing.**
- store at 25<sup>0</sup>C (77<sup>0</sup> F) excursions permitted between 15<sup>0</sup>-30<sup>0</sup>C (59<sup>0</sup>-86<sup>0</sup> F)

**Inactive ingredients:**

Anhydrous Lactose, Carnauba Wax, Colloidal Silicon Dioxide, Croscarmellose Sodium, D and C Yellow #10 Aluminum Lake, Iron Oxide Ochre, Methacrylic Acid Copolymer, Microcrystalline Cellulose, Polysorbate 80, Simethicone, Sodium Hydroxide, Sodium Lauryl Sulfate, Talc, Titanium

Dioxide, Triethyl Citrate.

## Questions?

If you have any questions or comments, or to report an adverse event, please contact (800) 795-9775.

## Principal Display Panel

PlusPHARMA

ASPIRIN

81 mg

NSAID

PAIN RELIEVER

LOW DOSE ASPIRIN REGIMEN

- Safety Coated-Enteric Coated
- Delayed Release Aspirin Tablets
- Small, Easy-to-Swallow Tablets

\*Compare to the active ingredient in Bayer Low Dose

\*This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Bayer Low Dose.

120 TABLETS

- 81 mg each

PlusPHARMA™

# ASPIRIN

NSAID 81 mg

**PAIN RELIEVER  
LOW DOSE ASPIRIN REGIMEN**

- Safety Coated - Enteric Coated
- Delayed Release Aspirin Tablets
- Small, Easy-to-Swallow Tablets

\*This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Bayer® Low Dose.

\*Compare to the Active Ingredient in Bayer® Low Dose

120 TABLETS • 81 mg each

## Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 81 mg (NSAID*)	Pain reliever
*nonsteroidal anti-inflammatory drug	

**Uses** • temporarily relieves minor aches and pains • for other uses, see your doctor, but do not use for more than 10 days without consulting your doctor because serious side effects may occur

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: • hives • facial swelling • asthma (wheezing) • shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug

• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • take more or for a longer time than directed • are age 60 or older • have 3 or more alcoholic drinks every day while using this product

**Drug Facts** (continued on back)

**Do not use if imprinted safety seal under cap is broken or missing**



Rev 12/12

0 0 0 0 4 1 1 0 0 0

Lot#

Exp. Date

PEEL  
HERE



### **Drug Facts** (continued)

**Do not use** • if you are allergic to aspirin or any other pain reliever/fever reducer

**Ask a doctor before use if** • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you have asthma • you are taking a diuretic • you have not been drinking fluids • you have lost a lot of fluid due to vomiting or diarrhea

**Ask a doctor or pharmacist before use if you are** • taking a prescription drug for diabetes, gout, or arthritis • taking any other drug • under a doctor's care for any serious condition

#### **Stop use and ask a doctor if**

• you experience any of the following signs of stomach bleeding:  
• feel faint • vomit blood • have bloody or black stools  
• have stomach pain that does not get better  
• pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present in the painful area • any new symptoms appear  
• ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions:** • Drink a full glass of water with each dose  
• **adults and children 12 years and over:** take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor  
• **children under 12 years:** consult a doctor

**Other information** • store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients:** Anhydrous Lactose, Carnauba Wax, Colloidal Silicon Dioxide, Croscarmellose Sodium, D&C Yellow #10 Aluminum Lake, Iron Oxide Ochre, Methacrylic Acid Copolymer, Microcrystalline Cellulose, Polysorbate 80, Simethicone, Sodium Hydroxide, Sodium Lauryl Sulfate, Talc, Titanium Dioxide, Triethyl Citrate.

**Questions?** If you have any questions or comments, or to report an adverse event, please contact (800) 795-9775.

**Distributed By:** Plus Pharma, Commack, NY 11725

## ASPIRIN

aspirin tablet, extended release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG LABEL	<b>Item Code (Source)</b>	NDC:51645-717
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (ASPIRIN)	ASPIRIN	81 mg

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE	
CARNAUBA WAX	
SILICON DIOXIDE	
CROSCARMELLOSE SODIUM	
D&C YELLOW NO. 10	
CELLULOSE, MICROCRYSTALLINE	
POLYSORBATE 80	
SODIUM HYDROXIDE	
SODIUM LAURYL SULFATE	
TALC	
TITANIUM DIOXIDE	
TRIETHYL CITRATE	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND (YELLOW)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	RAISED;HEART
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51645-717-08	120 in 1 BOTTLE, PLASTIC		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	01/10/2013	

**Labeler** - Gemini Pharmaceuticals, Inc. dba Plus Pharmaceuticals (055942270)

**Registrant** - Gemini Pharmaceuticals, Inc. dba Plus Pharmaceuticals (055942270)