

**UP AND UP ANTI DIARRHEAL ANTI GAS- loperamide hydrochloride, simethicone tablet**  
**Target Corporation**

-----  
**Target Corporation anti-diarrheal/anti-gas Drug Facts**

**Active ingredients (in each caplet)**

Loperamide HCl 2 mg

Simethicone 125 mg

**Purposes**

Anti-diarrheal

Anti-gas

**Uses**

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

- if you have bloody or black stool
- if you have difficulty swallowing

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- **do not use if printed blister unit is broken o**

**Inactive ingredients**

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

**Questions?**

**Call 1-888-547-7400**

**Package/Label Principal Display Panel**

Compare to active ingredients in Imodium® Multi-Symptom Relief

loperamide hydrochloride and simethicone tablets, 2 mg/125 mg

anti-diarrheal/anti-gas

multi-symptom relief caplet

relieves symptoms of diarrhea *plus:*

cramps and pressure, bloating, gas

ACTUAL SIZE

24 CAPLETS\*

24 CAPLETS\*

(\*CAPSULE-SHAPED TABLETS)



OR 162 LW C1

**Drug Facts**

**Active ingredients (in each caplet) Purposes**  
 Loperamide HCl 2 mg ..... Anti-diarrheal  
 Simethicone 125 mg ..... Anti-gas

**Uses** relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

**Warnings**  
**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl  
**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**  
 ■ if you have bloody or black stool  
 ■ if you have difficulty swallowing

**Ask a doctor before use if you have**  
 ■ fever ■ mucus in the stool ■ a history of liver disease  
 ■ a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**  
 ■ symptoms get worse  
 ■ diarrhea lasts for more than 2 days  
 ■ you get abdominal swelling or bulging. These may be signs of a serious condition  
 ■ you have difficulty swallowing the caplet

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**  
 ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea  
 ■ take only on an empty stomach (1 hour before or 2 hours after a meal)  
 ■ take with a full (8 oz.) glass of water  
 ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
---------------------------------------	---

children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
---------------------------------	--

children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
--------------------------------	--

children 2-5 years (34 to 47 lbs)	ask a doctor
-----------------------------------	--------------

children under 2 years (up to 33 lbs)	do not use
---------------------------------------	------------

**Other information**  
 ■ each caplet contains calcium 165 mg, sodium 3 mg  
 ■ store between 20°-25°C (68°-77°F). Protect from light  
 ■ do not use if printed blister unit is broken or torn

loperamide hydrochloride and simethicone tablets, 2 mg/125 mg  
**anti-diarrheal/anti-gas**



**Drug Facts (continued)**

**Inactive ingredients** acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, methylcellulose, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

**Questions?** Call 1-888-547-7400

**GLUTEN FREE**  
 \*\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® Multi-Symptom Relief.

Distributed by Target Corporation  
 Minneapolis, MN 55403  
 TM & ©2021 Target Brands, Inc.

245 05 1285 R00  
 C-001472-01-078



3 70030 14716 8

**UP AND UP ANTI DIARRHEAL ANTI GAS**

loperamide hydrochloride, simethicone tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-646
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg

**Inactive Ingredients**

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)

MALTODEXTRIN (UNII: 7CVR7L4A2D)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

STEARIC ACID (UNII: 4ELV7Z65AP)

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	L0R1;2;125
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-646-89	18 in 1 CARTON	03/14/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11673-646-62	24 in 1 CARTON	01/06/2021	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209837	03/14/2019	

**Labeler** - Target Corporation (006961700)

Revised: 1/2021

Target Corporation