

GOOD NEIGHBOR PHARMACY LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride liquid
Amerisource Bergen

Amerisource Bergen Loperamide Hydrochloride Oral Suspension Drug Facts

Active ingredient (in each 7.5 mL)

Loperamide HCl 1 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are

taking antibiotics

When using this product

- tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.

These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- only use attached measuring cup to dose product

adults and children 12 years and over	30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours
children 9-11 years (60-95 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours
children 6-8 years (48-59 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 30 mL (6 tsp) in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information

- each 30 mL (6 tsp) contains: sodium 15 mg
- store between 20-25°C (68-77°F)
- see side panel for lot number and expiration date

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Imodium® A-D active ingredient

Loperamide Hydrochloride Oral Suspension

anti-diarrheal

Controls the Symptoms of Diarrhea

mint flavor

1 mg loperamide hydrochloride per 7.5 mL

4 fl oz (120 mL)

Drug Facts

Active ingredient Purpose
(in each 7.5 mL)

Loperamide HCl
1 mg.....Anti-diarrheal

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have bloody or black stool

PEEL BACK HERE



GOOD NEIGHBOR PHARMACY® NDC 46122-111-26

Compare to Imodium® A-D active ingredient*

Loperamide Hydrochloride Oral Suspension
anti-diarrheal

Controls the Symptoms of Diarrhea

mint flavor

1 mg loperamide hydrochloride per 7.5 mL


4 fl oz (120 mL)

Do not use if printed plastic neckband is broken or missing.

GLUTEN FREE

Distributed By
AmerisourceBergen
1300 Morris Drive
Chestertown, PA 19087
Visit us at
www.goodneighborpharmacy.com

ABC# 142-796



0 87701 41126 7

*Good Neighbor Pharmacy® Loperamide Hydrochloride Oral Suspension is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Imodium® A-D.

: 64526 29 F2

Drug Facts (continued)

Ask a doctor before use if you have ■ fever
■ mucus in the stool ■ a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product ■ tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

■ symptoms get worse
■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bulging.
These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.


Keep out of reach of children. In case of overdose, get medical help or contact a Poison

Drug Facts (continued)

Control Center right away. (1-800-222-1222)

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
■ find right dose on chart. If possible, use weight to dose; otherwise use age. ■ shake well before using ■ only use attached measuring cup to dose product

adults and children 12 years and over	30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours
children 9-11 years (60-95 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours



GOOD NEIGHBOR PHARMACY LOPERAMIDE HYDROCHLORIDE			
loperamide hcl liquid			
Product Information			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:46 122-111
Route of Administration	ORAL	DEA Schedule	
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
LOPERAMIDE HYDROCHLORIDE (LOPERAMIDE)	LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL	
Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID			
CARBOXYMETHYLCELLULOSE SODIUM			
D&C YELLOW NO. 10			
FD&C BLUE NO. 1			
GLYCERIN			
CELLULOSE, MICROCRYSTALLINE			
PROPYLENE GLYCOL			
WATER			
SODIUM BENZOATE			
SUCRALOSE			
TITANIUM DIOXIDE			
XANTHAN GUM			
Product Characteristics			

Color	GREEN (opaque, viscous)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46 122-111-26	120 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA09 1292	02/17/20 12	

Labeler - Amerisource Bergen (007914906)

Revised: 4/2014

Amerisource Bergen