MAGNESIUM CITRATE- freskaro magnesium citrate oral solution - cherry liquid Pharma Nobis, LLC

Freskaro Magnesium Citrate Oral Solution - Cherry

Active Ingredient

Magnesium citrate 1.745g per fl oz

Purpose

Saline Laxtive

Use

for relief of occasional constipation (irregularity). Generally produces bowel movement in 1/2 to 6 hours

Warnings

kidney disease
a magnesium restricted diet
abdorominal pain, nausea, or vomiting
noticed a sudden change in bowel habits that persists over a period of 2 weeks
already used a laxative for a period longer than 1 week

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking any other drug.

Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

STOP USE AND ASK A DOCTOR IF

you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

IF PREGNANT OR BREAST-FEEDING

ask health professional before use.

KEEP OUT OF REACH OF CHILDREN.

In case of overdose, get medical help or contact a Poison Control Center right away.

Other Information

Store at temperatures between 46° and 86°F (8° and 30° C)

Directions

shake well before using drink a full glass (8 ounces) of liquid with each dose may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over - 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age - 2 to 3 fl oz in 24 hours maximum 3 fl oz in 24 hours children under 2 years of age - ask a doctor

Other Information

each fl oz contains: magnesium 290 mg

each fl oz contains: sodium 1 mg

Inactive Ingredients

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions?

1-833-551-0932

Drug Facts

Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 gSaline laxative

Uses: For relief of occasional constipation

(irregularity).

Generally produces bowel movement in ½ to 6 hours

Warnings:

Ask a doctor before use if you have

Kidney disease a magnesium restricted diet
 Abdominal pain, nausea, or vomiting
 Noticed a sudden change in bowel habits that persists over a period of 2 weeks

Already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are

■ Taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Do not use if tamper evident twist-off cap is missing, broken, or separated from neckring



Magnesium Citrate

Saline Laxative Oral Solution

- Sugar Free
- For relief of occasional constipation
- For use in bowel cleanings systems for surgery or exam preparation

Alcohol content 0.005% (From flavoring)

Cherry Flavor

10 fl. oz (296ml)

Drug Facts (continued)

Store at temperatures 46° and 86°F (8° and 30°C)

Directions: ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses adults and children 12

6.5 to 10 fl oz maximum 10 fl oz in 24 hours years of age and over children 6 to under 12 3 to 7 fl oz maximum 7 fl oz in 24 hours years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children 2 to under 6 years of age children under 2 years ask a doctor of age

Other Information:

each fl oz contains: magnesium 290 mg each fl oz contains: sodium 1 mg

Inactive ingredients: benzoic acid, citric acid disodium EDTA, flavor, sucralose, water

Questions? 1-833-551-0932

Distributed by: Pharma Nobis 7400 Alumax Dr. Texarkana, TX 75501



MAGNESIUM CITRATE

freskaro magnesium citrate oral solution - cherry liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:82645-312

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM CITRATE (UNII: RHO2601T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CITRATE	1.745 g in 29.6 mL

Inactive Ingredients Ingredient Name Strength BENZOIC ACID (UNII: 85KN0B0MIM) WATER (UNII: 059QF0KO0R) CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) **SUCRALOSE** (UNII: 96K6UQ3ZD4) **EDETATE DISODIUM ANHYDROUS (UNII: 8NLQ36F6MM)**

Packaging

#	Item Code		Package Description	Date	Date	
1	NDC:82645- 312-10		in 1 BOTTLE, PLASTIC; Type 0: Not a ation Product	12/27/2022		
M	larketing	Info	rmation			
M	larketing Marketing Category		rmation Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
	Marketing	•	Application Number or Monograph Citation			

Labeler - Pharma Nobis, LLC (118564114)

Registrant - Pharma Nobis, LLC (118856114)

Establishment						
Name	Address	,	Business Operations			
Pharma Nobis, LLC		118564114	label(82645-312), manufacture(82645-312), analysis(82645-312), pack(82645-312)			

Revised: 12/2023 Pharma Nobis, LLC