

TYLENOL COLD PLUS FLU SEVERE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, film coated

Select Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Tylenol® Cold Plus Flu Severe

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets every 4 hours▪ swallow whole; do not crush, chew or dissolve▪ do not take more than 10 caplets in 24 hours
children under 12 years	ask a doctor

Other information

- **each caplet contains:** sodium 3 mg
- store between 20-25°C (68-77°F)
- **do not use if pouch is torn or damaged**

Inactive ingredients

carnauba wax, croscarmellose sodium, D&C yellow no. 10 aluminum lake, flavor, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, sucralose, titanium dioxide

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by:

**JOHNSON & JOHNSON
CONSUMER INC.**

McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA

Product Repackaged and Distributed
with permission of manufacturer by:

Select Corporation, Carrollton, TX 75007

PRINCIPAL DISPLAY PANEL - 2 Caplet Pouch Single-Pack

**TYLENOL®
FOR ADULTS**

COLD + FLU SEVERE

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl, Guaifenesin
Pain Reliever-Fever Reducer, Cough Suppressant, Nasal Decongestant, Expectorant


2 Caplets
Single-Pack



Made in Italy
Distributed by:
**JOHNSON & JOHNSON
CONSUMER INC.**
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2018
Visit us at www.tylenol.com
or call toll-free 1-877-TYLENOL
(1-877-895-3665)



1712944083



Open Flap for **DRUG FACTS** →
Read all warnings and directions before use



Single - Pack: 2 Caplets

Product Repackaged and Distributed
with permission of manufacturer by:
Select Corporation, Carrollton, TX 75007

Drug Facts	
Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant
Uses	
<ul style="list-style-type: none"> for the temporary relief of the following cold/flu symptoms: <ul style="list-style-type: none"> minor aches and pains headache sore throat nasal congestion cough helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive temporarily reduces fever 	
Warnings	
<p>Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin reddening hives rash <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting.</p>	

Drug Facts (continued)
<p>consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease heart disease high blood pressure thyroid disease diabetes trouble urinating due to an enlarged prostate gland persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema cough that occurs with too much phlegm (mucus) <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</p> <p>When using this product do not exceed recommended dose</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> nervousness, dizziness, or sleeplessness occur pain, nasal congestion or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p>

Drug Facts (continued)				
<p>Keep out of reach of children.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed (see overdose warning) <table border="1"> <tr> <td>adults and children 12 years and over</td> <td> <ul style="list-style-type: none"> take 2 caplets every 4 hours swallow whole; do not crush, chew or dissolve do not take more than 10 caplets in 24 hours </td> </tr> <tr> <td>children under 12 years</td> <td>ask a doctor</td> </tr> </table> <p>Other information</p> <ul style="list-style-type: none"> each caplet contains: sodium 3 mg store between 20-25°C (68-77°F) do not use if pouch is torn or damaged <p>Inactive ingredients carnauba wax, croscarmellose sodium, D&C yellow no. 10 aluminum lake, flavor, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, sucralose, titanium dioxide</p> <p>Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)</p>	adults and children 12 years and over	<ul style="list-style-type: none"> take 2 caplets every 4 hours swallow whole; do not crush, chew or dissolve do not take more than 10 caplets in 24 hours 	children under 12 years	ask a doctor
adults and children 12 years and over	<ul style="list-style-type: none"> take 2 caplets every 4 hours swallow whole; do not crush, chew or dissolve do not take more than 10 caplets in 24 hours 			
children under 12 years	ask a doctor			

1712944083

TYLENOL COLD PLUS FLU SEVERE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:52904-944(NDC:50580-402)
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	acetaminophen (UNII: 362O9ITL9D) (acetaminophen - UNII:362O9ITL9D)	acetaminophen	325 mg
	dextromethorphan hydrobromide (UNII: 9D2RTI9KYH) (dextromethorphan - UNII:7355X3ROTS)	dextromethorphan hydrobromide	10 mg
	guaifenesin (UNII: 495W7451VQ) (guaifenesin - UNII:495W7451VQ)	guaifenesin	200 mg
	phenylephrine hydrochloride (UNII: 04JA59TNSJ) (phenylephrine - UNII:1WS297W6MV)	phenylephrine hydrochloride	5 mg

Inactive Ingredients		Strength
	Ingredient Name	
	carnauba wax (UNII: R12CBM0EIZ)	
	croscarmellose sodium (UNII: M28OL1HH48)	
	D&C yellow no. 10 aluminum lake (UNII: CQ3XH3DET6)	
	hydroxypropyl cellulose, unspecified (UNII: 9XZ8H6N6OH)	
	hypromellose, unspecified (UNII: 3NXW29V3WO)	
	magnesium stearate (UNII: 70097M6I30)	
	microcrystalline cellulose (UNII: OP1R32D61U)	
	polyethylene glycol, unspecified (UNII: 3WJQ0SDW1A)	
	sucralose (UNII: 96K6UQ3ZD4)	
	titanium dioxide (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	YELLOW	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	TYLENOL;SEVERE
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52904-944-04	1 in 1 BLISTER PACK	09/01/2011	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:52904-944-05	2 in 1 BLISTER PACK	09/01/2011	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:52904-944-25	25 in 1 CARTON	09/01/2011	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:52904-944-	50 in 1 CARTON	09/01/2011	

4	50	50 III 1 CARTON	09/01/2011	
4		2 in 1 POUCH; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341		09/01/2011	

Labeler - Select Corporation (053805599)

Revised: 3/2022

Select Corporation