

**GERI-KOT- sennosides tablet**  
**Geri-Care Pharmaceutical Corp**

-----  
**GC 451B 415**

**Active ingredient (in each tablet)**

Sennosides 8.6 mg

**Purpose**

Laxative

**Uses**

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

**Warnings**

**Do not use**

for more than 1 week unless directed by a doctor.

**Ask a doctor before use if you**

- have abdominal pain, nausea or vomiting
- have noticed a sudden change in bowel habits that lasts over 2 weeks

**Stop use and ask a doctor if**

you have rectal bleeding  
or fail to have a bowel movement after use of a laxative.  
These may indicate a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not exceed 8 tablets in 24 hours

| Age   | Starting Dose  | Maximum Dose  |
|---|--|---|
| adults and children<br>12 years of age and<br>older | 2 tablets once a day preferably at<br>bedtime; increase as needed, or as<br>directed by a doctor | 4 tablets in the<br>morning and 4 tablets<br>at bedtime |
| children under 12<br>years                          | ask a doctor   |   |

- **each tablet contains:** calcium 30 mg
- Tamper Evident: Do not use if imprinted seal under cap is missing or broken
- store at 20°C-25°C (68°F-77°F)

croscarmellose sodium, dicalcium phosphate, hypromellose, magnesium stearate, microcrystalline cellulose, silica, talc, triacetin

GERI CARE®

NDC 57896-415-01

**GERI-KOT**

STANDARDIZED SENNA CONCENTRATE

NATURAL VEGETABLE  
LAXATIVE

Compare to active  
ingredient in **SENOKOT**®\*

**100 Tablets 8.6 mg each**

### Drug Facts

#### Active ingredient (in each tablet)

Sennosides 8.6 mg.....Laxative

#### Uses • relieves occasional constipation (irregularity)

• generally produces a bowel movement in 6 to 12 hours

#### Warnings

**Do not use** for more than one week unless directed by a doctor.

**Ask a doctor before use** if you have

- stomach pain, nausea or vomiting
- noticed a change in bowel habits that lasts over two weeks

**Stop use and ask a doctor** if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions • do not exceed 8 tablets in 24 hours

| Age   | Starting Dose  | Maximum Dose                                      |
|---|--|---|
| Adults and children 12 years of age and older | 2 tablets once a day preferably at bedtime; increase as needed, or as directed by a doctor | 4 tablets in the morning and 4 tablets at bedtime |
| Children under 12 years                       | Ask a doctor   |   |

#### Other information • each tablet contains:

- calcium 30 mg • **Tamper Evident:** Do not use if imprinted seal under cap is missing or broken
- store at 20°C-25°C (68°F-77°F)

**Inactive ingredients:** croscarmellose sodium, dicalcium phosphate, hypromellose, magnesium stearate, microcrystalline cellulose, silica, talc, triacetin.

#### Questions or comments? Call 1-800-540-3765

\*This product is not manufactured or distributed by owner of the registered trademark SENOKOT®.

Dist. by: Geri-Care Pharmaceutical Corp.  
1295 Towbin Ave., Lakewood, NJ 08701

Product of India

REV 11228

## sennosides tablet

|                         |                |                    |               |
|-------------------------|----------------|--------------------|---------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:57896-415 |
| Route of Administration | ORAL           |                    |               |

| Ingredient Name   | Basis of Strength | Strength |
|---|-------------------|----------|
| <b>SENNOSIDES</b> (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX) | SENNOSIDES        | 8.6 mg   |

| Ingredient Name                                | Strength |
|--|----------|
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) |          |

**CROSCARMELLOSE SODIUM** (UNII: M28OL1HH48)

**DIBASIC CALCIUM PHOSPHATE DIHYDRATE** (UNII: O7TSZ97GEP)

**SILICON DIOXIDE** (UNII: ETJ7Z6XBU4)

**MAGNESIUM STEARATE** (UNII: 70097M6I30)

**TALC** (UNII: 7SEV7J4R1U)

**TRIACETIN** (UNII: XHX3C3X673)

**HYPROMELLOSES** (UNII: 3NXW29V3WO)

### Product Characteristics

|                 |       |                     |          |
|-----------------|-------|---------------------|----------|
| <b>Color</b>    | brown | <b>Score</b>        | no score |
| <b>Shape</b>    | ROUND | <b>Size</b>         | 9mm      |
| <b>Flavor</b>   |       | <b>Imprint Code</b> | PS23     |
| <b>Contains</b> |       |                     |          |

### Packaging

| # | Item Code        | Package Description                                 | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:57896-415-01 | 100 in 1 BOTTLE; Type 0: Not a Combination Product  | 12/01/2022           |                    |
| 2 | NDC:57896-415-10 | 1000 in 1 BOTTLE; Type 0: Not a Combination Product | 12/01/2022           |                    |
| 3 | NDC:57896-415-20 | 200 in 1 BOTTLE; Type 0: Not a Combination Product  | 12/01/2022           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | 505G(a)(3)                               | 12/01/2022           |                    |

**Labeler** - Geri-Care Pharmaceutical Corp (611196254)

**Registrant** - Geri-Care Pharmaceutical Corp (611196254)

Revised: 12/2023

Geri-Care Pharmaceutical Corp