# NASAL DECONGESTANT- pseudoephedrine hydrochloride tablet L.N.K. International, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### **Quality Plus 44-112**

#### Active ingredient (in each tablet)

Pseudoephedrine HCl 30 mg

#### Purpose

Nasal decongestant

#### Uses

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves sinus congestion and pressure

#### **Warnings**

#### Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains and MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- diabetes
- high blood pressure
- heart disease
- thyroid disease
- trouble urinating due to an enlarged prostate gland

#### When using this product

do not exceed recommended dose.

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with fever

#### If pregnant or breast-feeding,

ask a health professional before use.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

adults and children 12 years and older	take 2 tablets every 4 to 6 hours; do not take more than 8 tablets in 24 hours
children ages 6 to 12 years	take 1 tablet every 4 to 6 hours; do not take more than 4 tablets in 24 hours
children under 6 years	do not use this product in children under 6 years of age

#### Other information

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

• see end flap for expiration date and lot number

#### **Inactive ingredients**

croscarmellose sodium, dicalcium phosphate, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, silica gel, titanium dioxide, triacetin

Questions or comments? 1-800-426-9391

#### **Principal Display Panel**

NDC 50844-112-08

\*Compare to the active ingredient in Sudafed® Congestion

Non-Drowsy / Maximum Strength Nasal Decongestant Pseudoephedrine HCl 30 mg

#### **Relieves:**

- Nasal & Sinus Congestion
- Sinus Pressure

24 Tablets

#### **QUALITY PLUS**

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Sudafed® Congestion.

50844 REV0712L11208

# TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Distributed by LNK INTERNATIONAL, INC.

60 Arkay Drive, Hauppauge NY 11788

**USA** 

#### Non-Drowsy / Maximum Strength

# Nasal Decongestant

Pseudoephedrine HCl 30 mg

NDC 50844-112-08

208

\*Compare to the active ingredient in Sudafed® Congestion

Non-Drowsy / Maximum Strength

# Nasal Decongestant

Pseudoephedrine HCl 30 mg

#### Relieves:

Nasal & Sinus Congestion

- Sinus Pressure



#### 24 Tablets

60 Arkay Drive, Hauppauge, NY 11788 Distributed by LNK INTERNATIONAL, INC.

REV0712L11208 trademark Sudafed® Congestion. McNeil Consumer Healthcare, owner of the registered This product is not manufactured or distributed by



B-1603-112-08 REV0712L11208

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Questions or comments? 1-800-426-9391

polydextrose, polyethylene glycol, silica gel, titanium dioxide, triacetin lake, hypromellose, magnesium stearate, microcrystalline cellulose, Inactive ingredients croscsmellose sodium, dicalcium phosphate, FD&C yellow ∌6 aluminum lake, FD&C yellow ∌6 aluminum

> ■ see end flap for expiration date and lot number ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Uther information

сhildren under 6 years	do not use this product in children under 6 years of age
children ages 6 to 12 years	take 1 tablet every 4 to 6 hours; do not take more than 4 tablets in 24 hours
adults and children 12 years and older	take 2 tablets every 4 to 6 hours; do not take more than 8 tablets in 24 hours

contact a Poison Control Center right away. Keep out of reach of children. In case of overdose, get medical help or

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

- symptoms do not improve within 7 days or occur with fever ■ nervousness, dizziness, or sleeplessness occur
  - Stop use and ask a doctor if

When using this product do not exceed recommended dose.

- trouble uninating due to an enlarged prostate gland ■ high blood pressure ■ thyroid disease
  - heart disease Ask a doctor before use if you have

MAOI, ask a doctor or pharmacist before taking this product. MAOI drug. If you do not know if your prescription drug contains an conditions, or Parkinson's disease), or for 2 weeks after stopping the inhibitor (MAOI) (certain drugs for depression, psychiatric or emotion Do not use if you are now taking a prescription monoamine oxidase

> ■ temporarily relieves sinus congestion and pressure fever or other upper respiratory allergies

> > Active ingredient (in each tablet)

■ temporarily relieves nasal congestion due to the common cold, hay

Pseudoephedrine HCI 30 mg.

Nasal decongestant Purpose

> **РВОDUCT ІИГОВМАТІОИ** RETAIN CARTON FOR COMPLETE

Drug Facts

8087A

LEBG3130

**Qualty Plus 44-112** 

### pseudoephedrine hcl tablet

Product Information			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:50844-112
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
PSEUDO EPHEDR	INE HYDRO CHLO RIDE (PSEUDO EPHEDRINE)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

Inactive Ingredients			
Ingredient Name	Strength		
CROSCARMELLOSE SODIUM			
CALCIUM PHO SPHATE, DIBASIC, ANHYDRO US			
FD&C YELLOW NO. 6			
HYPROMELLOSES			
MAGNESIUM STEARATE			
CELLULOSE, MICRO CRYSTALLINE			
POLYDEXTROSE			
POLYETHYLENE GLYCOLS			
TITANIUM DIO XIDE			
TRIACETIN			

Product Characteristics				
Color	RED	Score	no score	
<b>Shape</b> R	ROUND	Size	8 mm	
Flavor		Imprint Code	44;112	
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:50844-112-22	4 in 1 CARTON			
1		12 in 1 BLISTER PACK; Combination Product Type = C112160			
2	NDC:50844-112- 08	2 in 1 CARTON			
2		12 in 1 BLISTER PACK; Combination Product Type = C112160			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	08/25/1981	

## Labeler - L.N.K. International, Inc. (038154464)

Revised: 11/2014 L.N.K. International, Inc.