

EQUATE IBUPROFEN - ibuprofen capsule, liquid filled
Walmart stores Inc

ACTIVE INGREDIENT(S)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*
(Present as the free acid and potassium salt)

***nonsteroidal anti-inflammatory drug**

PURPOSE

Pain reliever/Fever reducer

USE(S)

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

WARNINGS

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

DO NOT USE

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

ASK A DOCTOR BEFORE USE IF

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

ASK A DOCTOR OR PHARMACIST BEFORE USE IF

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

WHEN USING THIS PRODUCT

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

STOP USE AND ASK DOCTOR IF

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

OTHER REQUIRED WARNINGS

- each capsule contains: potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

PREGNANCY/BREASTFEEDING

Ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

DIRECTIONS

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

INACTIVE INGREDIENT(S)

Ammonium hydroxide, FD&C green no.3, Gelatin, Iron oxide black (ferrosoferric oxide), Medium chain triglycerides, Polyethylene glycol, Propylene glycol, Potassium hydroxide, Purified water, Shellac, Sorbitol, Sorbitan monooleate

PRINCIPAL DISPLAY PANEL

NDC 49035-133-12

equate™

Ibuprofen

Capsules, 200mg

Pain Reliever/ Fever Reducer

(NSAID)

200mg, 20 SOFTGELS

Liquid-Filled Capsules

equate™ NDC 49035-133-12
Compare to Advil® Liquid-Gels® Active Ingredient

Ibuprofen
Capsules, 200 mg
Pain Reliever/
Fever Reducer
(NSAID)

200 mg **20** SOFTGELS
Actual Size
Liquid-Filled Capsules

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing

Drug Facts	Purpose
Active ingredient (in each capsule) Solvated ibuprofen equal to 200 mg ibuprofen (NSAID)*	Pain reliever/fever reducer (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug
Uses	temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever
Warnings	KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued under label)

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

Excluded by MedWatch, Inc., Rockville, MD 20718
*This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil Liquid-Gels® (45888)

Lot No.:
PBF
Exp. Date:

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Directions ■ do not take more than directed ■ the smallest effective dose should be used

■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor

Questions or comments?

1-888-287-9115

Drug Facts (continued)

■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used

Drug Facts (continued)

■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor

Other information

■ each capsule contains: potassium 20 mg ■ store at 20°-25°C (68°-77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, propylene glycol, potassium hydroxide, purified water, shellac, sorbitan monoolate, sorbitol sorbitan

Questions or comments? 1-888-287-1915

LOT:
EXP:



458880

Manufactured by: New-Mark Stores, Inc., Berwyn, PA 17716
*This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil.

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP THIS BOX FOR IMPORTANT INFORMATION.

NDC 49035-133-12

Actual Size

200 mg
20 SOFTGELS[®]
Liquid-Filled Capsules

Pain Reliever/Fever Reducer (NSAID)

Capsules, 200 mg

Ibuprofen

equate™

Drug Facts

Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing

Active ingredient (in each capsule)

Purpose

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*. Pain reliever/Fever reducer (present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Uses

■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, liver cirrhosis or asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs



NDC 49035-133-14

equate™

Ibuprofen
 Capsules, 200mg
 Pain Reliever/ Fever
 Reducer (NSAID)
 Liquid-Filled Capsules
 200mg, 40 SOFTGELS

equate™ NDC 45085-185H4

Compare to Advil® Liquid-Gel® Active Ingredient**

Ibuprofen Capsules, 200 mg
 Pain Reliever/ Fever Reducer (NSAID)
 *Liquid-Filled Capsules

200 mg **40**
 SOFTGELS

Actual Size

Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing

Active ingredient (in each capsule)
 Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*Pain reliever/fever reducer (present as the free acid and potassium salt)
 *nonsteroidal anti-inflammatory drug

Purpose
 ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ toothache ■ backache
 ■ menstrual cramps ■ the common cold
 ■ muscular aches ■ minor pain of arthritis
 ■ temporarily reduces fever

Uses
 ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ toothache ■ backache
 ■ menstrual cramps ■ the common cold
 ■ muscular aches ■ minor pain of arthritis
 ■ temporarily reduces fever

Warnings
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
 Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
 Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing)
 ■ shock ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued under label)

Satisfaction guaranteed – For questions or comments please call 1-888-287-1915.

Distributed by: Wal-Mart Stores, Inc., Bentonville, AR 72716
 **This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil®.

Lot No.:
 Exp Date:

PEEL HERE

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Directions
 ■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
 ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used
 ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor
 ■ children under 12 years: ask a doctor

Questions or comments? 1-888-287-1915

Drug Facts (continued)

Directions ■ do not take more than directed ■ the smallest effective dose should be used

adults and children 12 years and over	■ take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor
children under 12 years	■ ask a doctor

Other information
 ■ each capsule contains: potassium 20 mg ■ store at 20°-25°C (68°-77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, propylene glycol, potassium hydroxide, purified water, shellac, sorbitan monoleate, sorbitol sorbitan

Questions or comments? 1-888-267-1915

 Satisfaction guaranteed - Or we'll replace it or give you your money back. No questions asked. For comments or to report an undesired reaction or side effect, please call 1-888-267-1915.

Compare to Active Liquid-Gel Capsules Ingredient™

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP THIS BOX FOR IMPORTANT INFORMATION.

Distributed by: Watson Pharmaceuticals, Inc., Garden City, NY 11731
 *This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil®

LOT: 6 81131187471 3
 EXP: 438880

Drug Facts Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing

Active ingredient (in each capsule)	Purpose
Solubilized ibuprofen equal to 200 mg (NSAID)* (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug	Pain reliever/Fever reducer

Uses

■ temporarily relieves minor aches and pains due to	■ headache	■ toothache	■ backache
■ menstrual cramps	■ the common cold	■ muscular aches	■ minor pain of arthritis
■ temporarily reduces fever			

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Drug Facts (continued)

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



NDC 49035-133-18
 equate™
 Ibuprofen
 Capsules, 200mg
 Pain Reliever/ Fever

Reducer (NSAID)
 Liquid-Filled Capsules
 200mg, 80 SOFTGELS

equate™ NDC 49035-133-26

Compare to Advil® Liquid-Gels® Active Ingredient**

Ibuprofen Capsules, 200 mg
 Pain Reliever/ Fever Reducer (NSAID)

*Liquid-Filled Capsules

200 mg 80 SOFTGELS

Actual Size

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing

Drug Facts

Active ingredient (in each capsule)
 Sublimated ibuprofen equal to 200 mg ibuprofen (NSAID)*. **Pain reliever/fever reducer** (present as the free acid and potassium salt) ***nonsteroidal anti-inflammatory drug**

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
 - temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Use before or after heart surgery

Drug Facts (continued under label)

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

Distributed by: Wal-Mart Stores, Inc., Bentonville, AR 72716
 **This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil®.

Lot No.: 088857
 Exp Date: 6 8113118748 0

SEE THE FRONT

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use.
- store at 20°-25°C (68°-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, propylene glycol, potassium hydroxide, purified water, shellac, sorbitan monooleate, sorbitol sodium

Questions or comments? 1-888-287-1915

NDC 49035-133-26
 equate™
 Ibuprofen
 Capsules, 200mg
 Pain Reliever/ Fever

Reducer (NSAID)
Liquid-Filled Capsules
200mg, 160 SOFTGELS

equate™ NDC 49035-133-26

Compare to Advil® Liqui-Gels® Active Ingredient**

Ibuprofen
Capsules, 200 mg
Pain Reliever/ Fever Reducer (NSAID)

Liquid-Filled Capsules

200 mg **160 SOFTGELS**

Actual Size

Do Not Use if seal under bottle cap imprinted with -SEALED for YOUR PROTECTION- is broken or missing

Drug Facts

Active ingredient (in each capsule) Purpose
Sulfonized ibuprofen equal to 200 mg Ibuprofen (NSAID)* Pain reliever/Fever reducer (present as the free acid and potassium salt) nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to:

- headache
- toothache
- menstrual cramps
- the common cold
- minor pain of arthritis
- muscular aches
- temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer:

- right before or after heart surgery

Drug Facts (continued under label)

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

Distributed by: Wel-Mart Stores, Inc., Bentonville, AR 72716
 **This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil®.

458880
 6 81131 29798 1
 Lot No. Exp. Date
 FEEL FREE

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use.
- store at 20°-25°C (68°-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, propylene glycol, potassium hydroxide, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments? 1-888-287-1915

EQUATE IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49035-133
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GELATIN (UNII: 2G86QN327L)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0K00R)	
SHELLAC (UNII: 46N107B71O)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	GREEN	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	133
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-133-12	20 in 1 BOTTLE		
2	NDC:49035-133-14	40 in 1 BOTTLE		
3	NDC:49035-133-18	80 in 1 BOTTLE		
4	NDC:49035-133-26	160 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079205	07/02/2014	

Labeler - Walmart stores Inc (051957769)**Establishment**

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Revised: 7/2014

Walmart stores Inc