CONRX PAIN RELIEVER- acetaminophen, as pirin and caffeine tablet Eagle Distributors, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ConRxTM pain Reliever

Drug Facts

Active ingredients (in each tablet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)*	Pain reliever
Caffeine 65 mg	Pain reliever aid

^{*} nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - a cold
 - arthritis
 - muscular aches
 - sinusitis
 - toothache
 - premenstrual & menstrual cramps

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- adult takes more than 8 tablets in 24 hours, which is the maximum daily amount
- child takes more than 5 tablets in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have not been drinking fluids
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- vou have asthma
- you have lost a lot of fluids due to vomiting or diarrhea

Ask a doctor or pharmacist before use if you are taking

- any other drug containing an NSAID (prescription or nonprescription)
- a blood thinning (anticoagulant) or steroid drug
- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - ringing in the ears or loss of hearing occurs
 - painful area is red or swollen
 - pain gets worse or lasts for more than 10 days
 - fever gets worse or lasts for more than 3 days
 - any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to

use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 tablets every 6 hours; not more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at controlled room temperature 20°-25°C (68°-77°F)
- read all product information before using. Keep this box for important information.

Inactive ingredients

benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions or comments?

1-800-570-8650

PRINCIPAL DISPLAY PANEL - 50 Pouch Box

See New Warnings Information & Directions

Compare to the Active Ingredients in $EXCERDRIN*_{\mathbb{R}}$

ConRxTM

EXTRA STRENGTH PAIN RELIEVER

Acetaminophen, Asprin (NSAID) and Caffeine

TO OPEN
PUSH IN TAB AND PULL OUT

Compare to the Active Ingredients in $EXCEDRIN*_{\mathbb{R}}$

50 Pouches of 2 Caplets Each

See New Warnings Information & Directions

Compare to the Active Ingredients in



EXTRA STRENGTH PAIN RELIEVER See New Warnings Information & Directions

Compare to the Active Ingredients in EXCEDRIN.*



EXTRA STRENGTH PAIN RELIEVER

Acetaminophen, Asprin (NSAID) and Caffeine

Acetaminophen, Asprin (NSAID) and Caffeine



TO OPEN
PUSH IN TAB AND PULL OUT

Compare to the Active Ingredients in

EXCEDRIN®

50 Pouches of 2 Caplets Each



DO Pouches of 2 Caplets Each

Acetaminophen, Asprin (NSAID) and Caffeine





Compare to the Active Ingredients in

See New Warnings Information & Directions

THIS DISPENSER CONTAINS TAMPER EVIDENT PACKETS.

DO NOT USE IF SEAL ON INDIVIDUAL PACKET IS BROKEN,			
Drug Facts			
Active ingredients (in each caplet)	Purposes		
Acetaminophen 250 mg	Pain reliever		
Aspirin 250 mg (NSAID)*	Pain reliever		
Caffeine 65 mg "nonsteroidal anti-inflammatory drug	Pain reflever aid		
Uses - temporarily relieves minor aches and pains due to: - headache - a colid - arthritis - muscular aches - sinusitis - toothache - premenstrual & menstrual cramps			

See New Warnings Information & Directions

Compare to the Active Ingredients in





Warninge

Reger's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare

but serious Briess.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives - facial swelling - asthma (wheezing) - shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- acult takes more than 8 tablets in 24 hours, which is the maximum daily amount

- · child takes more than 5 tablets in 24 hours
- with other drugs containing acetaminophe
- 3 or more alcoholic drinks every day while using this product
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older
 have had stomach ulcers or bleeding problems
 take a blood thinning (anticoagulant) or steroid drug

 are age out or oper - nave has some nucers or incesting processing or take other drugs containing preception in Soft (age).
 base a opport or others)
 have 3 or more alcoholic drinks every day white using this product
 have 3 or more alcoholic drinks every day white using this product was much caffeine as a cup or offee. Limit the use of caffeine-containing medications, floots, or beverages while taking this product because too much caffeine may cause nervousness, irritability, steeplissness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

- Ask a doctor before use if + you have liver disease
- Ask a doctor before use if

 you have liver disease

 you have a history of stomach problems, such as hearthurn

 you have a history of stomach problems, such as hearthurn

 you have a history of stomach problems, such as hearthurn

 you have lost a lot of fluids due to vomiting or diarrhea

 *you have lost a lot of fluids due to vomiting or diarrhea

Ask a doctor or pharmacist before use if you are taking

- any other drug containing an INSAID (prescription or nonprescription)
 a prescription drug for diabetes, gout, or arthritis
 any other drug, or are under a doctor's care for

 Stop use and ask a doctor if an altergic reaction occurs. Seek medical help right away. . any other drug, or are under a doctor's care for any serious condition
- you experience any of the following signs of stomach bleeding:
 feel faint
 vomit blood
 have bloody or black stools
 have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
 pain gets worse or lasts for more than 10 days
 + fever gets worse or lasts for more than 3 days

If pregnant or breast-feeding, ask a health professional before use, it is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions - do not use more than directed

- drink a full class of water with each dose
- adults and children 12 years and over; take 2 tablets every 6 hours; not more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information + store at controlled room temperature 20°-25°C (68°-77°F)

read all product information before using, Keep this box for important information

Inactive ingredients benzoic sold, carreade wax. FD&C blue #1, hydroxypropylcelulose, hygromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Acetaminophen, Asprin (NSAID) and Caffeine

*This product is not manufactured or distributed by: Novartis Consumer Health, Inc. Eagle Distributors,Inc.,/ConRx™ does not own the Excedrin® trademark.



CONRX PAIN RELIEVER

acetaminophen, aspirin, and caffeine tablet

Product Information			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:68737-234
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Acetaminophen (Acetaminophen)	Acetaminophen	250 mg	
Aspirin (Aspirin)	Aspirin	250 mg	
Caffeine (Caffeine)	Caffeine	65 mg	

Inactive Ingredients					
Ingredient Name	Strength				
CARNAUBA WAX					
BENZOIC ACID					
HYDRO XYPRO PYL CELLULO SE (TYPE H)					
FD&C BLUE NO. 1					
HYPROMELLOSES					
LIGHT MINERAL OIL					
CELLULO SE, MICRO CRYSTALLINE					
POLYSORBATE 20					

POVIDONES	
POLYETHYLENE GLYCOL 6000	
DIMETHICONE	
SORBITAN MONOLAURATE	
STEARIC ACID	
TITANIUM DIO XIDE	

Product Characteristics			
Color	BLUE	Score	2 pieces
Shape	OVAL	Size	18 mm
Flavor		Imprint Code	CRX
Contains			

F	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:68737-234-18	50 in 1 BOX				
1		2 in 1 POUCH				

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part343	02/15/2013		

Labeler - Eagle Distributors,Inc. (929837425)

Revised: 2/2013 Eagle Distributors,Inc.