

# DICYCLOMINE HYDROCHLORIDE- dicyclomine hydrochloride tablet

## Preferred Pharmaceuticals, Inc.

-----

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DICYCLOMINE HYDROCHLORIDE safely and effectively. See full prescribing information for DICYCLOMINE HYDROCHLORIDE.

DICYCLOMINE HYDROCHLORIDE capsules, for oral use  
DICYCLOMINE HYDROCHLORIDE tablets, for oral use  
Initial U.S. Approval: 1950

### INDICATIONS AND USAGE

Dicyclomine hydrochloride is an antispasmodic and anticholinergic (antimuscarinic) agent indicated for the treatment of functional bowel/irritable bowel syndrome (**Error! Hyperlink reference not valid.**) (1)

### DOSAGE AND ADMINISTRATION

Dosage for dicyclomine hydrochloride must be adjusted to individual patient needs (**Error! Hyperlink reference not valid.**). If a dose is missed, patients should continue the normal dosing schedule (**Error! Hyperlink reference not valid.**).

#### Oral in adults (2.1):

- Starting dose: 20 mg four times a day. After a week treatment with the starting dose, the dose may be escalated to 40 mg four times a day, unless side effects limit dosage escalation
- Discontinue dicyclomine hydrochloride if efficacy not achieved or side effects require doses less than 80 mg per day after two weeks of treatment

### DOSAGE FORMS AND STRENGTHS

- Dicyclomine hydrochloride capsules 10 mg (**Error! Hyperlink reference not valid.**)
- Dicyclomine hydrochloride tablets 20 mg (**Error! Hyperlink reference not valid.**)

### CONTRAINDICATIONS

- |   |  |
|---|--|
| • Infants less than 6 months of age ( <b>Error! Hyperlink reference not valid.</b> )                  | • Glaucoma ( <b>Error! Hyperlink reference not valid.</b> )  |
| • Nursing mothers ( <b>Error! Hyperlink reference not valid.</b> )                                    | • Obstructive uropathy ( <b>Error! Hyperlink reference not valid.</b> )                              |
| • Unstable cardiovascular status in acute hemorrhage ( <b>Error! Hyperlink reference not valid.</b> ) | • Obstructive disease of the gastrointestinal tract ( <b>Error! Hyperlink reference not valid.</b> ) |
| • Myasthenia gravis ( <b>Error! Hyperlink reference not valid.</b> )                                  | • Severe ulcerative colitis ( <b>Error! Hyperlink reference not valid.</b> )                         |
- (4)
- Reflux esophagitis (**Error! Hyperlink reference not valid.**)

### WARNINGS AND PRECAUTIONS

- Cardiovascular conditions: worsening of conditions (5.2)
- Peripheral and central nervous system: heat prostration can occur with drug use (fever and heat stroke due to decreased sweating); drug should be discontinued and supportive measures instituted (5.3)

- Psychosis and delirium have been reported in patients sensitive to anticholinergic drugs (such as elderly patients and/or in patients with mental illness): signs and symptoms resolve within 12 to 24 hours after discontinuation of dicyclomine hydrochloride (5.3)
- Myasthenia Gravis: overdose may lead to muscular weakness and paralysis. Dicyclomine hydrochloride should be given to patients with myasthenia gravis only to reduce adverse muscarinic effects of an anticholinesterase (5.4)
- Incomplete intestinal obstruction: diarrhea may be an early symptom especially in patients with ileostomy or colostomy. Treatment with dicyclomine hydrochloride would be inappropriate and possibly fatal (5.5)
- Salmonella dysenteric patients: due to risk of toxic megacolon (5.6)
- Ulcerative colitis: Dicyclomine hydrochloride should be used with caution in these patients; large doses may suppress intestinal motility or aggravate the serious complications of toxic megacolon (5.7)
- Prostatic hypertrophy: Dicyclomine hydrochloride should be used with caution in these patients; may lead to urinary retention (5.8)
- Hepatic and renal disease: should be used with caution (5.9)
- Geriatric: use with caution in elderly who may be more susceptible to dicyclomine hydrochloride's adverse events (5.10)

#### ----- ADVERSE REACTIONS -----

The most serious adverse reactions include cardiovascular and central nervous system symptoms. The most common adverse reactions (> 5% of patients) are dizziness, dry mouth, vision blurred, nausea, somnolence, asthenia and nervousness (**Error! Hyperlink reference not valid.**) (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Teva Pharmaceuticals USA, Inc. at 1-888-838-2872 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch). (6)**

#### ----- DRUG INTERACTIONS -----

- Antiglaucoma agents: anticholinergics antagonize antiglaucoma agents and may increase intraocular pressure (**Error! Hyperlink reference not valid.**)
- Anticholinergic agents: may affect the gastrointestinal absorption of various drugs; may also increase certain actions or side effects of other anticholinergic drugs (**Error! Hyperlink reference not valid.**)
- Antacids: interfere with the absorption of anticholinergic agents (**Error! Hyperlink reference not valid.**)

#### ----- USE IN SPECIFIC POPULATIONS -----

- Pregnancy: use only if clearly needed (**Error! Hyperlink reference not valid.**)
- Pediatric Use: Safety and effectiveness not established (**Error! Hyperlink reference not valid.**)
- Hepatic and renal impairment: caution must be taken with patients with significantly impaired hepatic and renal function (8.6)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 4/2024**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

#### 2.1 Oral Dosage and Administration in Adults

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

#### 5.2 Cardiovascular Conditions

#### 5.3 Peripheral and Central Nervous System

#### 5.4 Myasthenia Gravis

- 5.5 Intestinal Obstruction
- 5.6 Toxic Dilatation of Intestine megacolon
- 5.7 Ulcerative Colitis
- 5.8 Prostatic Hypertrophy
- 5.9 Hepatic and Renal Disease
- 5.10 Geriatric Population

## **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience
- 6.3 Adverse Reactions Reported with Similar Drugs with Anticholinergic/Antispasmodic Action

## **7 DRUG INTERACTIONS**

- 7.1 Antiglaucoma Agents
- 7.2 Other Drugs with Anticholinergic Activity
- 7.3 Other Gastrointestinal Motility Drugs
- 7.4 Effect of Antacids
- 7.5 Effect on Absorption of Other Drugs
- 7.6 Effect on Gastric Acid Secretion

## **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

## **14 CLINICAL STUDIES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

- 17.2 Use in Infants
- 17.3 Use in Nursing Mothers
- 17.4 Peripheral and Central Nervous System

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Dicyclomine hydrochloride is indicated for the treatment of patients with functional

bowel/irritable bowel syndrome.

## 2 DOSAGE AND ADMINISTRATION

Dosage must be adjusted to individual patient needs.

### 2.1 Oral Dosage and Administration in Adults

The recommended initial dose is 20 mg four times a day.

After one week treatment with the initial dose, the dose may be increased to 40 mg four times a day unless side effects limit dosage escalation.

If efficacy is not achieved within 2 weeks or side effects require doses below 80 mg per day, the drug should be discontinued. Documented safety data are not available for doses above 80 mg daily for periods longer than 2 weeks.

## 3 DOSAGE FORMS AND STRENGTHS

- Dicyclomine hydrochloride USP, 10 mg capsules: Clear Dark Blue cap/Clear Dark Blue body hard gelatin capsules, imprinted with white ink **WATSON** over **794** on cap and 10 mg on the body
- Dicyclomine hydrochloride USP, 20 mg tablets: Blue, round, unscored, flat-faced, beveled-edge tablets, debossed **WATSON** and **795** on the periphery on one side and plain on the other side

## 4 CONTRAINDICATIONS

Dicyclomine hydrochloride is contraindicated in infants less than 6 months of age [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**], nursing mothers [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**], and in patients with:

- unstable cardiovascular status in acute hemorrhage
- myasthenia gravis [see **Error! Hyperlink reference not valid. (5.4)**]
- glaucoma [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)** and **Error! Hyperlink reference not valid. (7.1)**]
- obstructive uropathy [see **Error! Hyperlink reference not valid. (5.8)**]
- obstructive disease of the gastrointestinal tract [see **Error! Hyperlink reference not valid. (5.5)**]
- severe ulcerative colitis [see **Error! Hyperlink reference not valid. (5.7)**]
- reflux esophagitis

## 5 WARNINGS AND PRECAUTIONS

### 5.2 Cardiovascular Conditions

Dicyclomine hydrochloride needs to be used with caution in conditions characterized by tachyarrhythmia such as thyrotoxicosis, congestive heart failure and in cardiac surgery, where they may further accelerate the heart rate. Investigate any tachycardia before

administration of dicyclomine hydrochloride. Care is required in patients with coronary heart disease, as ischemia and infarction may be worsened, and in patients with hypertension [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**].

### 5.3 Peripheral and Central Nervous System

The peripheral effects of dicyclomine hydrochloride are a consequence of their inhibitory effect on muscarinic receptors of the autonomic nervous system. They include dryness of the mouth with difficulty in swallowing and talking, thirst, reduced bronchial secretions, dilatation of the pupils (mydriasis) with loss of accommodation (cycloplegia) and photophobia, flushing and dryness of the skin, transient bradycardia followed by tachycardia, with palpitations and arrhythmias, and difficulty in micturition, as well as reduction in the tone and motility of the gastrointestinal tract leading to constipation [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**].

In the presence of high environmental temperature heat prostration can occur with drug use (fever and heat stroke due to decreased sweating). It should also be used cautiously in patients with fever. If symptoms occur, the drug should be discontinued and supportive measures instituted. Because of the inhibitory effect on muscarinic receptors within the autonomic nervous system, caution should be taken in patients with autonomic neuropathy. Central nervous system (CNS) signs and symptoms include confusional state, disorientation, amnesia, hallucinations, dysarthria, ataxia, coma, euphoria, fatigue, insomnia, agitation and mannerisms, and inappropriate affect.

Psychosis and delirium have been reported in sensitive individuals (such as elderly patients and/or in patients with mental illness) given anticholinergic drugs. These CNS signs and symptoms usually resolve within 12 to 24 hours after discontinuation of the drug.

Dicyclomine hydrochloride may produce drowsiness, dizziness or blurred vision. The patient should be warned not to engage in activities requiring mental alertness, such as operating a motor vehicle or other machinery or performing hazardous work while taking dicyclomine hydrochloride.

### 5.4 Myasthenia Gravis

With overdosage, a curare-like action may occur (i.e., neuromuscular blockade leading to muscular weakness and possible paralysis). It should not be given to patients with myasthenia gravis except to reduce adverse muscarinic effects of an anticholinesterase [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**].

### 5.5 Intestinal Obstruction

Diarrhea may be an early symptom of incomplete intestinal obstruction, especially in patients with ileostomy or colostomy. In this instance, treatment with this drug would be inappropriate and possibly harmful [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**].

Rarely development of Ogilvie's syndrome (colonic pseudo-obstruction) has been reported. Ogilvie's syndrome is a clinical disorder with signs, symptoms, and

radiographic appearance of an acute large bowel obstruction but with no evidence of distal colonic obstruction

### 5.6 Toxic Dilatation of Intestine megacolon

Toxic dilatation of intestine and intestinal perforation is possible when anticholinergic agents are administered in patients with Salmonella dysentery.

### 5.7 Ulcerative Colitis

Caution should be taken in patients with ulcerative colitis. Large doses may suppress intestinal motility to the point of producing a paralytic ileus and the use of this drug may precipitate or aggravate the serious complication of toxic megacolon [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**].

Dicyclomine hydrochloride is contraindicated in patients with severe ulcerative colitis [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**].

### 5.8 Prostatic Hypertrophy

Dicyclomine hydrochloride should be used with caution in patients with known or suspected prostatic enlargement, in whom prostatic enlargement may lead to urinary retention [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**].

### 5.9 Hepatic and Renal Disease

Dicyclomine hydrochloride should be used with caution in patients with known hepatic and renal impairment.

### 5.10 Geriatric Population

Dicyclomine hydrochloride should be used with caution in elderly who may be more susceptible to its adverse effects.

## 6 ADVERSE REACTIONS

The pattern of adverse effects seen with dicyclomine is mostly related to its pharmacological actions at muscarinic receptors [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**]. They are a consequence of the inhibitory effect on muscarinic receptors within the autonomic nervous system. These effects are dose-related and are usually reversible when treatment is discontinued.

The most serious adverse reactions reported with dicyclomine hydrochloride include cardiovascular and central nervous system symptoms [see **Error! Hyperlink reference not valid. (5.2, 5.3)**].

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The data described below reflect exposure in controlled clinical trials involving over 100 patients treated for functional bowel/irritable bowel syndrome with dicyclomine hydrochloride at initial doses of 160 mg daily (40 mg four times a day)

In these trials most of the side effects were typically anticholinergic in nature and were reported by 61% of the patients. Table 1 presents adverse reactions (*MedDRA 13.0* preferred terms) by decreasing order of frequency in a side-by-side comparison with placebo.

**Table 1: Adverse reactions experienced in controlled clinical trials with decreasing order of frequency**

<b>MedDRA Preferred Term</b>	<b>Dicyclomine Hydrochloride (40 mg four times a day) %</b>	<b>Placebo %</b>
Dry Mouth	33	5
Dizziness	40	5
Vision blurred	27	2
Nausea	14	6
Somnolence	9	1
Asthenia	7	1
Nervousness	6	2

Nine percent (9%) of patients were discontinued from dicyclomine hydrochloride because of one or more of these side effects (compared with 2% in the placebo group). In 41% of the patients with side effects, side effects disappeared or were tolerated at the 160 mg daily dose without reduction. A dose reduction from 160 mg daily to an average daily dose of 90 mg was required in 46% of the patients with side effects who then continued to experience a favorable clinical response; their side effects either disappeared or were tolerated.

## 6.2 Postmarketing Experience

The following adverse reactions, presented by system organ class in alphabetical order, have been identified during post approval use of dicyclomine hydrochloride. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Cardiac disorders: palpitations, tachyarrhythmias
- Eye disorders: cycloplegia, mydriasis, vision blurred
- Gastrointestinal disorders: abdominal distension, abdominal pain, constipation, dry mouth, dyspepsia, nausea, vomiting
- General disorders and administration site conditions: fatigue, malaise
- Immune System Disorders: drug hypersensitivity including face oedema, angioedema, anaphylactic shock
- Nervous system disorders: dizziness, headache, hallucinations insomnia, somnolence, syncope
- Psychiatric disorders: As with the other anti-cholinergic drugs, cases of delirium or

symptoms of delirium such as amnesia (or transient global amnesia), agitation, confusional state, delusion, disorientation, hallucination (including visual hallucination) as well as mania, mood altered and pseudodementia, have been reported with the use of Dicyclomine. Nervousness and insomnia have also been reported.

- Reproductive system and breast disorders: suppressed lactation
- Respiratory, thoracic and mediastinal disorders: dyspnoea, nasal congestion
- Skin and subcutaneous tissue disorder: dermatitis allergic, erythema, rash

### **6.3 Adverse Reactions Reported with Similar Drugs with Anticholinergic/Antispasmodic Action**

Gastrointestinal: anorexia,

Central Nervous System: tingling, numbness, dyskinesia, speech disturbance, insomnia

Peripheral Nervous System: With overdosage, a curare-like action may occur (i.e., neuromuscular blockade leading to muscular weakness and possible paralysis)

Ophthalmologic: diplopia, increased ocular tension

Dermatologic/Allergic: urticaria, itching, and other dermal manifestations

Genitourinary: urinary hesitancy, urinary retention in patients with prostatic hypertrophy

Cardiovascular: hypertension

Respiratory: apnea

Other: decreased sweating, sneezing, throat congestion, impotence. With the injectable form, there may be temporary sensation of light-headedness. Some local irritation and focal coagulation necrosis may occur following the intramuscular injection of dicyclomine hydrochloride.

## **7 DRUG INTERACTIONS**

### **7.1 Antiglaucoma Agents**

Anticholinergics antagonize the effects of antiglaucoma agents. Anticholinergic drugs in the presence of increased intraocular pressure may be hazardous when taken concurrently with agents such as corticosteroids. Use of dicyclomine hydrochloride in patients with glaucoma is not recommended [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**].

### **7.2 Other Drugs with Anticholinergic Activity**

The following agents may increase certain actions or side effects of anticholinergic drugs including dicyclomine hydrochloride: amantadine, antiarrhythmic agents of Class I (e.g., quinidine), antihistamines, antipsychotic agents (e.g., phenothiazines), benzodiazepines, MAO inhibitors, narcotic analgesics (e.g., meperidine), nitrates and nitrites, sympathomimetic agents, tricyclic antidepressants, and other drugs having



anticholinergic activity.

### **7.3 Other Gastrointestinal Motility Drugs**

Interaction with other gastrointestinal motility drugs may antagonize the effects of drugs that alter gastrointestinal motility, such as metoclopramide.

### **7.4 Effect of Antacids**

Because antacids may interfere with the absorption of anticholinergic agents including dicyclomine hydrochloride, simultaneous use of these drugs should be avoided.

### **7.5 Effect on Absorption of Other Drugs**

Anticholinergic agents may affect gastrointestinal absorption of various drugs by affecting on gastrointestinal motility, such as slowly dissolving dosage forms of digoxin; increased serum digoxin concentration may result.

### **7.6 Effect on Gastric Acid Secretion**

The inhibiting effects of anticholinergic drugs on gastric hydrochloric acid secretion are antagonized by agents used to treat achlorhydria and those used to test gastric secretion.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

Teratogenic Effects: Pregnancy Category B

Adequate and well-controlled studies have not been conducted with dicyclomine hydrochloride in pregnant women at the recommended doses of 80 to 160 mg/day. However, epidemiologic studies did not show an increased risk of structural malformations among babies born to women who took products containing dicyclomine hydrochloride at doses up to 40 mg/day during the first trimester of pregnancy. Reproduction studies have been performed in rats and rabbits at doses up to 33 times the maximum recommended human dose based on 160 mg/day (3 mg/kg) and have revealed no evidence of harm to the fetus due to dicyclomine. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### **8.3 Nursing Mothers**

Dicyclomine hydrochloride is contraindicated in women who are breastfeeding. Dicyclomine hydrochloride is excreted in breastmilk. Because of the potential for serious adverse reactions in breastfed infants from dicyclomine hydrochloride, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother [see **Error! Hyperlink reference not valid.** (**Error! Hyperlink reference not valid.**)].

### **8.4 Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

Dicyclomine hydrochloride is contraindicated in infants less than 6 months of age [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid.)**]. There are published cases reporting that the administration of dicyclomine hydrochloride syrup to infants has been followed by serious respiratory symptoms (dyspnea, shortness of breath, breathlessness, respiratory collapse, apnea and asphyxia), seizures, syncope, pulse rate fluctuations, muscular hypotonia, and coma, and death, however; no causal relationship has been established.

## 8.5 Geriatric Use

Clinical studies of dicyclomine hydrochloride did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range in adults, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

## 8.6 Renal Impairment

Effects of renal impairment on PK, safety and efficacy of dicyclomine hydrochloride has not been studied. Dicyclomine hydrochloride drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Dicyclomine hydrochloride should be administered with caution in patients with renal impairment.

## 8.7 Hepatic Impairment

Effects of renal impairment on PK, safety and efficacy of dicyclomine hydrochloride have not been studied. Dicyclomine hydrochloride should be administered with caution in patients with hepatic impairment.

# 10 OVERDOSAGE

In case of an overdose, patients should contact a physician, poison control center (1-800-222-1222), or emergency room.

The signs and symptoms of overdose include: headache; nausea; vomiting; blurred vision; dilated pupils; hot, dry skin; dizziness; dryness of the mouth; difficulty in swallowing; and CNS stimulation including convulsion. A curare-like action may occur (i.e., neuromuscular blockade leading to muscular weakness and possible paralysis).

One reported event included a 37-year-old who reported numbness on the left side, cold fingertips, blurred vision, abdominal and flank pain, decreased appetite, dry mouth, and nervousness following ingestion of 320 mg daily (four 20 mg tablets four times daily). These events resolved after discontinuing the dicyclomine.

The acute oral LD<sub>50</sub> of the drug is 625 mg/kg in mice.

The amount of drug in a single dose that is ordinarily associated with symptoms of

overdosage or that is likely to be life-threatening, has not been defined. The maximum human oral dose recorded was 600 mg by mouth in a 10-month-old child and approximately 1500 mg in an adult, each of whom survived. In three of the infants who died following administration of dicyclomine hydrochloride [see **Error! Hyperlink reference not valid.** (5.1)], the blood concentrations of drug were 200, 220, and 505 ng/mL.

It is not known if dicyclomine hydrochloride is dialyzable.


Treatment should consist of gastric lavage, emetics, and activated charcoal. Sedatives (e.g., short-acting barbiturates, benzodiazepines) may be used for management of overt signs of excitement. If indicated, an appropriate parenteral cholinergic agent may be used as an antidote.

## 11 DESCRIPTION

Dicyclomine hydrochloride, USP is an antispasmodic and anticholinergic (antimuscarinic) agent available in the following dosage forms:

- Dicyclomine hydrochloride capsules for oral use contain 10 mg dicyclomine hydrochloride, USP. Dicyclomine hydrochloride 10 mg capsules also contain inactive ingredients: sodium lauryl sulfate, D&C Red #28, FD&C Blue #1, FD&C Red #40, gelatin, magnesium stearate, pregelatinized starch, silicon dioxide.
- Dicyclomine hydrochloride tablets for oral use contain 20 mg dicyclomine hydrochloride, USP. Dicyclomine hydrochloride 20 mg tablets also contain inactive ingredients: colloidal silicon dioxide, corn starch, FD&C Blue #1, hypromellose, lactose monohydrate, sodium starch glycolate, and stearic acid.

Dicyclomine hydrochloride, USP is [bicyclohexyl]-1-carboxylic acid, 2-(diethylamino) ethyl ester, hydrochloride, with a molecular formula of  $C_{19}H_{35}NO_2 \cdot HCl$  and the following structural formula:



Molecular weight: 345.95

Dicyclomine hydrochloride, USP occurs as a fine, white, crystalline, practically odorless powder with a bitter taste. It is soluble in water, freely soluble in alcohol and chloroform, and very slightly soluble in ether.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Dicyclomine relieves smooth muscle spasm of the gastrointestinal tract. Animal studies indicate that this action is achieved via a dual mechanism:

- a specific anticholinergic effect (antimuscarinic) at the acetylcholine-receptor sites with approximately 1/8 the milligram potency of atropine (*in vitro*, guinea pig ileum); and
- a direct effect upon smooth muscle (musculotropic) as evidenced by dicyclomine's antagonism of bradykinin- and histamine-induced spasms of the isolated guinea pig ileum.

Atropine did not affect responses to these two agonists. *In vivo* studies in cats and dogs showed dicyclomine to be equally potent against acetylcholine (ACh)- or barium chloride (BaCl<sub>2</sub>)-induced intestinal spasm while atropine was at least 200 times more potent against effects of ACh than BaCl<sub>2</sub>. Tests for mydriatic effects in mice showed that dicyclomine was approximately 1/500 as potent as atropine; antisialagogue tests in rabbits showed dicyclomine to be 1/300 as potent as atropine.

## 12.2 Pharmacodynamics

Dicyclomine hydrochloride can inhibit the secretion of saliva and sweat, decrease gastrointestinal secretions and motility, cause drowsiness, dilate the pupils, increase heart rate, and depress motor function

## 12.3 Pharmacokinetics

### Absorption and Distribution

In man, dicyclomine is rapidly absorbed after oral administration, reaching peak values within 60-90 minutes. Mean volume of distribution for a 20 mg oral dose is approximately 3.65 L/kg suggesting extensive distribution in tissues.

Elimination The metabolism of dicyclomine was not studied. The principal route of excretion is via the urine (79.5% of the dose). Excretion also occurs in the feces, but to a lesser extent (8.4%). Mean half-life of plasma elimination in one study was determined to be approximately 1.8 hours when plasma concentrations were measured for 9 hours after a single dose. In subsequent studies, plasma concentrations were followed for up to 24 hours after a single dose, showing a secondary phase of elimination with a somewhat longer half-life.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies have not been conducted to evaluate the carcinogenic potential of dicyclomine. In studies in rats at doses of up to 100 mg/kg/day, dicyclomine produced no deleterious effects on breeding, conception, or parturition.

## 14 CLINICAL STUDIES

In controlled clinical trials involving over 100 patients who received drug, 82% of patients treated for functional bowel/irritable bowel syndrome with dicyclomine hydrochloride at initial doses of 160 mg daily (40 mg four times daily) demonstrated a favorable clinical response compared with 55% treated with placebo ( $p < 0.05$ ).

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Dicyclomine hydrochloride capsules, USP and dicyclomine hydrochloride tablets, USP are supplied as follows:

### Dicyclomine hydrochloride tablets, USP

20 mg tablets: Blue, round, unscored, flat-faced, beveled-edge tablets, debossed **WATSON** and **795** on the periphery on one side and plain on the other side, in;

Bottles of 20 - 68788-9525-2

Bottles of 30 - 68788-9525-3

Store at 20° - 25°C (68° - 77°F). [See USP controlled room temperature.]

Keep out of the reach of children.

Dispense in a well-closed container as defined in USP/NF.

## 17 PATIENT COUNSELING INFORMATION

### 17.2 Use in Infants

Inform parents and caregivers not to administer dicyclomine hydrochloride in infants less than 6 months of age [see **Error! Hyperlink reference not valid.Error! Hyperlink reference not valid.Error! Hyperlink reference not valid.**].

### 17.3 Use in Nursing Mothers

Advise lactating women that dicyclomine hydrochloride should not be used while breastfeeding their infants [see **Error! Hyperlink reference not valid. (Error! Hyperlink reference not valid., Error! Hyperlink reference not valid.)**].

### 17.4 Peripheral and Central Nervous System

In the presence of a high environmental temperature, heat prostration can occur with dicyclomine hydrochloride use (fever and heat stroke due to decreased sweating). If symptoms occur, the drug should be discontinued and a physician contacted. Dicyclomine hydrochloride may produce drowsiness or blurred vision. The patient should be warned not to engage in activities requiring mental alertness, such as operating a motor vehicle or other machinery or to perform hazardous work while taking dicyclomine hydrochloride [see **Error! Hyperlink reference not valid. (5.3)**].

Manufactured For:

**Teva Pharmaceuticals USA, Inc.**

Parsippany, NJ 07054

Rev. B 11/2020

**Repackaged By: Preferred Pharmaceuticals Inc.**

## PACKAGE LABEL PRINCIPAL DISPLAY PANEL

NDC 68788-9825

Dicyclomine Hydrochloride Tablets, USP

20 mg

Rx only

teva

Repackaged By: Preferred Pharmaceuticals Inc.

Dicyclomine HCl  
Tablets, USP

20mg

Generic for Bentyl

Each tablet contains: Dicyclomine Hydrochloride USP 20mg

Pkg Size: Exp Date:  
Lot#: Batch#: Ins:  
Mfg: Actavis Pharma, Inc.  
Prod#: Warning

Store at 20°-25°C (68°- 77°F). See USP Controlled Room Temperature. Keep this and all medication out of the reach of children. Keep only the amount of medicine you need for your condition. This tablet is round, white and imprinted with WATSON 795



Directions English  
May cause drowsiness.  
Take \_\_\_\_\_ tablet(s)  
every \_\_\_\_\_ hours.  
Use as directed by your doctor



Instrucciones Espanol:  
Puede causar  
somnolencia.  
Toma \_\_\_\_\_ tableta(s)  
cada \_\_\_\_\_ horas.  
Uso según lo dirigido  
por su doctor

CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed

Dicyclomine HCl Tablets, USP  
20mg  
Qty: Ins:  
Lot#: Bat#:

Prod# (NDC):  
Dicyclomine HCl Tablets, USP  
20mg  
Qty: Ins:  
Lot#: Bat#:  
Prod# (NDC):

Dicyclomine HCl Tablets, USP  
20mg  
Qty: Insurance NDC:  
Lot#: Bat#:

Dicyclomine HCl Tablets, USP  
20mg  
Qty: Ins:  
Lot#: Bat#:  
Prod# (NDC):

Log  
Chart  
Billing  
Patient

DICYCLOMINE HYDROCHLORIDE

dicyclomine hydrochloride tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68788-9525(NDC:0591-0795)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
DICYCLOMINE HYDROCHLORIDE (UNII: CQ903KQA31) (DICYCLOMINE - UNII:4KV4X8IF6V)		DICYCLOMINE HYDROCHLORIDE	20 mg
Inactive Ingredients			
Ingredient Name			Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STARCH, CORN (UNII: O8232NY3SJ)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
ALUMINUM OXIDE (UNII: LMI26O6933)			
HYPROMELLOSE 2910 (15 MPA.S) (UNII: 36SFW2JZ0W)			
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			

STEARIC ACID (UNII: 4ELV7Z65AP)				
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
Product Characteristics				
Color	BLUE	Score	no score	
Shape	ROUND	Size	7mm	
Flavor		Imprint Code	WATSON;795	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-9525-2	20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/21/2014	
2	NDC:68788-9525-3	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/21/2014	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA085223	02/21/2014	

**Labeler** - Preferred Pharmaceuticals, Inc. (791119022)

**Registrant** - Preferred Pharmaceuticals, Inc. (791119022)

## Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals, Inc.		791119022	REPACK(68788-9525)

Revised: 4/2024

Preferred Pharmaceuticals, Inc.