

**VICKS DAYQUIL NYQUIL HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD AND FLU PLUS CONGESTION- acetaminophen, dextromethorphan hbr, diphenhydramine hcl, phenylephrine hcl  
Procter & Gamble Manufactura S de RL de CV**

-----

**Vicks® DayQuil™ HOT REMEDY Vicks® NyQuil™ HOT REMEDY  
RAPID RELIEF  
MULTI-SYMPTOM  
COLD & FLU PLUS CONGESTION**

**Vicks® DayQuil™ HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD & FLU +  
CONGESTION**

## **Drug Facts**

### **DayQuil™**

#### **Active Ingredient (in each Packet)**

**Acetaminophen** 500 mg

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

#### **Purpose**

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

#### **Uses**

temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- nasal congestion
- sinus congestion & pressure
- reduces swelling of nasal passages

- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

## **Warnings**

### **Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Allergy Alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**When using this product,  
do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

---

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

---

**Other information**

- **each packet contains:** potassium 10 mg

- phenylketonurics: contains phenylalanine 64 mg per dose
- do not exceed 25°C.

**Inactive ingredients:**

acesulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, tribasic calcium phosphate

**Questions?**

**1-800-362-1683**

MADE IN MEXICO

DIST. BY PROCTER & GAMBLE,

CINCINNATI OH 45202

**Vicks® NyQuil™ HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD & FLU + CONGESTION**

**Drug Facts**

**NyQuil™**

**Active Ingredient (in each Packet)**

**Acetaminophen** 650 mg

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

**Purpose**

Pain reliever/fever reducer

Antihistamine/Cough suppressant

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor throat & bronchial irritation

- cough to help you sleep
- nasal congestion
- sinus congestion & pressure
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- runny nose
- sneezing

## **Warnings**

### **Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Allergy Alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease

- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **Do not use more than directed**
- excitability may occur especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

---

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

---

### Other information

- **each packet contains:**potassium 10 mg
- phenylketonurics: contains phenylalanine 68 mg per dose
- do not exceed 25°C

### Inactive ingredients

acesulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, tribasic calcium phosphate

### Questions?

1-800-362-1683

**Tamper Evident:**Do not use if sachet is opened or torn

MADE IN MEXICO

DIST. BY PROCTER & GAMBLE,

CINCINNATI OH 45202

**PRINCIPAL DISPLAY PANEL - 16 packet package**

MAX STRENGTH

Vicks® DayQuil™ HOT REMEDY

RAPID RELIEF

MULTI-SYMPTOM

COLD & FLU + CONGESTION

**Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl**

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Cough

*Non-Drowsy*

Vicks® NyQuil™ HOT REMEDY

RAPID RELIEF

MULTI-SYMPTOM

COLD & FLU + CONGESTION

**Acetaminophen, Diphenhydramine HCl, Phenylephrine HCl**

Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose, Cough

Nasal Congestion, Sinus Pressure

*Nighttime Relief*

HONEY LEMON FLAVORED

10 DAYQUIL POWDER PACKETS; 6 NYQUIL POWDER PACKETS; 16 TOTAL POWDER  
PACKETS



# VICKS DAYQUIL NYQUIL HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD AND FLU PLUS CONGESTION

acetaminophen, dextromethorphan hbr, diphenhydramine hcl, phenylephrine hcl kit

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58933-542
---------------------	----------------	---------------------------	---------------

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-542-16	1 in 1 PACKAGE; Type 0: Not a Combination Product	07/20/2022	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 PACKET	5 g

**Part 1 of 2****VICKS NYQUIL HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD AND FLU PLUS CONGESTION**

acetaminophen, diphenhydramine hcl, phenylephrine hcl powder, for solution

**Product Information****Item Code (Source)** NDC:58933-551**Route of Administration** ORAL**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 5 g
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 5 g
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 5 g

**Inactive Ingredients**

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>CARBOXYMETHYLCELLULOSE</b> (UNII: 05JZI7B19X)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>TRIBASIC CALCIUM PHOSPHATE</b> (UNII: 91D9GV0Z28)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	HONEY (LEMON)	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:58933-551-01	5 g in 1 PACKET; Type 0: Not a Combination Product	
---	------------------	--	--

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/20/2022	

## Part 2 of 2

### VICKS DAYQUIL HOT REMEDY RAPID RELIEF MULTI-SYMPATOM COLD AND FLU PLUS CONGESTION

acetaminophen, dextromethorphan hbr, phenylephrine hcl powder, for solution

## Product Information

Item Code (Source)	NDC:58933-550
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 5 g
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg in 5 g
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 5 g

## Inactive Ingredients

Ingredient Name	Strength
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>CARBOXYMETHYLCELLULOSE</b> (UNII: 05JZI7B19X)	
<b>TRIBASIC CALCIUM PHOSPHATE</b> (UNII: 91D9GV0Z28)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	

## Product Characteristics

Color	white	Score	
Shape		Size	

<b>Flavor</b>	HONEY (LEMON)	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-550-01	5 g in 1 PACKET; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/20/2022	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/20/2022	

**Labeler** - Procter & Gamble Manufactura S de RL de CV (812807550)

Revised: 10/2023

Procter & Gamble Manufactura S de RL de CV