

**BERKLEY AND JENSEN CLEARLAX- polyethylene glycol 3350 powder, for solution
BJWC**

BJWC ClearLax® Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - use once a day
 - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients

none

Questions or comments?

1-800-934-1204

Principal Display Panel

Compare to the active ingredient in MiraLAX®

CLEARLAX®

POLYETHYLENE GLYCOL 3350

POWDER FOR SOLUTION

OSMOTIC LAXATIVE

RELIEVES OCCASIONAL CONSTIPATION/IRREGULARITY

SOFTENS STOOL

GLUTEN FREE

17.9 OUNCES NET WT (510g)

30 ONCE-DAILY DOSES

UNFLAVORED POWDER

GRIT FREE

Compare to the active
ingredient in MiraLAX®*

berkley
jensen®

CLEARLAX®

POLYETHYLENE GLYCOL 3350
POWDER FOR SOLUTION
OSMOTIC LAXATIVE

- RELIEVES OCCASIONAL
CONSTIPATION/IRREGULARITY
- SOFTENS STOOL

17.9 OUNCES NET WT (510g)

30 ONCE-DAILY DOSES

: A5844 D3 F5



UNFLAVORED POWDER
GRIT FREE

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.

Dissolves in Any Beverage

Drug Facts

Active ingredient (in each dose) **Purpose**
 Polyethylene Glycol 3350,
 17 g (cap filled to line).....Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (continued)

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - use once a day
 - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° -25° C (68° -77° F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients none

Questions or comments?

1-800-934-1204

• Sugar Free

*This product is not manufactured or distributed by MSD Consumer Care, Inc., distributor of MiraLAX®.

Every Berkley Jensen product comes with our **100% Money-Back Guarantee.**

It's just one of the many ways a BJ's Membership gives you MORE.

Distributed by: BJ's Wholesale Club, 25 Research Drive, Westborough, MA 01581



: A5B44 D3 B5

BERKLEY AND JENSEN CLEARLAX

polyethylene glycol 3350 powder, for solution

Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:68391-306
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (POLYETHYLENE GLYCOL 3350)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68391-306-03	2 in 1 PACKAGE		
1		510 g in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090685	12/17/2009	

Labeler - BJWC (159082692)

Revised: 1/2015

BJWC