SINUFRIN NEILMED- oxymetazoline hydrochloride spray NeilMed Pharmaceuticals Inc

Drug Facts

Active Ingredients

0.05% Oxymetazoline Hydrochloride

Uses

temporarily relieves nasal congestion due to: common cold n sinusitis, hay fever, upper respiratory allergies

Warnings

Do not exceed recommended dosage. This product may cause temporary discomfort such as burning, stinging, sneezing, or an increase in nasal discharge. The use of this container by more than one person may spread infection.

Warnings

Do not use this product for more than 3 consecutive days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen. If symptoms persist, consult a doctor. Do not use this product if you have heart disease, high blood pressure, thyroid disease or diabetes unless directed by a doctor.

Directions for Dosing

Adults and children 6 to 12 years of age (with adult supervision): Per dose, 2 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period. Children under 6 years of age: consult a doctor.

Other Information

Store between 20 to 40 °C (68 to 104 °F) Retain carton for future reference on full labeling

Warnings

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Warnings

Do not use this product for more than 3 consecutive days. Use only as directed. Frequent or prolonged use may

cause nasal congestion to recur or worsen. If symptoms persist, consult a doctor. Do not use this product if you have heart disease, high

blood pressure, thyroid disease or diabetes unless directed by a doctor. Adult males with prostate disease, do not use this product.

Other Information

Keep head in an upright position for use. Do not tilt head backward while spraying. Wipe nozzle clean after each use.

Warnings

Stop using this product for more than 3 consecutive days. Stop and ask your doctor if symptoms of nasal congestion persist after using 3 consecutive days.

Warnings

If pregnant or breast-feeding, ask a health care professional before use.

Uses

temporarily relieves nasal congestion due to: common cold n sinusitis, hay fever, upper respiratory allergies.

Inactive Ingredients

Sodium chloride, sodium bicarbonate, propylene glycol, edetate disodium, benzalkonium chloride, purified water

Principal Display label

Drug Facts

Active ingredients

Purpose

Oxymetazoline hydrochloride 0.05% Nasal Decongestant Each full spray releases 0.1 mL of solution

Uses

temporarily relieves nasal congestion due to: common cold sinusitis hay fever upper respiratory allergies

1 Warnings

- Do not exceed recommended dosage.
- This product may cause temporary discomfort such as burning, stinging, sneezing, or an increase in nasal discharge.
- The use of this container by more than one person may spread infection.
- Do not use this product for more than 3 consecutive days.
 Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen. If symptoms persist, consult a doctor.
- Do not use this product if you have heart disease, high blood pressure, thyroid disease or diabetes unless directed by a doctor.
- Adult males with prostate disease, do not use this product.



Oxymetazoline HCl Nasal Solution Nasal Decongestant

Decongestant
Relief for up to 12 Hours

QUICK NASAL CONGESTION RELIEF

12 Hour Relief

0.5 fl oz (15 mL)

- If pregnant or breast feeding, ask a health professional before use
- Keep out of reach of children. If ingested, get medical help or contact a Poison Control Center right away.

(I) Directions for Dosing

- Adults and children 6 to 12 years of age (with adult supervision): Per dose, 2 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- Children under 6 years of age: consult a doctor

Other Information

- Store between 20 to 40 °C (68 to 104 °F)
- Retain carton for future reference on full labeling
- Keep head in an upright position for use.
- Do not tilt head backward while spraying.
- Wipe nozzle clean after each use.

inactive ingredients

Sodium chloride, sodium bicarbonate, propylene glycol, edetate disodium, benzalkonium chloride, purified water.


NDC 13709-231-07

Questions/Concerns 1 877 477 8633

Manufactured By NeilMed® Pharmaceuticals

601 Aviation Blvd, Santa Rosa, CA 95403 USA

Box label



Oxymetazoline Hydrochloride

Fast Acting

Nasal Decongestant

12 Hour Relief

Drug Facts

Active Ingredients Purpose
Oxymetazoline hydrochloride 0.05% Nasal Decongestant
Each full spray releases 0.1 mL of solution

Uses

- temporarily relieves nasal congestion due to:
- common cold sinusitis ■ hay fever
 - ■upper respiratory allergies

∆ Warnings

- Do not exceed recommended dosage.
 This product may cause temporary discomfort such as burning, stinging, sneezing, or an increase in nasal discharge.

 The use of this container by more than one person may
- spread infection.

 Do notuse this product for more than 3 consecutive days.
- Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen. If symptoms persist, consult a doctor.
- Do not use this product if you have heart disease, high blood pressure, thyroid disease or diabetes unless
- directed by a doctor.

 Adult males with prostate disease, do not use this product.

 If pregnant or breast-feeding, ask a health care
- professional before use.

 Keep out of reach of children. If swallowed, getmedical help or contact a Poison Control Center right away.

I Directions for Dosing

- Adults and children 6 to 12 years of age (with adult supervision): Per dose, 2 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- Children under 6 years of age: consult a doctor

Other Information

- Store between 20 to 40 °C (68 to 104 °F) Retain carton for future reference on full labeling
- Keep head in an upright position for use Do not tilt head backward while spraying.
- Wipe nozzle dean after each use

Inactive Ingredients

Sodium chloride, sodium bicarbonate, propylene glycol, edetate disodium, benzalkonium chloride, purified wate

Oxymetazoline HCl

Decongestant Relief for up to 12 Hours

NeilMed

SinuFrin

Decongestant 12 Hour Nasal Spray

QUICK NASAL CONGESTION

RELIEF

QUICK New! NASAL CONGESTION RELIEF

> **NeilMed®** SinuFrin-**Nasal Spray** starts to work in seconds, providing up to 12 hours of nasal decongestion.

No drowsiness 12 hour relief (Approximately)

Oxymetazoline HCl

Decongestant Relief for up to 12 Hours



QUICK NASAL CONGESTION RELIEF



QUICK NASAL CONGESTION RELIEF



USA - HEAD OFFICE / MFG
601 Aviation Blvd, Santa Rosa, CA 95403
Toll Free: 1(877) 477-8633
Tel: +1(707) 525-3784
Fax: +1(707) 525-3785
www.neilmed.com
CANADA
PO Box #3553, Markham, O N L3R 6G8
Toll Free: 1(877) 477-8633
Tel: +1(905) 480-0202
Fax: +1(905) 480-0201
UNITED KINGDOM

Fax: +1(905) 480-0201
UNITED KINGDOM
PO Box #2853, Coulsdon, Surrey, CR5 2WN, UK
Toll Free: +44 (0) 800 650 9447
Fax: +44 (0) 208 660 9447
Fax: +44 (0) 207 900 6414
WWW.neilmed.co.uk
IRELAND/EUROPE
PO Box #2853, Coulsdon, Surrey, CR5 2WN, UK
Toll Free: 00800 722 00027
Fax: +44 (0) 207 900 6414
AUSTRAIIA

rax: +44 (0)20/ 900 6414 AUSTRALIA 12/354 Chisholm Rd. Auburn, NSW 2144 Toll Free: 1800 158 900 Tel: +61 1300 652 148 Fax: +61 (03) 8677 9989 www.neilmed.com.au

www.neumed.com.au

NEW ZEALAND
PO Box #27310, Mount Roskill, Auckland 1440
Toll Free: 0800 634 5633
Tel: +64 (0)9 570 7279
Fax: +64 (0)9 353 1430

SINGAPORE
66 Tannery Lane, #01-02D
Sindo Industrial Building,
Singapore 347805
Tel: +65 (0)6 221 8948
Fax: +65 (0)6 491 5930

INDIA
Suite 1 C, #118, Neil Rao Towers,
Road #3 EPIP Zone,
Whitefield, Bangalore 560066
Tel: +91 80 4161 3512
Fax: +91 80 4160 5707



SINUFRIN NEILMED

oxymetazoline hydrochloride spray

Product Information

Product Type Route of Administration **HUMAN OTC DRUG**

Item Code (Source)

NDC:13709-231

NASAL

Active Ingredient/Active Moiety Ingredient Name OXYMETAZOLINE HYDROCHLORIDE (UNII: K89MJ0S5VY) (OXYMETAZOLINE HYDROCHLORIDE - UNII:8VLN5B44ZY) Basis of Strength OXYMETAZOLINE HYDROCHLORIDE 7.5 mg in 15 mL

Inactive Ingredients				
Ingredient Name	Strength			
SODIUM CHLORIDE (UNII: 451W47IQ8X)				
SODIUM BICARBONATE (UNII: 8MDF5V39QO)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)				
WATER (UNII: 059QF0KO0R)				

Product Characteristics			
Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:13709- 231-07	15 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)	07/15/2012		
2	NDC:13709- 231-04	15 mL in 1 BOX; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)	07/15/2012		
3	NDC:13709- 231-12	1 in 1 CARTON	04/21/2021		
3		30 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M012	07/15/2012			

Labeler - NeilMed Pharmaceuticals Inc (799295915)

Registrant - NeilMed Pharmaceuticals Inc (799295915)

Establishment					
Name	Address	ID/FEI	Business Operations		

NeilMed Pharmaceuticals Inc 799295915 manufacture(13709-231)

Revised: 1/2024 NeilMed Pharmaceuticals Inc