# EQUATE NAPROXEN SODIUM- naproxen sodium tablet, film coated Wal-Mart Stores Inc

-----

# Wal-Mart Naproxen Sodium Tablets 220 mg Drug Facts

# **Active ingredient (in each caplet)**

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

# **Purpose**

Pain reliever/fever reducer

### Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

### **Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

# Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

## When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

### If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### **Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older:	<ul> <li>take 1 caplet every 8 to 12 hours while symptoms last</li> <li>for the first dose you may take 2 caplets within the first hour</li> <li>do not exceed 2 caplets in any 8- to 12-hour period</li> <li>do not exceed 3 caplets in a 24-hour period</li> </ul>
Children under 12 years:	• ask a doctor

### Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

# **Inactive ingredients**

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

### Questions or comments?

1-888-287-1915

### **Principal Display Panel**

Compare to Aleve® Caplets Active Ingredient

2 75 COUNT BOTTLES INSIDE

TWIN PACK

Naproxen Sodium Tablets 220 mg

Pain Reliever/Fever Reducer (NSAID)

Strength to Last 12 Hours

220 mg

150 CAPLETS†

†Capsule Shaped Tablets

Actual Size



# equate® Naproxen **Sodium**

Tablets 220 mg

Pain Reliever/ **Fever Reducer** (NSAID)

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

equate<sup>®</sup>

NDC 49035-368-47

Compare to Aleve® Caplets **Active** Ingredient\*\*

equate®

Naproxen

Sodium

Tablets 220 mg

Pain Reliever/ Fever Reducer (NSAID)

Strength to Last 12 Hours

150 **CAPLETS**<sup>†</sup>

+Capsule Shaped Tablets



**Actual Size** 

equat

# Drug Facts

### Active ingredient (in each caplet)

Naproxen sodium 220 mg Pain reliever/ (naproxen 200 mg) (NSAID)\*..... fever reducer.

\*nonsteroidal anti-inflammatory drug

*Uses* ■ temporarily relieves minor aches and

pains due to:

- minor pain of arthritis backache
- muscular aches
- menstrual cramps
   headache
- the common cold
- toothache temporarily reduces fever

Purposes

### Warnings

Allergy alert Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing) shock
- skin reddening
- rash
   blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-287-1915.

Distributed by: Wal-Mart Stores, Inc., Bentonville, AR 72716. @ Wal-Mart Stores, Inc.

\*\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve® Caplets.



LOT NO.

EXP.

: A9414 2E C1



### Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Do not us

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers you have asthma

### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

### Drug Facts (continued)

### When using this product

■ take with food or milk if stomach upset occurs

■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Drug Facts (continued)

Directions ■ do not take more than directed

- the smallest effective dose should be used
- drink a full glass of water with each dose

  Adults and children 12 years and older:
- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period Children under 12 years: ask a doctor


#### Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-888-287-1915

KEEP CARTON FOR REFERENCE



# **EQUATE NAPROXEN SODIUM**

naproxen sodium tablet, film coated

### **Product Information**

1 Toutet Imol matter			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:49035-368
Route of Administration	ORAL	DEA Sche dule	

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
NAPRO XEN SO DIUM (NAPRO XEN)	NAPROXEN SODIUM	220 mg		

Inactive Ingredients			
Ingredient Name	Strength		
HYPROMELLOSES			
MAGNESIUM STEARATE			
CELLULO SE, MICRO CRYSTALLINE			
POLYETHYLENE GLYCOLS			
POVIDONES			
TALC			
TITANIUM DIO XIDE			

Product Characteristics			
Color	BLUE (Light Blue)	Score	no score
Shape	CAPSULE (oval)	Size	12mm
Flavor		Imprint Code	L368
Contains			

Packaging				
#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
1	NDC:49035-368-62	1 in 1 CARTON		
1		24 in 1 BOTTLE		
2	NDC:49035-368-78	1 in 1 CARTON		
2		100 in 1 BOTTLE		
3	NDC:49035-368-87	300 in 1 BOTTLE		
4	NDC:49035-368-47	2 in 1 CARTON		
4	NDC:49035-368-98	75 in 1 BOTTLE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	04/27/2007	

# Labeler - Wal-Mart Stores Inc (051957769)

Revised: 10/2013 Wal-Mart Stores Inc