MUSCLE PAIN RELIEF- acetaminophen tablet Rite Aid Corporation

Drug Facts

Active ingredient (in each caplet)
Acetaminophen USP, 650 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - backache
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps
 - headache
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks ever day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek a medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- adults
 - take 2 caplets every 8 hours with water
 - o swallow whole; do not crush, chew, split or dissolve
 - do not take more than 6 caplets in 24 hours
 - o do not use for more than 10 days unless directed by a doctor
- under 18 years of age: ask a doctor

Other information

store between 20-25°C (68-77°F)

Inactive ingredients

hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient of Tylenol® 8 HR Muscle Aches & Paint

SEE NEW WARNING

8 HR MUSCLE PAIN RELIEF

ACETAMINOPHEN

EXTENDED-RELEASE TABLETS USP 650 mg

PAIN RELIEVER/FEVER REDUCER

For the temporary relief of minor muscle aches & pain

Contain no aspirin

CAPLETS

*CAPSULE-SHAPED BI-LAYER TABLETS

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® 8 HR Arthritis Pain.

TAMPER EVIDENT: DO NOT USE IF FOIL INNER SEAL IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 www.riteaid.com

Product Label

Exp. Date Lot No.:

PLD-A684A FC007224

Orug Facts (continued)

■ store between 20-25°C (68-77°F) Other information

■ if you are allergic to acetaminophen or any of

Drug Facts (continued)

the inactive ingredients in this product

povidone, pregelatinized starch, sodium starch microcrystalline cellulose, polyethylene glycol cellulose, hypromellose, magnesium stearate Inactive ingredients hydroxyethy

Ask a doctor or pharmacist before use if you are Ask a doctor before use if you have liver disease.

taking the blood thinning drug warfarin.

1-877-753-3935 Monday-Friday 9AM-5PM EST Questions or comments? Call

alycolate, stearic acid

This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® 8 HR Muscle

RITE AID, 30 HUNTER LANE CAMP HILL, PA 1701 DISTRIBUTED BY: www.riteaid.com

attention is critical for adults as well as for children

medical help or contact a Poison Control Center

Overdose warning: In case of overdose, get

Keep out of reach of children.

professional before use.

If pregnant or breast-feeding, ask a health

These could be signs of a serious condition.

redness or swelling is present

new symptoms occur

(1-800-222-1222) right away. Quick medical

even if you do not notice any signs or symptoms.

do not take more than directed (see Overdose

swallow whole; do not crush, chew, split, or do not take more than 6 caplets in 24 hours

take 2 caplets every 8 hours with water

do not use for more than 10 days unless

children under 12 years: do not use

directed by a doctor

adults and children 12 years of age and over

warning)

SATISFACTION

TAMPER EVIDENT: DO NOT USE IF FOIL INNER SEAL IS BROKEN OR MISSING KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

50 CAPLETS*

Active ingredient 'in each caplet

Drug Facts

Purpose

Pain reliever fever reduces Acetaminophen USP, 650 mg.

 temporarily relieves minor aches and pains due to toothache minor pain of arthritis muscular aches

the common cold premenstrual and menstrual cramps headache

temporarily reduces fever

acetaminophen. Severe liver damage may occur if Liver warning: This product contains

SEE NEW WARNING

EXTENDED-RELEASE TABLETS USP 650 mg PAIN RELIEVER/FEVER REDUCER For the temporary relief of minor muscle aches & pair

Contains no aspirin

■ more than 6 caplets in 24 hours, which is the vou take:

3 or more alcoholic drinks every day while using with other drugs containing acetaminophen maximum daily amount

Compare to the active ingredient in Tylenol® 8 HR Muscle Aches & Paint

Allergy alert. Acetaminophen may cause severe this product

f a skin reaction occurs, stop use and seek medical skin reactions. Symptoms may include: skin reddening blisters rash help right away.

ACTUAL SIZE

*CAPSULE-SHAPED BI-LAYER TABLETS

appily refund your money

Do not use

(prescription or nonprescription). If you are not with any other drug containing acetaminophen sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

RITE **AID** 8 **HR Muscle Pain Relief**

Stop use and ask a doctor if

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

MUSCLE PAIN RELIEF

acetaminophen tablet

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:11822-6019

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) ACETAMINOPHEN 650 mg

Inactive Ingredients Ingredient Name Strength

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

HYDROXYETHYL CELLULOSE (140 MPA.S AT 5%) (UNII: 8136Y38GY5)

MAGNESIUM STEARATE (UNII: 70097M6I30)

STARCH, CORN (UNII: O8232NY3SJ)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

POVIDONE (UNII: FZ 989GH94E)

Contains

SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)

STEARIC ACID (UNII: 4ELV7Z65AP)

Product CharacteristicsColorwhiteScoreno scoreShapeCAPSULESize19mmFlavorImprint CodeG650

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:11822- 6019-2	1 in 1 BOX	07/30/2020	12/31/2025		
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product				
2	NDC:11822- 6019-1	1 in 1 BOX	07/30/2020	12/31/2025		
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product				

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211544	07/30/2020	12/31/2025

Labeler - Rite Aid Corporation (014578892)

Revised: 2/2023 Rite Aid Corporation